Case report

GM2 activator protein deficiency, mimic of Tay-Sachs disease

Sheena P. Kochumon^a, Dhanya Yesodharan^a, KP Vinayan^b, Natasha Radhakrishnan^c,
Jayesh J. Sheth^d, Sheela Nampoothiri^{a,*}

^a Department of Pediatric Genetics, Amrita Institute of Medical Sciences & Research Center, Aims Ponekkara PO, Kochi, 682041, Kerala, India

^b Division of Pediatric Neurology, Department of Neurology, Amrita Institute of Medical Sciences & Research Center, Aims Ponekkara PO Kochi 682041, India

^c Department of Ophthalmology, Amrita Institute of Medical Sciences & Research Center, Aims Ponekkara PO, Kochi, 682041, Kerala, India

^d Department of Biochemical and Molecular Genetics, FRIGE's Institute of Human Genetics, FRIGE House, Satellite, Ahmedabad, India

ARTICLE INFO

Article history:

Received 18 March 2017

Received in revised form 5 August 2017

Accepted 9 August 2017

Available online 14 August 2017

Keywords:

GM2 Gangliosidosis

GM2-Activator protein

GM2A

Cherry red spots

Tay-Sachs disease

ABSTRACT

GM2 Gangliosidosis is a group of autosomal recessive genetic disorders caused by intra-lysosomal deposition of ganglioside GM2 mainly in the neuronal cells. GM2-Activator protein deficiency is an extremely rare type of GM2 gangliosidosis (AB variant) caused by the mutation of *GM2A*. We report a case of a female child who presented with clinical features similar to classical Tay-Sachs disease, but with normal beta hexosaminidase enzyme levels. Molecular study revealed a novel homozygous intronic mutation which confirmed the diagnosis of GM2 Activator protein deficiency. GM2 Activator protein deficiency is a mimic of Classical Tay-Sachs disease and should be a differential diagnosis in children who present with neuroregression, cherry red spots without hepatosplenomegaly and with normal beta hexosaminidase enzyme levels.

© 2017 Published by Elsevier, a division of RELX India, Pvt. Ltd on behalf of Indian Epilepsy Society.

1. Introduction

GM2-Activator protein deficiency (OMIM 272750) is an extremely rare neuroregressive disorder which is clinically indistinguishable from classical Tay-Sachs disease. Konrad Sandhoff coined it as GM2 gangliosidosis AB variant in 1971. GM2 gangliosidosis AB variant is an autosomal recessive genetic disorder caused by the mutation of *GM2A* encoding for GM2 Activator protein.^{1,2} Only 16 patients with GM2 Activator protein deficiency have been reported so far and the underlying mutation was proven in only 9 cases.^{3–11} We report a homozygous intronic mutation proven case of GM2 activator protein deficiency which is the 10th case report proven by molecular studies.

2. Case report

A nineteen month old female child was referred to the Department of Pediatric Genetics for evaluation of neuroregression. She was the second child born to third degree consanguineous parents. She was born after an uneventful antenatal and post natal period with the birth weight of 3500 gm. Her development was age appropriate till seventh month. She had already attained head control and could sit without support. At seventh month, her parents had noticed squint following which they noticed regression of previously acquired motor and mental milestones. She started developing excessive startle response from 14 months of age and also had two episodes of seizures at 14th and 18th months.

On examination, she was found to be floppy with complete head lag and was not fixing on objects. Excessive startle response was elicited and there was no organomegaly. Fundus examination revealed pale disc with bilateral classical cherry red spots. MRI Brain done at 11 months of age was normal. Patient underwent three EEGs during the whole course of her illness. The first EEG which was done at 16th months of age was reported as a normal sleep record, without any awake record. The second EEG at 18th months of age did report generalized long interval periodic complexes with correlation with the myoclonic jerks. The third

* Corresponding author.

E-mail addresses: sheenspk@gmail.com (S.P. Kochumon), dhanyayesodharan@aims.amrita.edu (D. Yesodharan), vinayankp@aims.amrita.edu (K. Vinayan), natashar@aims.amrita.edu (N. Radhakrishnan), sheeladr@gmail.com (S. Nampoothiri).

Fig. 1. Electropherogram showing a homozygous mutation in the intron 3 of *GM2A* (c.243-2A > T) in the proband and heterozygous status in the parents confirming the *GM2* activator protein deficiency.

EEG at 20th months of age showed only diffuse beta activity, probably related to drugs. There were no periodic complexes in this last EEG. Her seven year old elder sibling was normal.

In view of positive consanguinity, neuroregression, bilateral cherry red spots, exaggerated startle response and absence of organomegaly, possibility of Tay – Sachs disease was considered. Enzyme analysis was sent for confirmation of the diagnosis. The total hexosaminidase, hexosaminidase A and chitotriosidase estimations were within normal limits. The other storage disorders presenting with neuroregression and cherry red spots viz, Gauchers disease, Niemann pick and sialidosis were ruled out by enzymatic analysis.

Since the clinical phenotypes of infantile form of Tay- Sachs disease is indistinguishable from *GM2* activator protein deficiency, we investigated in favour of the latter by molecular analysis. Mutation study of the *GM2A* revealed homozygous mutation in the intron 3 of *GM2A* (c.243-2 A > T) and her parents are heterozygous carriers for the same mutation, thereby confirming the diagnosis of *GM2* activator protein deficiency in the proband (Fig. 1).

3. Discussion

GM2 Gangliosidoses are a group of lysosomal storage disorder caused by excessive intra – lysosomal deposition of *GM2* ganglioside mainly in the neuronal cells. There are 3 types of *GM2* Gangliosidoses. The most common type is Tay – Sachs disease (variant B) caused by the mutation of the *HEXA* which is associated with the deficiency of β hexosaminidase A (Hex – A) and normal levels of β hexosaminidase B (Hex – B) enzyme activity. Sandhoff disease (variant O) is caused by the mutation of *HEXB* with deficient activity of β subunit of Hex – A and Hex – B. *GM2*

activator protein deficiency (variant AB) is an extremely rare type due to defective *GM2*-Activator protein resulting from the mutation of the *GM2A*.¹

Hexosaminidase is a dimer composed of 2 subunits. The alpha sub unit is encoded by the *HEXA* gene located on chromosome 15. The beta sub unit is encoded by the *HEXB* gene located on chromosome 5. Hex – A is a heterodimer of α β (alpha beta) and Hex – B is a homodimer of β (beta beta). The *GM2A* is located on chromosome 5q31.3–q33.1 and has 4 exons.^{1,12} *GM2* activator protein (*GM2* – AP) is 22 kDa glycoprotein which is an essential cofactor for β hexosaminidase A for the conversion of *GM2* to *GM3*. The *GM2* activator protein is encoded by the *GM2A*. Mutation in *HEXA*, *HEXB* or *GM2A* can lead to classical Tay- Sachs disease, Sandhoff disease and *GM2* activator protein deficiency respectively. The *GM2* activator initially binds with *GM2* gangliosides and this complex binds with β hexosaminidase A, thereby leading to the degradation of *GM2* gangliosides. Hence the defective *GM2* Activator protein is unable to produce a functional ganglioside *GM2* Activator complex even with normal or elevated levels of Hexosaminidase A and B.^{1,13} The binding of *GM2* activator to a wide variety of negatively charged glycosphingolipids may indicate that the activator protein has functions other than assisting the enzymatic hydrolysis of *GM2*.¹⁴ Failure of enzymatic hydrolysis results in the progressive intra – lysosomal accumulation of *GM2* gangliosides in the neuronal cells and spinal cord leading to severe progressive psychomotor regression. Affected infants are normal at birth and later present with clinical features consistent with classical Tay – Sachs disease with psychomotor regression, exaggerated startle response and refractory seizures. Exact incidence of AB variant is not known. It is an extremely rare disease and only 16 cases have been reported so far, of which 9 cases only are proven by molecular studies (Table 1).^{3–11}

Table 1Comparison of present case with previous proven cases.^{3–11}

	Schroder et al., 1991, Xie et al., 1992 ^{3,4}	Schroder et al., 1993 ⁵	Schepers et al., 1996 ⁶	Schepers et al., 1996 ⁶	Chen et al., 1999 ⁷	Kolodny et al., 2008 ⁸	Renaud et al., 2015 ⁹	Kustermann et al., 2015 ¹⁰	Sheth et al., 2016 ¹¹	Our Case
Ethnicity	Afro American	Indian	Saudi Arab	Spanish	Laotian Hmong	Indian	Hmong	Turkish	Indian	Indian
Consanguinity	NA	—	+	+	—	—	—	+	+	+
Sex	F	F	F	F	M	F	F	F	M	F
Age of onset (months)	9	5	8	7	5	11	3	9	12	7
Presenting Symptoms	Neuro regression	Neuro regression	Motor Weakness, head lag	Neuro regression	Delayed motor milestones, weakness	Development delay, seizures	Poor visual fixation, global developmental delay	Ataxia, Developmental stagnation	Global developmental delay	Neuro regression
Squint	NA	NA	NA	NA	NA	NA	NA	NA	NA	+
Nystagmus	NA	+	+	+	NA	NA	NA	NA	+	+
Bilateral Cherry red spots	+	+	+	+	+	+	+	+	+	+
Hepatomegaly	NA	NA	+	NA	NA	NA	NA	+	—	—
Splenomegaly	NA	NA	NA	NA	NA	NA	NA	NA	—	—
Seizures	NA	+	+	+	+	+	+	—	NA	+
Hyperacusis	+	+	+	+	+	+	+	+	+	+
EEG	NA	NA	NA	NA	NA	NA	NA	NA	NA	Generalized periodic complexes
MRI brain	NA	NA	Diffuse Brain atrophy	Demyelination of both cerebral hemispheres & cerebellum	Abnormal signal in basal ganglia and white matter	NA	Delayed myelination with abnormal signal intensity in both thalami	periventricular dysmyelination	Abnormal signal intensity in putamen and thalamus	Normal
Hex-A & Hex-B Enzyme	Normal	Normal	Normal	Normal	Normal	Normal	Normal	Normal	Normal	Normal
Mutation in GM2A gene (homozygous)	c.412T > C (p. C107R) ³ c.412T > C (p. C138R) ⁴	c.506G > C (p.R169P)	c.262_264delAA (p.88Kdel)	c.410delA (p. H137PfsX34)	c.160G > T (p.E54X)	c.522T > G (p. L174R)	c. 160G > T (p.E54X)	c.164C > T (p. P55L)	c.472G > T (p.E158X)	c.243-2 A > T (IVS3-2 A > T)

NA – Not Available, F –Female, M – Male.

4. Conclusion

The diagnosis of GM2 Activator protein deficiency should be considered in infants who present with classical features of Tay-Sachs disease with normal levels of beta hexosaminidase A enzyme activity. Confirmation of the diagnosis is absolutely essential for genetic counselling and prenatal diagnosis, as the risk of recurrence in subsequent pregnancy is 25%.

Contributor's credits

SK prepared the manuscript & did detailed literature search. JS did the molecular studies and interpretation. DY did literature search and had helped in manuscript preparation. VP had evaluated the patient contributed in manuscript. NR had evaluated the case in detail and had helped in manuscript. SN had conceived the idea of drafting this paper and has done the final drafting and will act as the guarantor of the manuscript.

Conflict of interest

None stated.

Source of funding

None.

References

1. Gravel RA, Clarke JTR, Kaback MM, et al. The GM2 gangliosidosis. In: Scriver CR, Beaudet AL, Sly WS, Valle D, eds. *The Metabolic and Molecular Bases of Inherited Disease*. 7th ed. New York: McGraw-Hill; 1995:2839–2879.
2. Sakuraba H, Itoh K, Shimmoto M, et al. GM2 gangliosidosis AB variant. Clinical and biochemical studies of a Japanese patient. *Neurology*. 1999;52:372–377.
3. Schroder M, Schnabel D, Suzuki K, et al. A mutation in the gene of a glycolipid binding protein (GM2activator) that causes GM2 gangliosidosis variant AB. *FEBS Lett*. 1991;290:1–3.
4. Xie B, Wang W, Mahuran DJ. A cys138-to-Arg substitution in the GM2 activator protein is associated with the AB variant form of GM2 gangliosidosis. *Am J Hum Genet*. 1992;50:1046–1052.
5. Schroder M, Schnabel D, Hurwitz R, et al. Molecular genetics of GM2 gangliosidosis AB variant: a novel mutation and expression in BHK cells. *Hum Genet*. 1993;92:437–440.
6. Schepers U, Glombitza G, Lemm T, et al. Molecular analysis of a GM2-Activator deficiency in two patients with GM2-Gangliosidosis AB variant. *Am J Hum Genet*. 1996;59:1048–1056.
7. Chen B, Rigat B, Curry C, et al. Structure of the GM2A gene: identification of an exon 2 nonsense mutation and a naturally occurring transcript with an In-frame deletion of exon 2. *Am J of Hum Genet*. 1999;65:77–87.
8. Kolodny E, Sathe S, Zeng BJ, et al. A novel GM2- activator deficiency mutation as a cause of AB variant GM2-Gangliosidosis. *Mol Genet Metab*. 2008;93:27–28.
9. Renaud D, Brodsky M. GM2-Gangliosidosis, AB variant: clinical, ophthalmological, MRI and molecular findings. *JIMD Rep*. 2016;25:83–86.
10. Kustermann W, Brackmann F, Gusek-Schneider G, et al. Rare variant of GM2 gangliosidosis due to activator protein deficiency: a case report. *Neuropediatrics*. 2015;46:PS01–PS21.
11. Sheth J, Datar C, Mistri M, et al. GM2 gangliosidosis AB variant: novel mutation from India —a case report with a Review. *BMC Pediatr*. 2016;16:88.
12. Burg J, Conzelmann E, Sandhoff K, et al. Mapping of the gene coding for the human GM2 activator protein to chromosome 5. *Ann Hum Genet*. 1985;49:41–45.
13. Wright CS, Li SC, Rastinejad F. Crystal structure of human GM2-activator protein with a novel beta-cup topology. *J Mol Biol*. 2000;304:411–422.
14. Hama Y, Li Y, Li SC. Interaction of GM2 activator protein with glycosphingolipids. *J Biol Chem*. 1997;272:2828–2833.