Pancreatotomy-guided laser lithotripsy in a patient with difficult ductal stone

► **Fig. 1** Magnetic resonance cholangiopancreatography, showing an abrupt stop at the pancreatic body, followed by major dilation of the remaining body and tail.

► **Fig. 2** Endoscopic ultrasound revealed a large intraductal stone in the pancreatic body (7.4 mm), and a dilated Wirsung in the body and tail.

► **Fig. 3** Pancreatography showing irregularity in the proximal body, suggesting a large intraductal stone.

► **Fig. 4** Pancreatotomy image showing a large intraductal stone impacted on a main duct stricture.

A 63-year-old man with heavy alcohol consumption was referred to our institution for upper abdominal pain, weight loss, and a computed tomography scan showing signs of chronic pancreatitis (parenchymal calcifications and atrophy of the pancreatic body/tail).

A magnetic resonance cholangiopancreatography was performed, showing Wirsung dilation, namely of the tail, and an abrupt stop in the pancreatic body of unknown cause (► **Fig. 1**). Endoscopic ultrasound revealed an intraductal stone in the pancreatic body (7.4 mm), and a dilated Wirsung in the body and tail (► **Fig. 2**). The patient underwent endoscopic retrograde cholangiopancreatography (ERCP).

Pancreatography showed an irregular Wirsung contour in the head and irregularity in the body–tail transition, suggestive of an intraductal stone (► **Fig. 3**).

Pancreatic sphincterotomy was performed, and the calculus was crossed with the guidewire, but it was impossible to cross it with a 6-mm dilation balloon. After hydrostatic balloon dilation of the pancreatic head (up to 6 mm), a pancreatoscope (Spyglass Direct Visualization System; Boston Scientific, Marlborough, Massachusetts, USA) was advanced over a 0.025-inch guidewire to reach a large intraductal stone of 7–8 mm in size (► **Fig. 4**, ► **Video 1**). After targeting the stone, laser bursts (Holmium laser, Auriga XL; Boston Scientific) of less than 5 seconds were delivered through the aqueous medium using a 365-μm diameter fiber (energy level 1200 mJ; frequency of 12 Hz). After stone fragmentation, ductal clearance was achieved with an 8.5-mm extraction balloon. Two pancreatic stents (12 cm, 7 Fr) were placed.

▶ Video 1 Intraductal pancreatoscopy with holmium laser lithotripsy was performed until complete stone fragmentation was achieved.

▶ Fig. 5 Pancreatogram image showing resolution of the pancreatic stricture after successful endotherapy.

At follow-up 3 months later, repeat ERCP showed frank improvement of the head stricture, without filling defects in the remaining Wirsung (▶ Fig. 5). The patient remained asymptomatic during follow-up (6 months) without further interventions.

Published experience is limited, but pancreatoscopy-guided laser lithotripsy for calcific chronic pancreatitis is a promising technique that can be used as a supplementary approach to extracorporeal shock wave lithotripsy or as a single modality in a small number of stones obstructing the main pancreatic duct [1–4].

Endoscopy_UCTN_Code_TTT_1AR_2AI

Competing interests

Jorge Canena is a consultant for Boston Scientific but did not receive any financial arrangements for this research or any assistance with manuscript preparation.

The authors

Gonçalo Alexandrino, Luís Lourenço, Catarina G. Rodrigues, David Horta, Jorge Reis, Jorge Canena

Department of Gastroenterology, Hospital Prof. Doutor Fernando Fonseca, Amadora, Portugal

Corresponding author

Gonçalo Alexandrino, MD

Department of Gastroenterology, Hospital Prof. Doutor Fernando Fonseca, IC 19

Amadora 2720-276, Portugal

Fax: +351-21-4345566

goncaloalexandrino@gmail.com

References

- [1] Attwell AR, Patel S, Kahaleh M et al. ERCP with per-oral pancreatoscopy-guided laser lithotripsy for calcific chronic pancreatitis: a multicenter U.S. experience. *Gastrointest Endosc* 2015; 82: 311–318
- [2] Shah J. Innovations in intraductal endoscopy: cholangioscopy and pancreatoscopy. *Gastrointest Endosc Clin N Am* 2015; 25: 779–792
- [3] Ang TL. Chronic pancreatitis with pancreatic duct stricture and calculi treated by fully covered self-expandable metal stent placement and intraductal pancreatoscopy-guided laser lithotripsy. *Endoscopy* 2017; 49: E145–E146
- [4] Maydeo A, Kwek BE, Bhandari S et al. Single-operator cholangioscopy-guided laser lithotripsy in patients with difficult biliary and pancreatic ductal stones (with videos). *Gastrointest Endosc* 2011; 74: 1308–1314

Bibliography

DOI <https://doi.org/10.1055/a-0574-2278>

Published online: 8.3.2018

Endoscopy 2018; 50: E130–E131

© Georg Thieme Verlag KG

Stuttgart · New York

ISSN 0013-726X

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>

CORRECTION

Alexandrino G, Lourenço L, Rodrigues CG et al. Pancreatoscopy-guided laser lithotripsy in a patient with difficult ductal stone.

Endoscopy 2018, doi:10.1055/a-0574-2278

In the above mentioned article one author was missing in the authors' list.

Correct is: Gonçalo Alexandrino, Luís Lourenço, Catarina G. Rodrigues, David Horta, Jorge Reis, Jorge Canena.

This was corrected in the online version on April 6, 2018.