Closure of a mucosal defect with clips and rubber band: a technical trick to improve edge apposition in large mucosal defects

Endoscopic mucosal resection (EMR) of duodenal adenomas is recommended by European Society for Gastrointestinal Endoscopy (ESGE) guidelines as offering complete resection with relatively low morbidity compared with surgery or endoscopic submucosal dissection (ESD) [1, 2]. Nevertheless, resection of duodenal adenomas carries a high risk of perforation (5%–10%) and delayed bleeding (15%) [3]. Systematic closure of mucosal defects could reduce the risk of delayed bleeding by protecting the resected area from bile and pancreatic secretions. Nevertheless, apposition of the two mucosal edges is not always feasible and can carry a risk of muscular damage from the clip itself.

We report here the case of a 63-year-old patient who was referred for endoscopic removal of a large (4-cm) duodenal adenoma with high grade dysplasia (► **Fig. 1**). We performed complete piecemeal EMR. To reduce the risk of delayed bleeding, we decided to close the defect with hemoclips. Because of the large area resected, it appeared impossible to achieve complete apposition of the two edges of the mucosal defect by standard clipping. As used for traction in colorectal ESD [4, 5], we added a small rubber band to the first clip, which was fixed on the inner edge of the lesion. A second clip was used to grasp the rubber band and was then attached to the opposite edge of the lesion (► **Fig. 2**). Thanks to the retraction provided by the rubber band, the two edges of the lesion came closer and it was subsequently possible to close all of the mucosal defect by adding two further clips onto the retracted area (► **Video 1**). The patient was discharged 48 hours after the intervention, with no adverse events having occurred.

Therefore, the technique of traction with a rubber band allows the two lateral

► **Fig. 1** Endoscopic views showing a duodenal adenoma under: **a** white light; **b** narrow-band imaging (NBI).

► **Video 1** Closure of a large mucosal defect with clips and rubber band.

edges of the resected area to be apposed, thereby facilitating complete closure. This appears to be a safe and effective technique for closing large mucosal defects after EMR.

Endoscopy_UCTN_Code_TTT_1AO_2AG

Competing interests

None

► **Fig. 2** Schematic of the procedure. **a** Mucosectomy of a large duodenal nonampullary sporadic adenoma is performed using a front-viewing endoscope. **b** The first clip with the attached rubber band is fixed onto the anterior side of the mucosal defect. **c** Elastic traction is applied using a second clip in order to close together the two opposing mucosal edges. **d** Further clips are then applied to completely close the mucosal defect.

The authors

Alexandru Lupu^{1,2}, Jérémie Jacques³, Jérôme Rivory¹, Florian Rostain¹, Romain Legros³, Thierry Ponchon^{1,4}, Mathieu Pioche^{1,4}

- 1 Department of Endoscopy and Gastroenterology, Pavillon L, Edouard Herriot Hospital, Lyon, France
- 2 Department of Endoscopy and Gastroenterology, Fundeni clinical institute, Bucharest, Romania.
- 3 Department of Endoscopy and Gastroenterology, Limoges university Hospital, Limoges, France
- 4 Inserm U1032 LabTau, Lyon, France

Corresponding author

Mathieu Pioche, MD

Endoscopy unit – Digestive Disease department, Pavillon L – Edouard Herriot Hospital, 69437 Lyon, France
mathieu.pioche@chu-lyon.fr

References

- [1] Klein A, Ahlenstiel G, Tate DJ et al. Endoscopic resection of large duodenal and papillary lateral spreading lesions is clinically and economically advantageous compared with surgery. *Endoscopy* 2017; 49: 659 – 667
- [2] Marques J, Baldaque-Silva F, Pereira P et al. Endoscopic mucosal resection and endoscopic submucosal dissection in the treatment of sporadic nonampullary duodenal adenomatous polyps. *World J Gastrointest Endosc* 2015; 7: 720 – 727
- [3] Lépilliez V, Chemaly M, Ponchon T et al. Endoscopic resection of sporadic duodenal adenomas: an efficient technique with a substantial risk of delayed bleeding. *Endoscopy* 2008; 40: 806 – 810
- [4] Utzeri E, Jacques J, Charissoux A et al. Traction strategy with clips and rubber band allows complete en bloc endoscopic submucosal dissection of laterally spreading tumors invading the appendix. *Endoscopy* 2017; 49: 820 – 822
- [5] Jacques J, Legros R, Charissoux A et al. A combination of pocket, double-clip countertraction, and isolated HybridKnife as a quick and safe strategy for colonic endoscopic submucosal dissection. *Endoscopy* 2017; 49: E134 – E135

Bibliography

DOI <https://doi.org/10.1055/a-0591-2039>
Published online: 13.4.2018
Endoscopy 2018; 50: 726–727
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

ENDOSCOPY E-VIDEOS
<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>