Rendezvous biliary recanalization with combined percutaneous transhepatic cholangioscopy and double-balloon endoscopy

► **Fig. 1** Percutaneous transhepatic cholangiogram showing complete obstruction of the choledochojejunal anastomosis in a patient who had undergone pancreaticoduodenectomy 7 years previously.

Despite advances in biliary stenting in patients with altered gastrointestinal anatomy, it is still a challenging procedure [1]. We present a case where percutaneous transhepatic cholangioscopy (PTCS) was combined with double-balloon endoscopy (DBE) for biliary stenting in a patient with complete obstruction of a choledochojejunostomy.

A 71-year-old woman, who had a history of distal cholangiocarcinoma and had undergone pancreaticoduodenectomy 7 years previously, experienced recurrent cholangitis. DBE-assisted balloon dilation had been performed 7 months previously for stricture of the choledochojejunal anastomosis. However, she developed complete obstruction of the anastomosis (► **Fig. 1**). A 7.2-Fr percutaneous transhepatic biliary drainage (PTBD) catheter was initially placed, and the fistula tract was dilated up to 12 Fr within 4 weeks. DBE-assisted endoscopic retrograde cholangiopancreatography was then attempted. First, the double-balloon endoscope (EI-580BT; Fujifilm, Tokyo, Japan) was advanced to the afferent limb, and a percutaneous transhepatic cholangiogram revealed complete obstruction of

► **Fig. 2** Left panel: The choledochojejunal anastomosis has an appearance similar to an ulcer scar. Right panel: Transillumination from the percutaneous transhepatic cholangioscope guides direct precutting using the double-balloon endoscope.

► **Video 1** Biliary recanalization, using a rendezvous technique with combined percutaneous transhepatic cholangioscopy and double-balloon endoscopy, for a completely obstructed choledochojejunostomy.

the anastomosis. Next, a PTCS scope (BF type P260F; Olympus, Tokyo, Japan) was inserted via the PTBD route. However, a guidewire (0.018-inch, Pathfinder Exchange; Boston Scientific Japan, Tokyo, Japan) through the PTCS scope could not pass the anastomosis (► **Video 1**). Therefore, we attempted direct precutting (KD-V451M; Olympus) at the anas-

tomosis, using the double-balloon endoscope and guided by transillumination from the percutaneous transhepatic cholangioscope' (► **Fig. 2**, ► **Video 1**). A small incision was carefully made in order to create a fistula (► **Fig. 3**). This was followed by successful passage of the guidewire (0.032-inch, Radifocus Guidewire M; Terumo, Tokyo, Japan)

► **Fig. 3** Radiograph showing direct precutting at the choledochojejunal anastomosis, under fluoroscopic guidance and transillumination from the percutaneous transhepatic cholangioscope. Left inset: enteroscopy view. Right inset: percutaneous transhepatic cholangioscopy view.

► **Fig. 4** Radiograph showing the guidewire passing through the obstruction. Inset: percutaneous transhepatic cholangioscopy view.

► **Fig. 5** The guidewire is grasped by means of a snare under fluoroscopic and percutaneous transhepatic cholangioscopic guidance.

► **Fig. 6** Insertion of the percutaneous transhepatic biliary drainage catheter, using a rendezvous technique, across the previously obstructed choledochojejunal anastomosis. Inset: percutaneous transhepatic cholangioscopic view.

completely through the anastomotic obstruction (► **Fig. 4**, ► **Video 1**). We then grasped the guidewire with an ultraslim basket catheter (Zero Tip Retrieval Basket; Boston Scientific) using the cholangioscope (► **Fig. 5**, ► **Video 1**). Finally, a 12-Fr PTBD catheter was placed across the obstruction without any complications (► **Fig. 6**, ► **Video 1**).

The rendezvous technique in combination with PTCS and DBE facilitates biliary recanalization of complete biliary obstruction [1,2]. However, blind incision has the risk of gastrointestinal tract per-

foration or bile leakage. Although caution should be exercised, incision guided by transillumination from the peroral transhepatic cholangioscope is a safe and less invasive technique compared with surgery.

Endoscopy_UCTN_Code_TTT_1AR_2AJ

Competing interests

None

The authors

Hiroshi Kawakami^{1,2}, Teshin Ban^{1,2}, Yoshimasa Kubota^{1,2}, Shinya Ashizuka², Ichiro Sannomiya², Naoya Imamura^{2,3}, Takeomi Hamada^{2,3}

- 1 Department of Gastroenterology and Hepatology, Faculty of Medicine, University of Miyazaki, Miyazaki, Japan
- 2 Department of Gastroenterology and Hepatology, Center for Digestive Disease and Division of Endoscopy, University of Miyazaki Hospital, Miyazaki, Japan
- 3 Division of Hepato-Biliary-Pancreas Surgery, Department of Surgery, Faculty of Medicine, University of Miyazaki, Miyazaki, Japan

Corresponding author

Hiroshi Kawakami, MD, PhD

Department of Gastroenterology and
Hepatology, Center for Digestive Disease
and Division of Endoscopy, University of
Miyazaki Hospital, 5200, Kiyotake, Kihara,
Miyazaki 889-1692, Japan
Fax: +81-985-859802
hiropo@med.miyazaki-u.ac.jp

References

- [1] Shimatani M, Takaoka M, Ikeura T et al. Rendezvous technique: double-balloon endoscopy and SpyGlass direct visualization system in a patient with severe stenosis of a choledochojejunal anastomosis. *Endoscopy* 2014; 46 (Suppl. 01): E275 – E276
- [2] Bukhari MA, Haito-Chavez Y, Ngamruengphong S et al. Rendezvous biliary recanalization of complete biliary obstruction with direct peroral and percutaneous transhepatic cholangioscopy. *Gastroenterology* 2018; 154: 23 – 25

Bibliography

DOI <https://doi.org/10.1055/a-0591-2109>

Published online: 13.4.2018

Endoscopy 2018; 50: E146–E148

© Georg Thieme Verlag KG

Stuttgart · New York

ISSN 0013-726X

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>