Reintervention for stent occlusion after endoscopic ultrasound-guided hepaticogastrostomy with novel use of a precut needle-knife

Endoscopic ultrasound-guided hepaticogastrostomy (EUS-HGS) has gained popularity as an alternative biliary drainage method [1, 2]; however, reintervention after EUS-HGS remains to be elucidated. In EUS-HGS, use of a biliary stent that is longer than 100 mm is recommended in order to prevent stent migration [2, 3]. However, such stent placement occasionally makes reintervention challenging owing to the long length of the stent in the gastric lumen. A few reports have described technical efforts involved in reintervention after EUS-HGS [4, 5]. We describe a patient who underwent successful reintervention via a novel use of a precut needle-knife.

A 74-year-old woman with recurrent pancreatic cancer after pancreaticoduodenectomy presented with recurrent cholangitis. An 8 × 100 mm covered metal stent (Niti-S biliary covered stent; TaeWoong Medical, Seoul, South Korea) had been previously deployed during EUS-HGS for biliary obstruction at the hepatic hilum. Stent occlusion occurred 4 months after EUS-HGS. Abdominal computed tomography showed a dilated intrahepatic bile duct, and stent occlusion was confirmed on endoscopy (► Fig. 1). Revisionary stent placement was attempted.

First, the advancement of an endoscopic retrograde cholangiopancreatography (ERCP) catheter was attempted via the proximal end of the HGS stent; however, the long stent length in the gastric lumen rendered catheter insertion impossible. Therefore, reintervention through the stent mesh was attempted. A 0.035-inch guidewire (Jagwire; Boston Scientific, Marlborough, Massachusetts, USA) was successfully passed through the stent mesh (► Fig. 2); however, an ERCP catheter could not be passed. Subsequently, a diathermic dilator was utilized, but it failed to break the stent mesh. Next, the use of a precut needle-knife (NeedleCut3V; Olympus, Tokyo, Japan) was

► **Fig. 1** Stent occlusion after endoscopic ultrasound-guided hepaticogastrostomy. **a** Abdominal computed tomography showed a dilated intrahepatic bile duct. **b** Gastrosocopy showed an occluded hepaticogastrostomy stent.

► **Fig. 2** A 0.035-inch guidewire (Jagwire; Boston Scientific, Marlborough, Massachusetts, USA) was passed successfully through the mesh of the previously deployed hepaticogastrostomy stent (Niti-S biliary covered stent, 8 × 100 mm; TaeWoong Medical, Seoul, South Korea).

considered. Using this knife, the stent mesh was broken easily (► Fig. 3), and a 7-Fr plastic stent (Flexima; Boston Scientific) was successfully deployed via the stent mesh into the left intrahepatic bile duct (► Fig. 4, ► Video 1). Cholangitis resolved in a few days.

The use of a precut needle-knife is simple and may be considered as a useful treatment option for reintervention after EUS-HGS.

Endoscopy_UCTN_Code_CPL_1AL_2AD

Competing interests

None

The authors

Kosuke Minaga, Mamoru Takenaka, Ayana Okamoto, Shunsuke Omoto, Takeshi Miyata, Hajime Imai, Masatoshi Kudo

Department of Gastroenterology and Hepatology, Kindai University Faculty of Medicine, Osaka-Sayama, Japan

► **Fig. 3** A precut needle-knife (NeedleCut3V; Olympus, Tokyo, Japan) was inserted over the guidewire and could break the stent mesh easily.

► **Fig. 4** A 7-Fr biliary plastic stent (70 mm long, Flexima; Boston Scientific, Marlborough, Massachusetts, USA) was deployed successfully via the stent mesh into the left intrahepatic bile duct.

► **Video 1** Using a precut needle-knife, the mesh of the previously deployed hepaticogastrostomy stent was broken easily. Thereafter, a 7-Fr biliary plastic stent was deployed successfully via the stent mesh into the left intrahepatic bile duct.

Corresponding author

Mamoru Takenaka, MD

Department of Gastroenterology and Hepatology, Kindai University Faculty of Medicine, 377-2 Ohno-Higashi, Osaka-Sayama 589-8511, Japan
Fax: +81-72-3672880
mamoxyo45@gmail.com

References

- [1] Wang K, Zhu J, Xing L et al. Assessment of efficacy and safety of EUS-guided biliary drainage: a systematic review. *Gastrointest Endosc* 2016; 83: 1218–1227
- [2] Nakai Y, Isayama H, Yamamoto N et al. Safety and effectiveness of a long, partially covered metal stent for endoscopic ultrasound-guided hepaticogastrostomy in patients with malignant biliary obstruction. *Endoscopy* 2016; 48: 1125–1128
- [3] Okuno N, Hara K, Mizuno N et al. Stent migration into the peritoneal cavity following endoscopic ultrasound-guided hepaticogastrostomy. *Endoscopy* 2015; 47 (Suppl. 01): E311
- [4] Ogura T, Masuda D, Takeuchi T et al. Simplified reintervention method of EUS-guided hepaticogastrostomy stent obstruction. *Gastrointest Endosc* 2016; 83: 831
- [5] Minaga K, Takenaka M, Miyata T et al. Through-the-mesh technique after endoscopic ultrasonography-guided hepaticogastrostomy: a novel re-intervention method. *Endoscopy* 2016; 48: E369–E370

Bibliography

DOI <https://doi.org/10.1055/a-0596-7171>
Published online: 13.4.2018
Endoscopy 2018; 50: E153–E154
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>