


## Resection of large sessile serrated polyps by cold piecemeal endoscopic mucosal resection: Serrated COLD Piecemeal Endoscopic mucosal resection (SCOPE)



**Fig. 1** Resection of a large sessile serrated polyp by cold piecemeal endoscopic mucosal resection (SCOPE) technique. **a** A 40-mm sessile serrated adenoma/polyp in the ascending colon; the edges are enhanced by chromoendoscopy. **b** Complete resection was achieved using the SCOPE technique. **c** No recurrence was observed at follow-up colonoscopy. **d** Normal mucosa in continuity with a serrated polyp highlights complete resection of the lesion (B1, normal mucosa; B2, serrated polyp).

Sessile serrated adenomas/polyps (SSA/Ps) are frequently found in the proximal colon, where the wall is thinner and easily damaged by diathermy during polypectomy, which also carries a risk of delayed bleeding, perforation, and post-polypec-

tomy syndrome. SSA/Ps are often flat with subtle, irregular edges making endoscopic assessment of their extent difficult [1]. This can lead to incomplete resection and risk of post-colonoscopy cancer [2].

Currently, cold snare resection (CSR) is considered the preferred technique to resect small polyps. It is safe, time efficient, and user friendly [3]. Recently, case series have highlighted the safety and efficiency of CSR for larger adeno-

► **Table 1** Patient and polyp characteristics.

Case #	Age, years	Location	Number of SSA/P	Size, mm (number of polyps)	Follow-up, months	Outcome
1	64	Ascending colon	1	30	12	No recurrence
2	62	Hepatic flexure	1	30	9	No recurrence
3	68	Ascending colon	1	30	7	5-mm residual polyp; cold snared
4	68	Ascending colon	1	20	7	No recurrence
5	42	Hepatic flexure	1	30	6	No recurrence
6	31	Cecum – transverse colon	7	10 (5), 15 (1), 20 (1)	12	No recurrence
7	39	Cecum – transverse colon	7	10 (4), 20 (3)	6	No recurrence
8	77	Ascending colon	2	12 (1), 18 (1)	8	No recurrence
9	34	Cecum – transverse colon	4	10 (2), 15 (2)	12	No recurrence
10	29	Ascending colon	4	10 (2), 15 (2)	7	No recurrence

SSA/P, sessile serrated adenoma/polyp.

► **Video 1** A 30-mm sessile serrated polyp was resected using the serrated cold piecemeal endoscopic mucosal resection (SCOPE) technique.

The SCOPE technique was applied successfully in 10 consecutive patients with 29 large SSA/Ps. We achieved complete resection in all cases (► **Table 1**). Minor oozing was noted in almost all cases; however, no hemostatic interventions were required. There were no adverse events during or after resection. Histology showed complete resection of polyps (► **Fig. 1**). In one polyp (3.4%), a small area of residual tissue was observed at the follow-up examination; this was resected using cold snaring.

Endoscopy\_UCTN\_Code\_TTT\_1AQ\_2AD

### Competing interests

None

### The authors

**Rajaratnam Rameshshanker<sup>\*</sup>, Zacharias Tsiamoulos<sup>\*</sup>, Andrew Latchford, Morgan Moorghen, Brian P. Saunders**

Wolfson Unit for Endoscopy, St. Mark's Hospital and Academic Institute, Harrow, United Kingdom

\* These authors contributed equally to this work.

mas [4,5]. In this series, we report our preliminary experience in achieving complete resection of large SSA/Ps using a cold piecemeal endoscopic mucosal resection (SCOPE) technique. Following detection of an SSA/P, the polyp surface was assessed. The polyp was then lifted using a submucosal injection of 0.1% hyaluronate and methylene blue, and resected using a small cold snare (9mm, Exacto; US Endoscopy, Mentor, Ohio, USA) in a piecemeal man-

ner (► **Video 1**). A gradual increase in snare closure pressure was applied to mechanically transect each polyp piece. Each polyp was resected with a small rim of adjacent normal mucosa (1–2 mm) in order to achieve a complete resection margin. The polypectomy defect edges were scrutinized for any remaining polyp and trimmed using the snare, or cold avulsed with a biopsy forceps (► **Fig. 1**, ► **Video 1**).

## Corresponding author

### Rajaratnam Rameshshanker, MD

Wolfson Unit for Endoscopy, St. Mark's Hospital and Academic Institute, Watford road, Harrow HA1 3UJ, United Kingdom  
Fax: +44-208-2354033  
r.rameshshanker@nhs.net

## References

- [1] Kahi CJ, Li X, Eckert GJ et al. High colonoscopic prevalence of proximal colon serrated polyps in average-risk men and women. *Gastrointest Endosc* 2012; 75: 515–520
- [2] Zauber AG, Winawer SJ, O'Brien MJ et al. Colonoscopic polypectomy and long-term prevention of colorectal-cancer deaths. *N Engl J Med* 2012; 366: 687–696

- [3] Hewett DG. Cold snare polypectomy: optimizing technique and technology (with videos). *Gastrointest Endosc* 2015; 82: 693–696
- [4] Piraka C, Saeed A, Waljee AK et al. Cold snare polypectomy for non-pedunculated colon polyps greater than 1 cm. *Endosc Int Open* 2017; 5: E184–E189
- [5] Muniraj T, Sahakian A, Ciarleglio MM et al. Cold snare polypectomy for large sessile colonic polyps: a single-center experience. *Gastroenterol Res Pract* 2015; 2015: 175959

### Bibliography

DOI <https://doi.org/10.1055/a-0599-0346>  
Published online: 9.5.2018  
*Endoscopy* 2018; 50: E165–E167  
© Georg Thieme Verlag KG  
Stuttgart · New York  
ISSN 0013-726X

### ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



*Endoscopy E-Videos* is a free access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>