Narrow-band imaging detects synchronous oropharyngeal squamous carcinoma during treatment of an early esophageal squamous cell carcinoma

► **Fig. 1** Esophageal epidermoid lesion. **a, c** Appearance with white-light imaging. **b, d** Appearance with narrow-band imaging.

Squamous cell carcinoma (SCC) can appear synchronously in the upper aerodigestive tract and in the esophagus by the mechanism of field cancerization, as tobacco and alcohol are both major risk factors for these lesions [1,2]. Narrow-band imaging (NBI) and Lugol chromoendoscopy are the standard techniques for assessment of superficial esophageal lesions, although Lugol staining can lead to adverse events, such as mucosal irritation, allergic reactions, and aspiration, especially in the oropharyngeal area. NBI has proven to be superior to white-light imaging (WLI), with significantly higher diagnostic sensitivity in the detection of superficial mucosal lesions of the esophagus and pharynx [3], by showing altered intrapapillary capillary loop (IPCL) pattern [4].

► **Fig. 2** Resected specimen and endoscopic appearance after resection.

► **Fig. 3** Narrow-band imaging of lesions in pharynx, larynx, palate, and tonsil.

► **Video 1** Synchronous oropharyngeal squamous carcinoma detected by narrow-band imaging during treatment of an early esophageal squamous cell carcinoma.

These lesions appeared typical with large IPCLs of various shapes (B1 type). Biopsies were taken from all lesions. The histology result from the resected esophageal specimen was in situ SCC without micro-invasion and R0 resection. Palate and tonsil biopsies also showed in situ SCC.

When an esophageal SCC lesion is found, the endoscopist should do a complete examination of the oropharyngeal area in order to detect synchronous lesions, which can be multiple in patients with long tobacco and alcohol use. NBI chromoendoscopy is safe, simple, and reproducible with no risk of aspiration, and should be used as a first-intention imaging technique.

Endoscopy_UCTN_Code_CCL_1AB_2pAB

Competing interests

Dr. Katzarov received a travel grant from the European Society of Gastrointestinal Endoscopy to join our team at Edouard Herriot Hospital for 4 months.

We present here the case of a 58-year-old man who was referred for endoscopic evaluation of the esophagus.

Endoscopic examination found a reddish 2.5-cm anfractuous lesion (► **Fig. 1 a**) on the front wall of the proximal esophagus. NBI showed IPCL type B1 (► **Fig. 1 b**). The margins were delineated and endoscopic

submucosal dissection (ESD) was performed, applying the tunnel and clip line technique (► **Fig. 2**) [5]. When withdrawing the endoscope using NBI, multiple SCC lesions were found in the upper aerodigestive tract and oral cavity (► **Video 1**): on palate, right tonsil, pharynx, and left piriform sinus (► **Fig. 3**).

The authors

**Alexander Katzarov^{1,2}, Alexandru Lupu¹,
Jérémie Jacques³, Jérôme Rivory¹,
Florian Rostain¹, Thomas Walter^{1,4},
Mathieu Pioche^{1,5}**

- 1 Department of Endoscopy and Gastroenterology, Pavillon L, Edouard Herriot Hospital, Lyon, France
- 2 Department of Gastroenterology and Hepatology, Military Medical Academy, Sofia, Bulgaria
- 3 Department of Endoscopy and Gastroenterology, Limoges University Hospital, Limoges, France
- 4 Digestive Oncology division, Pavillon E, Edouard Herriot Hospital, Lyon, France
- 5 Inserm U1032 LabTau, Lyon, France

Corresponding author

Mathieu Pioche, MD
Endoscopy Unit – Digestive Disease
Department, Pavilion L – Edouard Herriot
Hospital, 69437 Lyon Cedex, France
Fax: +33-4-72110147
mathieu.pioche@chu-lyon.fr

References

- [1] Ni XG, Zhang QQ, Zhu JQ et al. Hypopharyngeal cancer associated with synchronous oesophageal cancer: risk factors and benefits of image-enhanced endoscopic screening. *J Laryngol Otol* 2018; 132: 154–161
- [2] Jaiswal G, Jaiswal S, Kumar R et al. Field cancerisation: concept and clinical implications in head and neck squamous cell carcinoma. *J Exp Ther Oncol* 2013; 10: 209–214
- [3] Muto M, Minashi K, Yano T et al. Early detection of superficial squamous cell carcinoma in the head and neck region and esophagus by narrow band imaging: a multicenter randomized controlled trial. *J Clin Oncol* 2010; 28: 1566–1572
- [4] Oyama T, Inoue H, Arima M et al. Prediction of the invasion depth of superficial squamous cell carcinoma based on micro vessel morphology: magnifying endoscopic classification of the Japan Esophageal Society. *Esophagus* 2017; 14: 105–112
- [5] Jacques J, Legros R, Rivory J et al. The “tunnel + clip” strategy standardized and facilitates oesophageal ESD procedures: a prospective, consecutive bi-centric study. *Surg Endosc* 2017; 31: 4838–4847

Bibliography

DOI <https://doi.org/10.1055/a-0640-2464>
Published online: 3.7.2018
Endoscopy 2018; 50: E264–E266
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>