Thoracoscopic and endoscopic cooperative surgery (TECS): a novel less invasive technique for resection of gastric tube cancer after esophagectomy

► **Fig. 1** Endoscopic view of a depressed lesion (arrows) with an irregular surface observed in Patient #1.

► **Fig. 2** Endoscopic ultrasonographic image of a low-echoic tumor with deep invasion of the submucosal layer (arrows).

► **Fig. 3** Endoscopic view of a slightly depressed lesion (arrows) observed in Patient #2.

With recent improvements in the survival of patients after esophagectomy, the occurrence of secondary malignancies arising in the gastric tube has been increasing [1,2]. However, resection of the reconstructed gastric tube with lymphadenectomy for gastric tube cancer (GTC) is an invasive procedure associated with high morbidity and mortality [3]. Local resection without lymphadenectomy may be reasonable in high risk patients. We have developed a novel thoracoscopic and endoscopic cooperative surgery (TECS) technique as a minimally invasive alternative.

Two patients underwent TECS for GTC after esophagectomy. Patient #1 had a 25mm, undifferentiated, submucosal invasive adenocarcinoma (► **Fig. 1**, ► **Fig. 2**), which was considered difficult to resect completely by endoscopic submucosal dissection (ESD). Patient #2 had undergone ESD for a 10mm depressed GTC (► **Fig. 3**); however, histopathological analysis revealed noncurative resection because of submucosal invasion to 1150 μ m, with a positive vertical margin. The absence of metastasis was confirmed preoperatively using computed tomography. The TECS steps (► **Video 1**) were: 1) exposure of the gastric tube by

► **Video 1** Thoracoscopic and endoscopic cooperative surgery procedure in Patient #2 with gastric tube cancer after noncurative endoscopic submucosal dissection.

thoracoscopy; 2) mucosal incision around the involved site and submucosal trimming using ESD technique, followed by full-thickness incision by endoscopy (ITknife 2; Olympus, Tokyo, Japan); 3) thoracoscopic resection with Harmonic scalpel (Ethicon, Somerville, New Jersey, USA); 4) suturing.

Both TECS procedures were performed under general anesthesia with orotra-

cheal intubation and were successfully completed in 250 and 420 minutes, respectively. Blood loss was minimal. No perioperative complications such as anastomotic leaks, bleeding, or stricture were observed. Oral intake was started on postoperative days 7 and 8. The patients were discharged 14 and 25 days after the procedure.

ESD for GTC is sometimes technically difficult with fibrosis [4], and the long-term outcomes for noncurative patients are reportedly less satisfactory [5]. Although careful follow-up for possible metastasis is necessary in our patients, TECS was a feasible, safer, and less invasive therapeutic option for patients with noncurative GTC.

Endoscopy_UCTN_Code_TTT_1AT_2AF

Competing interests

None

The authors

Yoshiki Tsujii¹, Makoto Yamasaki², Yoshito Hayashi¹, Koji Tanaka², Tomoki Makino², Yuichiro Doki², Tetsuo Takehara¹

- 1 Department of Gastroenterology and Hepatology, Osaka University Graduate School of Medicine, Suita, Japan
- 2 Department of Gastroenterological Surgery, Osaka University Graduate School of Medicine, Suita, Japan

Corresponding author

Tetsuo Takehara, MD, PhD

Department of Gastroenterology and Hepatology, Osaka University Graduate School of Medicine, 2-2 Yamadaoka, Suita, Osaka 565-0871, Japan
Fax: +81-6-68793629
takehara@gh.med.osaka-u.ac.jp

References

- [1] Matsubara T, Yamada K, Nakagawa A. Risk of second primary malignancy after esophagectomy for squamous cell carcinoma of the thoracic esophagus. *J Clin Oncol* 2003; 21: 4336–4341
- [2] Okamoto N, Ozawa S, Kitagawa Y et al. Metachronous gastric carcinoma from a gastric tube after radical surgery for esophageal carcinoma. *Ann Thorac Surg* 2004; 77: 1189–1192
- [3] Sugiura T, Kato H, Tachimori Y et al. Second primary carcinoma in the gastric tube constructed as an esophageal substitute after esophagectomy. *J Am Coll Surg* 2002; 194: 578–583
- [4] Nishide N, Ono H, Kakushima N et al. Clinical outcomes of endoscopic submucosal dissection for early gastric cancer in remnant stomach or gastric tube. *Endoscopy* 2012; 44: 577–583
- [5] Nonaka S, Oda I, Sato C et al. Endoscopic submucosal dissection for gastric tube cancer after esophagectomy. *Gastrointest Endosc* 2014; 79: 260–270

Bibliography

DOI <https://doi.org/10.1055/a-0751-2540>
Published online: 7.11.2018
Endoscopy 2019; 51: E5–E6
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>