



## Cervical esophageal adenocarcinoma arising from heterotopic gastric mucosa, treated with endoscopic submucosal dissection



► **Fig. 1** Cervical esophageal adenocarcinoma arising from heterotopic gastric mucosa (HGM). **a** Conventional endoscopy revealed circumferential HGM of the cervical esophagus and a 20-mm protruding lesion in the posterior wall. **b** Indigo carmine spraying revealed a flat lesion in the reddish mucosa, but the demarcation line was unclear. **c** Narrow band imaging (NBI) revealed a clearly demarcated brownish area consistent with the reddish area (arrowheads). **d** Magnifying NBI revealed villous and glandular structures of various sizes and a dense distribution of abnormal capillaries.



► **Fig. 2** Histologically, tumor cells showed well-differentiated adenocarcinoma in the muscularis mucosa.

showed adenocarcinoma, with no evidence of deep submucosal invasion. Therefore, ESD was performed using a dual knife (KD-655L; Olympus, Tokyo, Japan) and an ITknife nano (KD-612; Olympus, Tokyo, Japan) (► **Video 1**). The resected specimen measured 51 × 35 mm and contained a macroscopically measured 32 × 27-mm tumor surrounded by HGM. Histologically, the tumor cells showed well-differentiated adenocarcinoma in the muscularis mucosa (► **Fig. 2**). Immunohistochemically, the tumor cells were positive for mucin (MUC) 2, MUC5AC, and MUC6. The estimated incidence of HGM of the esophagus is 0.75%–11% based on endoscopic studies [3,4]. Primary adenocarcinoma arising from HMG is rare and several cases of mucosal cancer have been reported [1,2,5]. The macroscopic appearance was protruding or polypoid in most cases [5]; therefore, adenocarcinoma should be suspected when a protrusion is identified in HGM in the esophagus.


Endoscopy\_UCTN\_Code\_TTT\_1AO\_2AG

### Competing interests

None

Esophageal adenocarcinoma arising from heterotopic gastric mucosa (HGM) is extremely rare [1,2]. We describe a case of esophageal adenocarcinoma arising from HGM in the cervical esophagus that was treated with endoscopic submucosal dissection (ESD). A 58-year-old man found to have an esophageal tumor on endoscopy was referred to our hospital for further examination and treatment. Conventional endoscopy revealed circumferential HGM of the cervical esophagus and a 20-mm

protruding lesion in the posterior wall. Poorly defined reddish mucosa surrounded the protruding lesion (► **Fig. 1 a**). Spraying with indigo carmine visualized the flat lesion in the reddish mucosa, but the demarcation line was unclear (► **Fig. 1 b**). Narrow-band imaging (NBI) revealed a clearly demarcated brownish area consistent with the reddish area at 18–21 cm from the upper incisors, with villous and glandular structures of various sizes and dense distribution of abnormal capillaries (► **Fig. 1 c, d**). Biopsy



**Video 1** Endoscopic submucosal dissection of cervical esophageal adenocarcinoma arising from heterotopic gastric mucosa.

## The authors

**Yasuhiro Oono<sup>1,2</sup>, Shinmura Kensuke<sup>1</sup>, Yusuke Yoda<sup>1</sup>, Keisuke Hori<sup>1</sup>, Hiroaki Ikematsu<sup>1</sup>, Tomonori Yano<sup>1</sup>**

- 1 Department of Gastroenterology and Endoscopy, National Cancer Center Hospital East, Chiba, Japan
- 2 Department of Internal Medicine, Tokyo Metropolitan Ebara Hospital, Tokyo, Japan

## Corresponding author

**Yasuhiro Oono, MD**

Department of Gastroenterology and Endoscopy, National Cancer Center Hospital East, 6-5-1, Kashiwanoha, Kashiwa, Chiba 277-8577, Japan  
Fax: +81-4-71346928  
yohno@east.ncc.go.jp

## References

- [1] Noguchi T, Takeno S, Takahashi Y et al. Primary adenocarcinoma of the cervical esophagus arising from heterotopic gastric mucosa. *J Gastroenterol* 2001; 36: 704–709
- [2] Yasar B, Tarcin O, Benek D et al. Intramucosal adenocarcinoma arising from ectopic gastric mucosa in the upper esophagus treated successfully with endoscopic mucosal resection. *J Gastrointest Cancer* 2014; 45 (Suppl. 01): 201–204
- [3] Weickert U, Wolf A, Schroder C et al. Frequency, histopathological findings, and clinical significance of cervical heterotopic gastric mucosa (gastric inlet patch): a prospective study in 300 patients. *Dis Esophagus* 2011; 24: 63–68
- [4] Yu L, Yang Y, Cui L et al. Heterotopic gastric mucosa of the gastrointestinal tract: prevalence, histological features, and clinical characteristics. *Scand J Gastroenterol* 2014; 49: 138–144
- [5] Kadota T, Fujii S, Oono Y et al. Adenocarcinoma arising from heterotopic gastric mucosa in the cervical esophagus and upper thoracic esophagus: two case reports and literature review. *Expert Rev Gastroenterol Hepatol* 2016; 10: 405–414

## Bibliography

DOI <https://doi.org/10.1055/a-0767-6253>  
Published online: 23.11.2018  
Endoscopy 2019; 51: E28–E29  
© Georg Thieme Verlag KG  
Stuttgart · New York  
ISSN 0013-726X

## ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



*Endoscopy E-Videos* is a free access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at  
<https://mc.manuscriptcentral.com/e-videos>