Rendezvous within biloma technique combining percutaneous and endoscopic approaches: a novel biliary recanalization method

► **Fig. 1** Abdominal computed tomography showing a large biloma under the diaphragm.

► **Fig. 2** Contrast injection from the percutaneous route showing the right intrahepatic B6 bile duct in communication with the biloma.

Biliary injury and biloma are complications that can occur with hepatobiliary surgery [1]. Endoscopic or percutaneous recanalization for postoperative biliary injury have been reported as minimally invasive treatments [2], but remains challenging for severe biliary stricture [3–5]. Here, we present a case of biliary injury successfully treated by combining endoscopic and percutaneous approaches within the biloma.

A 61-year-old man who underwent partial hepatectomy for hepatocellular carcinoma

► **Fig. 3** Cholangiographic views during endoscopic retrograde cholangiopancreatography showing: **a** the disrupted bile duct of the right posterior branch; **b** a 0.025-inch guidewire that could be successfully advanced into the biloma through the stricture but could not be passed through the disrupted B6.

presented with high fever and dyspnea. Computed tomography (CT) revealed a large biloma under the diaphragm (► **Fig. 1**), and percutaneous drainage was performed. Contrast agent injection via the percutaneous route showed a communication between the biloma and the right intrahepatic bile duct (► **Fig. 2**). Biliary recanalization was attempted with endoscopic retrograde cholangiopancreatography (ERCP). Cholangiography showed a disrupted right posterior branch bile duct (► **Fig. 3**). A 0.025-inch guidewire was advanced into the biloma through the stricture but could not be passed through the disrupted B6 bile duct. Percutaneous transhepatic biliary drainage (PTBD) of B6 was performed.

Via the PTBD route, the guidewire could reach the biloma, but not the right hepatic duct. Therefore, rendezvous biliary recanalization within the biloma combining endoscopic and percutaneous approaches was attempted. After coiling the guidewire within the biloma under ERCP guidance, an EN snare (Merit Medi-

cal, West Jordan, Utah, USA) designed with three interlaced loops to retrieve objects was inserted via the PTBD route (► **Fig. 4**). The guidewire was grasped with the EN snare, guided through B6, and drawn out via the PTBD route. Finally, a 6-Fr plastic stent was placed through the biloma with its tip positioned at B6 (► **Fig. 5**; ► **Video 1**). The PTBD catheter was removed 3 days post-procedure and the patient was discharged without any complications.

This “rendezvous within biloma” technique could be a minimally invasive rescue option when either the percutaneous or endoscopic approach alone is difficult.

Endoscopy_UCTN_Code_TTT_1AR_2AJ

Competing interests

None

► **Fig. 4** The EN snare, which had been inserted via the percutaneous transhepatic biliary drainage route, grasping the guidewire, which had been coiled within the biloma under endoscopic retrograde cholangiopancreatography guidance: **a** on fluoroscopic view; **b** in schematic view. **c** Photograph of the EN snare showing its three interlaced loops that can be used to retrieve and manipulate foreign objects in the body. Source ► **Fig. 4c**: Merit Medical Systems, INC

► **Fig. 5** The 6-Fr plastic stent that had been placed through the biloma with its tip positioned at B6: **a** on fluoroscopic view; **b** in schematic view.

References

- [1] Lo CM, Fan ST, Liu CL et al. Biliary complications after hepatic resection: risk factors, management, and outcome. *Arch Surg* 1998; 133: 156–161
- [2] Yabe S, Kato H, Mizukawa S et al. Predictive factors for outcomes of patients undergoing endoscopic therapy for bile leak after hepatobiliary surgery. *Dig Endosc* 2017; 29: 353–361
- [3] Jang SI, Rhee K, Kim H et al. Recanalization of refractory benign biliary stricture using magnetic compression anastomosis. *Endoscopy* 2014; 46: 70–74
- [4] Kawakami H, Abo D, Kawakubo K et al. Rendezvous biliary recanalization combining percutaneous and endoscopic techniques using a diathermic dilator for bile duct obstruction. *Endoscopy* 2014; 46: E460–E461
- [5] Tonozuka R, Mukai S, Tsuchiya T et al. Recanalization after biliojejunostomy by use of a new digital per-oral cholangioscope through the hepaticogastrostomy route. *VideoGIE* 2016; 1: 63–65

The authors

Ayana Okamoto¹, Kosuke Minaga¹, Mamoru Takenaka¹, Tomoe Yoshikawa¹, Toshimitsu Iwasaki², Masakatsu Tsurusaki³, Masatoshi Kudo¹

- 1 Department of Gastroenterology and Hepatology, Kindai University Faculty of Medicine, Osaka-Sayama, Japan
- 2 Department of Surgery, Kindai University Faculty of Medicine, Osaka-Sayama, Japan
- 3 Department of Radiology, Kindai University Faculty of Medicine, Osaka-Sayama, Japan

Corresponding author

Kosuke Minaga, MD

Department of Gastroenterology and Hepatology, Kindai University Faculty of Medicine, 377-2 Ohno-Higashi, Osaka-Sayama, 589-8511, Japan
Fax: +81-72-3672880
kousukeminaga@med.kindai.ac.jp

The guidewire which was coiled within the biloma via the ERCP route was grasped with the snare.

Video 1 The “rendezvous within biloma” technique combining percutaneous and endoscopic approaches was used to treat post-operative biliary injury with biloma. This technique could be a minimally invasive rescue option in patients where either the percutaneous or endoscopic approach alone is difficult.

Bibliography

DOI <https://doi.org/10.1055/a-0800-8216>

Published online: 14.12.2018

Endoscopy 2019; 51: E42–E44

© Georg Thieme Verlag KG

Stuttgart · New York

ISSN 0013-726X

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at <https://mc.manuscriptcentral.com/e-videos>