Successful endoscopic submucosal dissection of a large cavernous hemangioma in the colon

Endoscopic submucosal dissection (ESD) is regarded as a common treatment for complete resection of early gastrointestinal neoplasms [1]. However, few cases have been reported on ESD for resection of cavernous hemangioma in the digestive tract. Here we present a patient with a globular, pedunculated, cavernous hemangioma in the descending colon that was removed successfully en bloc by ESD without any bleeding (► **Video 1**).

A 50-year-old woman was referred to our hospital for melena. Initial colonoscopy examination revealed a submucosal tumor approximately 20 mm in diameter in the descending colon, mainly characterized by a soft, globular, pedunculated submucosal lesion with a red-purple nodular surface (► **Fig. 1 a, b**). Further endoscopic ultrasonography indicated that the lesion originated from the submucosa. The lesion showed high echogenicity, mixed with a small anechogenic area on the inside, and with a decreased blood flow signal (► **Fig. 1 c**).

For treatment, we first performed endoscopic incision of the colonic mucosa using a Hybrid knife (Erbe, Tübingen, Germany) after submucosal injection (► **Fig. 2 a**). The submucosal dissection was very

carefully performed to avoid damage to the body of the cavernous hemangioma and a clear field was maintained. The lesion was then successfully removed from the colon wall by ESD, as described previously (► **Fig. 2 b, c, e**). We used three clips (two from Anrei Medical, Hangzhou, China and one from Micro-Tech, Nanjing, China) for closure of the mucosal defect (► **Fig. 2 d**). Finally, histological examina-

tion of the resected specimen confirmed cavernous hemangioma (► **Fig. 2 f**). The patient was discharged with no further symptoms after 2 days of intravenous antibiotic treatment.

In summary, we believe that this is the first published report of a colonic cavernous hemangioma that has been completely removed by ESD.

► **Video 1** A large cavernous hemangioma in the colon was successfully removed by endoscopic submucosal dissection.

► **Fig. 1** Colonoscopy and endoscopic ultrasonography examination of the submucosal lesion. **a, b** A submucosal tumor approximately 20 mm in diameter characterized by a soft, globular, pedunculated submucosal lesion with a red-purple nodular surface. **c** Endoscopic ultrasonography indicated that the lesion originated from the submucosa and showed high echogenicity, mixed with a small anechogenic area on the inside, and with decreased blood flow signal (yellow arrow).

► **Fig. 2** Colonic cavernous hemangioma was completely removed by endoscopic submucosal dissection (ESD). **a** An endoscopic incision of the mucosa was performed after submucosal injection. **b, c, e** The lesion was successfully removed from the colon wall by ESD. **d** Closure of the mucosal defect. **f** Histological examination of the resected specimen.

Endoscopy_UCTN_Code_CCL_1AD_2AF

Competing interests

None

The authors

Kan Chen^{1,*}, Min Yan^{2,*}, Feng Liu¹

- 1 Department of Gastroenterology, Shanghai Tenth People's Hospital, Tongji University School of Medicine, Shanghai, China
- 2 Department of Pathology, Shanghai Tenth People's Hospital, Tongji University School of Medicine, Shanghai, China

* These authors contributed equally to this work.

Corresponding author

Feng Liu, MD

Department of Gastroenterology, Shanghai Tenth People's Hospital, Tongji University School of Medicine, Middle Yanchang Road No. 301, Jingan District, Shanghai 200072, China
Fax: +86-21-66300588
drliuffeng@hotmail.com

Reference

- [1] Mavrogenis G, Hochberger J, Deprez P et al. Technological review on endoscopic submucosal dissection: available equipment, recent developments and emerging techniques. *Scand J Gastroenterol* 2017; 52: 486–498

Bibliography

DOI <https://doi.org/10.1055/a-0875-3519>
Published online: 12.4.2019
Endoscopy 2019; 51: E195–E196
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X