Traction-assisted endoscopic submucosal dissection for a gastric lesion involving the pyloric ring and duodenal bulb

An 81-year-old woman underwent an upper gastrointestinal endoscopy because of anemia. Two lesions were detected: a 30-mm type 0-Ip located in the lesser curvature of the antrum, and a 25-mm type 0-Is involving the upper hemisphere of the pylorus and the proximal duodenal bulb. Biopsy specimens revealed hyperplastic polyp and adenocarcinoma, respectively. Endoscopic submucosal dissection (ESD) was proposed (► **Video 1**). The procedure was performed using an insulated-tipped knife (IT-knife 2; Olympus, Tokyo, Japan), with the patient under deep sedation. First, a submucosal injection on the distal part was performed and initial approach to the lesion from the duodenal bulb was attempted; however, neither the forward nor retroflexion view allowed a good endoscopic approach to the lesion. Therefore, clip traction was performed using dental floss, pulling the whole lesion into the gastric lumen (► **Fig. 1**). ESD was accomplished from the distal to the proximal side (► **Fig. 2**), achieving en bloc resection of both lesions on the same pathological specimen (► **Fig. 3**). At the end of

ESD, moderate bleeding was observed from two vessels in the scar, and five hemoclips were applied for successful hemostasis (► **Fig. 4**). Major nonbleeding vessels were coagulated using Coagrasper Hemostatic Forceps (Olympus) to prevent delayed hemorrhage. The procedure time was 60 minutes. The patient did not develop complications and was discharged after 3 days.

Histopathological examination revealed a hyperplastic polyp without dysplasia in the 30-mm lesion, and a moderately differentiated intramucosal adenocarcinoma, without lymphovascular invasion and with negative horizontal margins in the 25-mm lesion (expanded curative criteria).

The pylorus and duodenal bulb have been reported among the most challenging locations for performing ESD, with a greater risk of complications [1–3]. The complete resection rate decreases for tumors that are located in the upper hemisphere, have duodenal extension, and have a large circumferential extent of resection [3]. Different traction systems have been developed to assist ESD.

► **Fig. 1** Traction with dental floss and clip was performed from the duodenal part of the lesion.

► **Fig. 2** Endoscopic submucosal dissection was performed from the distal to the proximal side.

► **Video 1** Traction-assisted endoscopic submucosal dissection for a gastric lesion involving the pyloric ring and duodenal bulb.

► **Fig. 3** En bloc resection was achieved for both lesions in the same pathological specimen.

► **Fig. 4** After completing the resection, moderate bleeding was observed from two vessels in the scar, and five hemoclips were applied.

Among them, dental floss and clip traction is a simple, feasible, and cost-effective method that allows the procedure time to be shortened while achieving en bloc resection [4, 5].

Endoscopy_UCTN_Code_TTT_1AO_2AN

Competing interests

None

The authors

Marta Rodríguez-Carrasco¹, Gonçalo Nunes², Diogo Libânio^{1,3}, Pedro Pimentel-Nunes^{1,3,4}, Mário Dinis-Ribeiro^{1,3}

- 1 Gastroenterology Department, Portuguese Oncology Institute of Porto, Porto, Portugal
- 2 Gastroenterology Department, Hospital Garcia de Orta, Almada, Portugal

- 3 MEDCIDS – Department of Community Medicine, Information and Decision in Health, Faculty of Porto, University of Medicine, Porto, Portugal
- 4 Surgery and Physiology Department, Faculty of Medicine of the University of Porto, Porto, Portugal

Corresponding author

Marta Rodríguez-Carrasco, MD

Gastroenterology Department, Portuguese Oncology Institute of Porto, Rua Dr. Bernardino de Almeida, 4200-072 Porto, Portugal
Fax: +351-22-5513646
martarc7@gmail.com

References

- [1] Imagawa A, Okada H, Kawahara Y et al. Endoscopic submucosal dissection for early gastric cancer: results and degrees of technical difficulty as well as success. *Endoscopy* 2006; 38: 987–990
- [2] Lim CH, Park JM, Park CH et al. Endoscopic submucosal dissection of gastric neoplasia involving the pyloric channel by retroflexion in the duodenum. *Dig Dis Sci* 2012; 57: 148–154
- [3] Bae JH, Kim GH, Lee BE et al. Factors associated with the outcomes of endoscopic submucosal dissection in pyloric neoplasms. *Gastrointest Endosc* 2015; 81: 303–311
- [4] Koike Y, Hirasawa D, Fujita N et al. Usefulness of the thread-traction method in esophageal endoscopic submucosal dissection: randomized controlled trial. *Dig Endosc* 2015; 27: 303–309
- [5] Jeon WJ, You IY, Chae HB et al. A new technique for gastric endoscopic submucosal dissection: peroral traction-assisted endoscopic submucosal dissection. *Gastrointest Endosc* 2009; 69: 29–33

Bibliography

DOI <https://doi.org/10.1055/a-1011-3769>
Published online: 27.9.2019
Endoscopy 2020; 52: E96–E97
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>