Whole-fornix endoscopic submucosal dissection for gastric mucosal adenocarcinoma

► **Fig. 1** A huge superficial elevated lesion was observed in the entire fornix, after marking.

► **Fig. 2** The dissecting plane of the submucosal layer was distinctly observed by pulling the clip-and-line system.

► **Fig. 3** The use of multiple clip-and-line procedures facilitated submucosal dissection from both the anterior and posterior sides.

► **Fig. 4** Mucosal defect after endoscopic submucosal dissection.

► **Fig. 5** The lesion was resected en bloc.

A 69-year-old man with a history of radiation therapy for gastric malignant lymphoma had undergone follow-up esophagogastroduodenoscopy (EGD) at a previous institution, and a superficial elevated lesion was found at the fornix. Examination of a biopsied specimen revealed well-differentiated adenocarcinoma. Endoscopic submucosal dissection (ESD) was attempted, but the procedure was stopped because Mallory–Weiss syndrome occurred in and around the lesion during endoscopic observation immediately before starting ESD. The patient was then referred to our hospital for fur-

ther treatment. EGD at our institution revealed a huge superficial lesion occupying the whole fornix (► **Fig. 1**). As no evidence of invasive cancer was found, we performed endoscopic resection. A multi-bending two-channel scope (GIF-2TQ260M; Olympus Medical Systems, Tokyo, Japan) was used because this device can closely approach the fornix. We performed traction-assisted ESD using the clip-and-line technique [1–4]. After performing a mucosal incision on the anterior side of the lesion using a FlushKnife BT (DK2618JB15; Fujifilm Medical, Tokyo, Japan) and an

ITknife2 (KD-611L; Olympus Medical Systems), we grasped the anterior side of the specimen with the clip-and-line technique. The dissecting plane of the submucosal layer was distinctly observed by pulling the line (► **Fig. 2**), and submucosal dissection was then easily performed. After creating a circumferential incision, a second clip-and-line procedure was applied to facilitate submucosal dissection of the anterior wall [5] (► **Fig. 3**). Perforation occurred during dissection, but the defect was promptly closed by endoscopic clipping. Using a third clip-and-line procedure on the greater curvature side of the specimen, the specimen was resected en bloc (► **Fig. 4**, ► **Fig. 5**). The pathological diagnosis of the resected specimen was an intramucosal adenocarcinoma with a diameter of 110×48 mm.

Endoscopy_UCTN_Code_CPL_1AH_2AZ

Acknowledgement

We thank Angela Morben, DVM, ELS, from Edanz Group (<https://en-author-services.edanzgroup.com/>), for editing a draft of this manuscript.

Video 1 Whole-fornix endoscopic submucosal dissection for gastric mucosal adenocarcinoma.

- [3] Shichijo S, Yamasaki Y, Takeuchi Y. Case of colonic adenoma involving a diverticulum resected by a traction-assisted endoscopic submucosal dissection technique. *Dig Endosc* 2017; 29: 729–730
- [4] Shichijo S, Takeuchi Y, Matsuno K et al. Pulley traction-assisted colonic endoscopic submucosal dissection: a retrospective case series. *Dig Dis* 2019; 37: 473–477
- [5] Yamasaki Y, Harada K, Okada H. Traction-assisted endoscopic submucosal dissection for a giant rectal tumor: multiple clip-and-threads technique. *Dig Endosc* 2018; 30: 697–699

Bibliography

DOI <https://doi.org/10.1055/a-1085-9472>
 Published online: 22.1.2020
Endoscopy 2020; 52: E243–E244
 © Georg Thieme Verlag KG
 Stuttgart · New York
 ISSN 0013-726X

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>

Competing interests

The authors declare that they have no conflict of interest.

The authors

Satoki Shichijo, Yoji Takeuchi, Hiromu Fukuda, Akira Maekawa, Takashi Kanesaka, Noriya Uedo, Ryu Ishihara

Department of Gastrointestinal Oncology,
 Osaka International Cancer Institute, Osaka,
 Japan

Corresponding author

Satoki Shichijo, MD, PhD

Department of Gastrointestinal Oncology,
 Osaka International Cancer Institute,
 3-1-69 Otemae, Cyuo-ku, Osaka 541-8567,
 Japan
 Fax: +81-6-69814067
shichijyou-tyk@umin.ac.jp

References

- [1] Yoshida M, Takizawa K, Suzuki S et al. Conventional versus traction-assisted endoscopic submucosal dissection for gastric neoplasms: a multicenter, randomized controlled trial (with video). *Gastrointest Endosc* 2018; 87: 1231–1240
- [2] Oyama T. Counter traction makes endoscopic submucosal dissection easier. *Clin Endosc* 2012; 45: 375–378