Chemical Constituents of the Terrestrial Stems of *Ephedra sinica* and their PPAR- γ Ligand-Binding Activity

Authors

Yukiko Matsuo¹, Mayu Sasaki¹, Haruhiko Fukaya¹, Katsunori Miyake¹, Riko Takeuchi², Hidetoshi Kumata², Yoshihiro Mimaki¹

Affiliations

- 1 School of Pharmacy, Tokyo University of Pharmacy and Life Science, Hachioji, Japan
- 2 School of Life Sciences, Tokyo University of Pharmacy and Life Science, Hachioji, Japan

Key words

Ephedra sinica, Ephedraceae, sesquiterpene, bisabolane, flavonoid, PPAR- γ

received 04.11.2019

revised 17.12.2019

accepted 06.01.2020

Bibliography

DOI <https://doi.org/10.1055/a-1094-9229>

Planta Med Int Open 2020; 7: e12–e16

© Georg Thieme Verlag KG Stuttgart · New York

ISSN 2509-9264

Correspondence

Dr. Yukiko Matsuo

School of Pharmacy, Tokyo University of Pharmacy and Life Science

1432-1 Horinouchi

192-0392 Hachioji

Japan

matsuoy@toyaku.ac.jp

Supporting Information for this article is available online at: <http://www.thieme-connect.de/products>.

Abstract

Bioassay-guided fractionation of the MeOH extract of *Ephedra sinica* terrestrial stems, using a PPAR- γ ligand binding assay, resulted in the isolation of 10 compounds, including one new bisabolane-type sesquiterpenoid (**10**). The structure of the new compound was determined by extensive spectroscopic analysis, including two-dimensional (2D) NMR. Among the isolated compounds, the sitosterol derivatives (**1** and **2**), flavonoid glucoside (**7**), and the new sesquiterpenoid (**10**), showed significant PPAR- γ ligand-binding activity.

Introduction

Ephedra sinica is described in the Japanese Pharmacopoeia (17th edition) as one of the original plants from which the crude drug Ephedra Herb has been obtained [1]. Ephedra Herb has long been used as an antitussive and expectorant in traditional Japanese medicine [2, 3]. The terrestrial stems of *E. sinica* are well known to contain ephedrine alkaloids, (-)-ephedrine and (+)-pseudoephedrine, and tannins ephedrannin A and B [4–6]. The ephedrine alkaloids in Ephedra Herb are mainly responsible for exerting pharmaceutical effects as well as adverse effects [7, 8]. In addition, only a few bioactive constituents of Ephedra Herb, except for ephedrine alkaloids, have been reported [9]. Pharmacological studies of minor constituents of Ephedra Herb, such as sesquiterpenes and monoterpenes, are needed. As part of our continuous search for bioactive secondary metabolites of traditional medicines, a MeOH extract of *E. sinica* was found to exhibit peroxisome proliferator-activated receptor (PPAR)- γ ligand-binding activity. To identify the secondary metab-

olites in *E. sinica* having PPAR- γ ligand-binding activity, bioassay-guided fractionation of the MeOH extract of *E. sinica* terrestrial stems was carried out. This led to the isolation of 10 compounds (**1–10**), including a new bisabolane-type sesquiterpenoid (**10**) (► **Fig. 1**). This paper is a report on the structural determination of the new sesquiterpene obtained by extensive spectroscopic analysis, including 2D NMR data. The isolated compounds were evaluated for their PPAR- γ ligand-binding activity.

Results and Discussion

The terrestrial stems of *E. sinica* (5.4 kg dry weight) were extracted with MeOH at 50 °C. After removing the solvent, the MeOH extract (665 g) was applied to a porous polymer polystyrene resin (Diaion HP-20) column. The MeOH, EtOH, and EtOAc-eluted portions showed PPAR- γ ligand-binding activity (► **Fig. 2**). Thus, the MeOH and EtOH-soluble fractions were passed through column chroma-

► **Fig. 1** Isolated compounds from *E. sinica*.

tography on silica gel and octadecylsilanized (ODS) silica gel producing compounds **1–10**. Compounds **1–9** were identified as β -sitosterol (**1**) [10], 7β -hydroxysitosterol (**2**) [11], 7α -hydroxysitosterol (**3**) [11], kaempferol-3-*O*-rhamnoside (**4**) [12], kaempferol-3-*O*-(4''-*trans*-*p*-coumaroyl)-rhamnopyranoside (**5**) [13], kaempferol-3-*O*-(4''-*cis*-*p*-coumaroyl)-rhamnopyranoside (**6**) [14], herbacetin-7-*O*-glucopyranoside (**7**) [6], mahuanin A (**8**) [15], and ephedrannin A (**9**) [16], respectively (► **Fig. 1**).

Compound **10** ($C_{16}H_{22}O_4$) was obtained as a colorless oil, which was determined by high-resolution electrospray ionization time-of-flight mass spectrometry (HR-ESI-TOF-MS, m/z 301.1413 [$M + Na$] $^+$, calcd. for $C_{16}H_{22}NaO_4$ 301.1416) and ^{13}C NMR (16 carbon signals) data. The IR spectrum of **10** showed absorption bands of carbonyl groups at 1700 and 1644 cm^{-1} . The ^{13}C NMR spectrum exhibited two carbonyl carbon signals at δ_C 172.0 and 168.1, and the HMBC spectrum displayed a correlation peak between δ_C 168.1 and a methyl singlet signal at δ_H 3.74. On the other hand, treatment of **10** with bis(trimethylsilyl)trifluoroacetamide (BSTFA) in pyridine produced the corresponding trimethylsilyl (TMS) derivative of **10** ($C_{19}H_{30}O_4Si$; GC/MS, m/z : 350.1 [$M + TMS$]). These data implied the presence of two carboxy groups in the molecule of **10**, one is free and another is methyl ester. The 1H NMR and 1H - 1H COSY spectra disclosed the following spin-coupling correlations: δ_H 7.16

(H-2)/ 2.73 (H-3a) and 2.20 (H-3b)/2.89 (H-4)/ 2.12 (H-5a) and 1.74 (H-5b) /2.54 (H-6a) and 2.24 (H-6b), and the H-2 olefinic proton and H₂-6 methylene protons showed long-range correlations with the quaternary olefinic carbon at δ_C 129.2 (C-1) and the carbonyl carbon of the carboxy group at δ_C 172.0 (C-7). These data provided evidence of the presence of a cyclohexene group bearing a carboxy group at C-1. The other the spin-coupling correlations δ_H 1.06 (Me-14) and 1.05 (Me-15)/2.44 (H-13)/6.09 (H-12)/6.35 (H-11)/7.16 (H-10) were consistent with a 4-methyl-2-pentene group. The HMBC correlations from H₂-3, H-4, H₂-5, and H-11 to the olefinic carbon at δ_C 131.8 (C-8) and from H-4 to the carbonyl carbon of the methyl carboxylate group at δ_C 168.1 (C-9) indicated that C-4 of the cyclohexene group, C-9 carbonyl group, and C-10 of the 4-methyl-2-pentene group were linked to the C-8 olefinic carbon, and a double bond was present between C-8 and C-10 (► **Fig. 3**). The spin-coupling constant of H-10/H-11 ($J = 11.4$ Hz) and H-11/H-12 ($J = 15.0$ Hz) and NOE correlations between H-10 and H-12 and between H-11 and H₂-3 allowed the geometry at C-8 and C-11 to be established as *8E* and *11E*. In ESI-TOF/MS-MS analysis of **10**, the prominent fragment ion-peak at m/z 207 originated from a loss of the $C_{(11)}H-C_{(12)}H-C_{(13)}H-Me_{(14)}(Me_{(15)})$ moiety of **10**. Other minor peaks observed at m/z 245, 235, 233, 201, 179, and 163 corresponded to the proposed fragments as depicted in the **Supporting Informa-**

► **Fig. 2** PPAR- γ ligand-binding activity of the MeOH extract of *E. sinica* and fractions. The PPAR- γ ligand-binding activity of each fraction was measured at sample concentrations of 1000 and 10 μ g/mL and that of pioglitazone at 5.0 μ M. Data are represented as the mean $\pm$ SEM of three experiments performed in triplicate; *** p < 0.0001 vs. control, ** p < 0.001 vs. control, * p < 0.01 vs. control.

► **Fig. 3** ^1H - ^1H COSY and selected HMBC correlations of **10**.

tion. Thus, the structure of **10** was established as a bisabolane-type sesquiterpenoid as shown in ► **Fig. 1**. Compound **10** showed no specific rotation and was assumed to be a racemate.

The isolated compounds **1**, **2**, and **4–10** were evaluated for their PPAR- γ ligand-binding activity using a nuclear receptor cofactor assay system (► **Fig. 4**). Kaempferol was used as a positive control. Compounds **1**, **2**, **7**, and **10** exhibited significant PPAR- γ ligand-binding activity, among which the new bisabolane-type sesquiterpenoid **10** was the most potent. In the kaempferol derivatives (**4–7**), the glycosylation at the C-2 hydroxy group by a rhamnosyl moiety (**4–6**) reduced the PPAR- γ ligand activity.

In conclusion, a new bisabolane-type sesquiterpenoid (**10**) and nine known compounds (**1–9**) were isolated from the MeOH extract of *E. sinica* terrestrial stems. Compounds **1**, **2**, **7**, and **10** showed significant PPAR- γ ligand-binding activity, among which the new bisabolane derivative **10** was the most potent. In a previous study, terpenoids with a cyclohexene ring were reported to act as a native retinoid X receptor agonist [17]. With expectation, **10** exhibited PPAR- γ ligand-binding activity.

Materials and Methods

General experimental procedures

The instruments and experimental conditions were the same as those described in a previous paper [18]. All solvents used for extraction and isolation were high grade (> 99%) (Fujifilm Wako Pure Chemical). Purities of all isolated compounds (> 95%) were confirmed by NMR and TLC analysis.

Plant material

The dried terrestrial parts (5.4 kg) of *E. sinica* Stapf (Ephedraceae) were collected from the Medicinal Plant Garden of the Tokyo University of Pharmacy and Life Sciences, Tokyo, Japan on July 10, 2015. This genetic resource was introduced to the garden in December 1981. The plant material was authenticated by one of the authors (K. M.) and was also identified according to the ITS rDNA sequence [19]. A voucher specimen has been deposited in the Herbarium of Tokyo University of Pharmacy and Life Sciences (KS-2015–001).

Extraction and isolation

The terrestrial stems of *E. sinica* (5.4 kg) were extracted with MeOH (50 L) at 50 °C. After evaporation of MeOH under reduced pressure at 40 °C, the MeOH extract (665 g) was subjected to a Diaion HP-20 column (2200 g, 85 mm i.d. $\times$ 600 mm) and successively partitioned by eluting with MeOH-H₂O (3:7), MeOH-H₂O (1:1), MeOH, EtOH, and EtOAc (each 10 L) in order of decreasing polarity. The EtOH-eluted fraction (55 g) was passed through a silica gel column (2000 g, 85 mm i.d. $\times$ 600 mm) eluted with gradient mixtures of hexane-EtOAc (4:1; 3:1; 2:1; 1:1) and MeOH, which produced 9 fractions (Frs. A–I). Fr. C was separated by a silica gel column (1800 g, 80 mm i.d. $\times$ 400 mm) eluted with hexane-EtOAc (7:1; 4:1; 1:1) and, sequentially, an ODS silica gel column eluted with MeCN–

► **Fig. 4** PPAR- γ ligand-binding activity of **1**, **2**, and **4–10**. PPAR- γ ligand-binding activities of **1**, **2**, **4–10**, and kaempferol were measured at a sample concentration of 0.5 mM. Data are represented as the mean $\pm$ SEM of three experiments performed in triplicate; *** p < 0.0001 vs. control, * p < 0.01 vs. control.

► **Table 1** ^1H and ^{13}C NMR (500 and 125 MHz, CDCl_3) spectroscopic assignments of **10**.

Position	δ_{H}		J (Hz)	δ_{C}
1	—			129.2
2	7.16	br d	2.5	142.2
3	2.73	ddd	19.5, 11.2, 2.5	30.5
	2.20	m		
4	2.89	m		33.4
5	2.12	qd	12.4, 5.0	26.4
	1.74	br d	12.4	
6	2.54	ddd	16.9, 2.3, 2.0	24.7
	2.24	m		
7	—			172.0
8	—			131.8
9	—			168.1
10	7.16	br d	11.4	140.2
11	6.35	ddd	15.0, 11.4, 1.1	121.9
12	6.09	ddd	15.0, 7.1, 0.6	151.4
13	2.44	sd	7.1, 1.1	31.9
14	1.06	d	6.7	22.0
15	1.05	d	6.7	22.0
16	3.74	s		51.4

H_2O (3:1; 4:1; 5:1) to yield **1** (44.8 mg), **2** (1.5 mg), and **3** (4.3 mg). The MeOH-eluted portion (120 g) was chromatographed on silica gel (2700 g, 45 mm i.d. $\times$ 430 mm) eluted with MeCN- H_2O (1:3; 1:2; 1:1; 2:1) to give 10 subfractions (Frs. a-j). Fr. a was subjected to a Sephadex LH-20 column (470 g, 25 mm i.d. $\times$ 300 mm) eluted with MeOH- H_2O (1:1; 2:1) to give **4** (58.0 mg), **7** (6.7 mg), and **8** (17.0 mg). Fr. e was subjected to a Sephadex LH-20 column (470 g, 25 mm i.d. $\times$ 300 mm) eluted with MeOH- H_2O (2:1) and a silica gel column eluted with CHCl_3 -MeOH- H_2O (100:10:1; 50:10:1; 30:10:1) to yield **5** (25.2 mg), **6** (6.2 mg), and **9** (7.1 mg). Fr. i was subjected to a Sephadex LH-20 column (470 g, 25 mm i.d. $\times$ 300 mm) eluted

with MeOH- H_2O (2:1) and an ODS silica gel column eluted with MeCN- H_2O (2:1) to yield **10** (14.9 mg). Separation and isolation were guided a PPAR- γ ligand-binding activity and TLC analysis of the fractions. (► **Fig. 1S–8S**).

Compound **10**

An amorphous solid; $[\alpha]_{\text{D}}^{25}$ -0.22 (c 0.10, CHCl_3); IR ν_{max} (film) cm^{-1} : 3414 (OH), 1700 and 1644 ($\text{C}=\text{O}$); ^1H and ^{13}C NMR (500 and 125 MHz, CDCl_3), see ► **Table 1**; HR-ESI-TOF-MS (m/z : 301.1413 $[\text{M} + \text{Na}]^+$, calcd. for $\text{C}_{16}\text{H}_{22}\text{NaO}_4$: 301.1416).

Derivatization of **10**

A mixture of **10** (0.5 mg) and BSTFA (50 μL) in pyridine (50 μL) was sealed in a glass vial and placed at 60 $^\circ\text{C}$ for 3 h. The reaction mixture was evaporated under a gentle stream of nitrogen to give the TMS derivative of **10**.

PPAR- γ ligand-binding activity

PPAR- γ agonist activity was examined using a nuclear receptor co-factor assay system (EnBio RCAS for PPAR- γ ; EnBioTec Laboratories) according to the manufacturer's instructions. Briefly, a peptide of cyclic AMP response element-binding protein (CBP) was immobilized on the bottom of a microtiter plate. After adding the recombinant human PPAR- γ solution to the wells, DMSO (purity > 99.5%) as a control, or a positive control or isolated compounds were added, respectively. The binding of the PPAR- γ ligand complex to the CBP on the plate was detected by measuring the absorbance at 450 nm. This assay involves a cell-free system using nuclear receptors and cofactors. Pioglitazone (purity > 98.0%) and kaempferol (purity > 97.0%) were used as positive controls (TCI). All other reagents or solvents used were of biochemical reagent grade.

Statistical analyses

Data are represented as the mean $\pm$ standard error of the mean (SEM) of three experiments performed in triplicate. Dunnett's test was used and the level of significance is indicated by p values.

Supporting information

1D and 2D NMR and MS/MS spectra of **10** are available as Supporting Information.

Acknowledgements

We thank Hirokazu Ando and Yohei Sasaki for identifying *E. sinica* according to the ITS rDNA sequence.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- [1] Ministry of Health Law, Japan The Japanese pharmacopoeia. 17th edition, Tokyo: Ministry of Health Law; 2016: 1589
- [2] Minamizawa K, Goto H, Shimada Y, Terasawa K, Haji A. Effects of eppikahangeto, a Kampo formula, and Ephedrae herba against citric acid-induced laryngeal cough in guinea pigs. *J Pharm Sci* 2006; 101: 118–125
- [3] Abourashed EA, Alfay AT, Khan IA, Walker L. Ephedra in perspective – a current review. *Phytother Res* 2003; 17: 703–712
- [4] Kim HK, Choi YH, Chang WT, Verpoorte R. Quantitative analysis of ephedrine analogues from Ephedra species using ¹H-NMR. *Chem Pharm Bull* 2003; 51: 1382–1385
- [5] Amakura Y, Yoshimura M, Yamakami S, Yoshida T, Wakana D, Hyuga M, Goda Y. Characterization of phenolic constituents from Ephedra herb extract. *Molecules* 2013; 18: 5326–5334
- [6] Nawwar MA, El-Sissi HI, Barakat HH. Flavonoid constituents of *Ephedra alata*. *Phytochemistry* 1984; 23: 2937–2939
- [7] Shekelle P, Hardy ML, Morton SC, Maglione M, Suttrop M, Roth E, Jungvig L, Mojica WA, Gagné J, Rhodes S, McKinnon E. Ephedra and ephedrine for weight loss and athletic performance enhancement: Clinical efficacy and side effects. *Evid Rep Technol Assess (Summ)* 2003; 76: 1–4
- [8] Kobayashi Y. Analgesic effects and side effects of ephedra herb extract and ephedrine alkaloids-free ephedra herb extract. *Yakugaku Zasshi* 2017; 137: 187–194
- [9] Hyuga S, Hyuga M, Yoshimura M, Amakura Y, Goda Y, Hanawa T. Herbacetin, a constituent of ephedrae herba, suppresses the HGF-induced motility of human breast cancer MDA-MB-231 cells by inhibiting c-Met and Akt phosphorylation. *Planta Med* 2013; 79: 1525–1530
- [10] Morales G, Sierra P, Mancilla A, Paredes A, Loyola LA, Gallardo O, Borquez J. Secondary metabolites from four medicinal plants from northern Chile: antimicrobial activity and biotoxicity against *Artemia salina*. *J Chin Chem Soc* 2003; 48: 13–18
- [11] Cui EJ, Park JH, Park HJ, Chung IS, Kim JY, Yeon SW, Baek NI. Isolation of sterols from cowpea (*Vigna sinensis*) seeds and their promotion activity on HO-1. *J Korean Soc Appl Biol Chem* 2011; 54: 362–366
- [12] Bernal F, Cuca L, Yamaguchi L, Coy-Barrera E. LC-DAD-UV and LC-ESI-MS-based analyses, antioxidant capacity, and antimicrobial activity of a polar fraction from *Iryanthera ulei* leaves. *Rec Nat Prod* 2013; 7: 152–156
- [13] Kaouadji M, Morand JM, Garcia J. Further acylated kaempferol rhamnosides from *Platanus acerifolia* buds. *J Nat Prod* 1993; 56: 1618–1621
- [14] Si CL, Yu GJ, Du ZG, Huang XF, Fan S, Du HS, Hu WC. A new cis-p-coumaroyl flavonol glycoside from the inner barks of *Sophora japonica* L. *Holzforchung* 2016; 70: 39–45
- [15] Rawat MSM, Prasad D, Joshi RK, Pant G. Proanthocyanidins from *Prunus armeniaca* roots. *Phytochemistry* 1999; 50: 321–324
- [16] Kim IS, Park YJ, Yoon SJ, Lee HB. Ephedrannin A and B from roots of *Ephedra sinica* inhibit lipopolysaccharide-induced inflammatory mediators by suppressing nuclear factor-κB activation in RAW 264.7 macrophages. *Int Immunopharmacol* 2010; 10: 1616–1625
- [17] Okitsu T, Sato K, Iwatsuka K, Sawada N, Nakagawa K, Okano T, Wada A. Replacement of the hydrophobic part of 9-cis-retinoic acid with cyclic terpenoid moiety results in RXR-selective agonistic activity. *Bioorg Med Chem* 2011; 19: 2939–2949
- [18] Matsuo Y, Maeda S, Ohba C, Fukaya H, Mimaki Y. Vetiverianines A, B, and C: Sesquiterpenoids from *Vetiveria zizanioides* Roots. *J Nat Prod* 2016; 79: 2175–2180
- [19] Long C, Kakiuchi N, Takahashi A, Komatsu K, Cai S, Mikage M. Phylogenetic Analysis of the DNA Sequence of the Non-Coding Region of Nuclear Ribosomal DNA and Chloroplast of Ephedra Plants in China. *Planta Med* 2004; 70: 1080–1084