Stratification of Breast Cancer Risk in Terms of the Influence of Age and Mammographic density

Stratifizierung des Brustkrebsrisikos hinsichtlich der Einflüsse von Alter und mammografischer Dichte

Authors

Stefanie Weigel¹, Walter Heindel¹, Caroline Dietz², Ulrike Meyer-Johann³, Axel Graewingholt⁴, Hans Werner Hense⁵

Affiliations

- 1 Institute of Clinical Radiology and Reference Center for Mammography Münster, University and University Hospital Münster, Germany
- 2 Screening Unit, Leverkusen, Germany
- 3 Screening Unit, Bielefeld, Germany
- 4 Screening Unit, Paderborn, Germany
- 5 Institute of Epidemiology and Social Medicine, University of Münster, Germany

Key words

breast cancer screening, digital mammography, breast density, age, interval cancer, population surveillance

received 10.03.2019

accepted 22.12.2019

Bibliography

DOI <https://doi.org/10.1055/a-1100-0016>

Published online: 27.2.2020

Fortschr Röntgenstr 2020; 192: 678–685

© Georg Thieme Verlag KG, Stuttgart · New York

ISSN 1438-9029

Correspondence

Prof. Stefanie Weigel

Institute of Clinical Radiology, University and University Hospital Münster, Albert-Schweitzer-Campus 1, 48149 Münster, Germany

Tel.: ++49/2 51/8 34 56 50

Fax: ++49/2 51/8 34 56 60

weigels@uni-muenster.de

ABSTRACT

Purpose Analysis of the influence of the singular risk factors age and breast density on the 2-year incidence of breast cancer among participants in the German mammography screening program.

Materials and Methods The multicenter study includes 111 456 subsequent round digital mammographic screening examinations from four screening units with prospective visual categorization of breast density. Based on detection in screening and during the 2-year interval after negative screening participation (interval cancers), 2-year breast cancer

incidences (2 YBCI) (‰) were calculated in the 5-year age groups (5 YAG) of the target group 50–69 years and in the BI-RADS density categories ACR 1–4. Multivariate statistical evaluations were carried out using logistic regression models. **Results** With an increase in the 5 YAG, the 2 YBCI increased by 5.0 ‰, 6.7 ‰, 8.5 ‰ to 9.7 ‰, and was significantly different among 55–59, 60–64 and 65–69-year-old women compared to the youngest reference group 50–54 years (odds ratio (OR): 1.34; 1.68; and 1.93; p-value < 0.0001). With an increase in density categories 1–4, the 2 YBCI increased from 2.6 ‰, to 5.8 ‰, 9.6 ‰, and 9.7 ‰. The 2 YBCI differed significantly in breast density categories 2, 3, 4 from reference group 1 (OR: 2.17; 3.65; and 3.76; p-value < 0.0001). Only within the two main breast density groups 2 (frequency 44.3 %) and 3 (44.7 %), a significant increase in the 2 YBCI was observed across the 5 YAG (category 2: 3.7–8.9 ‰; category 3: 5.8–11.7 ‰; p-value < 0.001 each). The 2 YBCI was above the median of 7.5 ‰ in women with breast density category 2 and aged 65–69 years, as well as in women with breast density categories 3 and 4 aged 55–69 years. A 2 YBCI below the median was seen in women between 50–54 years regardless of breast density, as well as women in category 1 in all age groups.

Conclusion Within the main breast density categories 2 and 3 (almost 90 % of participants), incidences increase with age to double. A consistently low incidence is found regardless of breast density at a young screening age and in women with the lowest breast density.

Key Points:

- The risk of breast cancer is modified by age in density categories.
- Women aged 50–54 years have a low risk in all density categories.
- Women in category ACR 1 of any age group have a low risk.

Citation Format

- Weigel S, Heindel W, Dietz C et al. Stratifizierung des Brustkrebsrisikos hinsichtlich der Einflüsse von Alter und mammografischer Dichte. Fortschr Röntgenstr 2020; 192: 678–685

ZUSAMMENFASSUNG

Ziel Analyse des Einflusses der singulären Risikofaktoren Lebensalter und Brustdicke auf die 2-Jahres-Inzidenz des Mammakarzinoms unter Teilnehmerinnen am deutschen Mammografie-Screening-Programm.

Material und Methode Die multizentrische Studie umfasst 111 456 digitale mammografische Folgerunden-Untersuchungen aus 4 Screening-Einheiten mit prospektiver visueller Kategorisierung der Brustdicke. Basierend auf der Detektion im Screening und im 2-Jahres-Intervall nach negativer Screening-Teilnahme (Intervallkarzinome) wurden 2-Jahres-Brustkrebsinzidenzen (2-JBI) (%) in den 5-Jahres-Altersgruppen (5-JAG) der Zielgruppe 50–69 Jahre und in den Dichtekategorien ACR 1–4 berechnet. Multivariate statistische Auswertungen erfolgten mit logistischen Regressionsmodellen.

Ergebnisse Mit steigender 5-JAG stieg die 2-JBI von 5,0 %, 6,7 % und 8,5 % auf 9,7 % und unterschied sich unter den 55–59-, 60–64- und 65–69-jährigen Frauen signifikant von der jüngsten Referenzgruppe 50–54 Jahre (Odds-Ratio (OR):

1,34, 1,68 bzw. 1,93; $p < 0,0001$). Mit Anstieg der ACR-Kategorien 1–4 nahm die 2 JBI von 2,6 %, auf 5,8 %, 9,6 % und 9,7 % zu. Die 2-JBI unterschied sich in ACR 2, 3 und 4 signifikant von der Referenzgruppe ACR 1 (OR: 2,17, 3,65 bzw. 3,76; $p < 0,0001$). Nur innerhalb der beiden Hauptgruppen ACR 2 (Häufigkeit 44,3 %) und ACR 3 (44,7 %) zeigte sich ein signifikanter Anstieg der 2-JBI über die 5-JAG (ACR 2: 3,7 %–8,9 %; ACR 3: 5,8 %–11,7 %; $p < 0,001$). 2-JBI lagen über dem Median von 7,5 % bei Frauen mit ACR 2 und Alter 65–69 Jahre sowie bei ACR 3 und ACR 4 im Alter 55–69 Jahre. 2-JBI unter dem Median hatten Frauen zwischen 50–54 Jahren, unabhängig von der Brustdicke, sowie Frauen der Kategorie ACR 1 in allen Altersgruppen.

Schlussfolgerung Innerhalb der Hauptkategorien ACR 2 und ACR 3 (fast 90 % der Teilnehmerinnen) steigen die Inzidenzen mit zunehmendem Alter bis zu einer Verdopplung. Eine konsistent niedrige Inzidenz findet sich unabhängig von der Brustdicke bei jungem Screening-Alter und bei Frauen mit geringster Brustdicke.

Introduction

Breast cancer is the most common fatal cancer among women [1]. Population-based mammography screening programs have been implemented to allow earlier diagnosis of breast cancer by means of systematic imaging and to help to reduce the mortality rate as a result of earlier initiation of treatment [2]. There is an estimated reduction in breast cancer mortality of 20–40 % for women between 50 and 69 years of age participating in mammography screening [3].

Established risk factors for breast cancer include age, precancerous precursor lesions, genetic predisposition, positive personal and family history, hormone therapies, ionizing radiation, alcohol consumption, and postmenopausal obesity [3]. While structured intensified surveillance for early detection is performed in the case of a genetic predisposition for breast cancer [4], the justification for early detection examinations as part of population-based mammography screening programs is based on the risk factor “age”. National statistics show that age is a relevant risk factor for breast cancer. Therefore, 1 in 110 women 35 years old will develop breast cancer in the next 10 years. The risk is highest at 65 years of age. 1 in 27 women is affected by the age of 75 [5].

Studies have also shown that higher breast density is also a strong independent biomarker for an increased risk of breast cancer [6–9]. Breast density can be determined from mammography and relates to the ratio of radiopaque epithelium and stroma to radiolucent fatty tissue. In addition, higher breast density has a masking effect on mammography [6, 7].

Scientific approaches focus on the development of risk-adjusted early detection concepts for women also outside of high-risk groups for breast cancer [10]. However, knowledge of the combination of risk constellations is currently still limited. Detection requires consideration of breast cancer diagnoses in the interval between screening examinations. Although this information can

be obtained from population-based cancer registries including tumor-specific data, such as staging, grading, etc., data regarding imaging criteria, such as mammographic density values, is not available there.

Therefore, the goal of the present study is to analyze the effect of age and breast density on the incidence of breast cancer in a 2-year period among participants in the German mammography screening program.

Materials and Methods

Screening procedure and data collection

The national mammography screening program is based on the European guidelines [2]. The target group is women aged 50–69 years. All eligible women in the population are mailed an invitation to participate in two-year intervals.

111 456 of 111 470 subsequent digital screening examinations that were generated at four screening units between January 2009 and December 2010 and comprised ACR categorization were included in the study. Independent double reading provided the evaluation of abnormal mammography findings [11] as well as independent visual categorization of breast density according to the American College of Radiology Breast Imaging Reporting and Data System Version 4. The percentage of dense tissue increases by 25 % per category: ACR 1 lipomatous, ACR 2 fibroglandular, ACR 3 heterogeneously dense, ACR 4 extremely dense [12]. For study purposes the higher ACR category was used if the independent second reading yielded a different density category for an examination.

All examinations were obtained using the digital technique in two views (cranio-caudal, medio-lateral-oblique) (Lorad Selenia, Hologic; MammoDiagnost, Philips; Mammomat 3000 Nova, Sie-

mens AG; MicroDose Mammography, Sectra Medical Systems; Senographe DS, GE Healthcare; Senographe Essential, GE Healthcare).

According to the defined screening procedure, a consensus conference including both readers and the physician responsible for the program was held in the case of an abnormal mammography finding by at least one interpreting physician in order to decide whether further diagnostic workup is indicated. Diagnostic workup was performed in accordance with regulations by the physician responsible for the program [11].

Screening was documented by screening software with an interface to the national cancer registry in North Rhine-Westphalia. The national cancer registry receives electronic messages about all cases of breast cancer detected for the first time as part of and outside of mammography screening (incidence report). Breast cancers are recorded as invasive breast cancer (ICD-10 C50) and ductal carcinoma in situ (DCIS; ICD-10 D05). A differentiation is made between cases detected during screening and cases of interval cancer. The interval is defined as the 24-month period following participation in the mammography screening program with a negative examination result. The study included corresponding collection of interval cancer data including December 2012.

Statistical calculations

Breast cancer incidences were calculated as the sum of the breast cancer cases detected during screening and the cases of interval cancer detected in the 2-year interval after a negative screening result in relation to the total number of screened women.

The breast cancer incidences were calculated in the 5-year age groups and in the density categories ACR 1–4 per 1000 screening participants.

The program sensitivity (PS) was calculated as the number of screening-detected breast cancer cases divided by the sum of screening-detected breast cancer cases plus interval cancers.

Multivariate statistical evaluations were performed with logistic regression models in which the four age groups and the four ACR categories were defined as binary dummy variables. In addition, an adjustment to the four screening units was performed. The results of the regression models are presented as odds ratios with 95 % confidence intervals. Moreover, a p-value for an age or ACR variable with four attribute levels was determined. The software SAS for Windows 9.4 was used for the evaluations.

Results

Stratification by age

The total group of 111,456 examinations of all age groups had a 2-year breast cancer incidence of 7.5 %. The percentages of the 5-year age groups among screening participants ranged from 21 % to 29 %.

The 2-year breast cancer incidences were 5.0 %, 6.7 %, 8.5 %, and 9.7 %, respectively for the 5-year age groups in increasing order. Older participants had a significantly higher 2-year incidence than the reference group of 50–54-year-olds (odds ratio

(OR: 1.34 (55–59 years); 1.68 (60–64 years); 1.93 (65–69 years); p-value < 0.0001) (► Table 1).

The group as a whole had a program sensitivity of 77.7 %. The values were lower in women under the age of 60 and higher in women 60 or older (► Table 1).

Stratification by breast density category

The means of the visual density categories according to ACR for the first and second reading were 2.32 ± 0.71 and 2.31 ± 0.71 , respectively. A discrepancy resulting in a higher classification between the first and second reading was seen in 16.7 % (18,622/111,456) of the examinations. 30.7 % of these cases with a higher category were category ACR 2, 57.6 % were category ACR 3, and 11.7 % were category ACR 4.

In the group as a whole, ACR categories 2 and 3 were most common with a percentage of 44 % and 45 %, respectively, while ACR 1 and ACR 4 were less common with a percentage of 6 % and 5 %, respectively.

The 2-year breast cancer incidences for ACR categories 1 to 4 were 2.6 %, 5.8 %, 9.6 % and 9.7 %, respectively. The breast cancer incidence for ACR 2, 3 and 4 differed significantly from ACR 1 as the reference group (OR = 2.17, 3.65, and 3.76; p-value < 0.0001) (► Table 2).

The largest density groups ACR 2 and ACR 3 had program sensitivities of 84.0 % and 74.7 %, respectively (► Table 2).

Stratification by age group and breast density category

In the subgroups resulting from the simultaneous stratification by age group and ACR category, the 2-year breast cancer incidence varied from 1.6 % among 55–59 year-olds in category ACR 1 to 13.3 % among 60–64-year-olds in category ACR 4. Increased breast cancer incidences over the total median (> 7.5 %) were seen in the case of the following combinations: ACR 2 and the oldest age group, ACR 3 and all age groups except the youngest, ACR 4 and all age groups except the youngest. Lower 2-year incidences for breast cancer were seen in women in the youngest age group (50–54 years) in every ACR category and women in category ACR 1 in every age group (► Table 3).

In both of the main groups (ACR 2 and ACR 3), a significant increase in the 2-year breast cancer incidence among the 5-year age groups was seen (ACR 2: 3.7 % to 8.9 %; ACR 3: 5.8 % to 11.7 %; p-value < 0.001) (► Fig. 1).

► Fig. 2 shows the discrepancies in the frequency distribution of breast cancers in the 2-year time period depending on breast density and age.

Discussion

The singular risk factors “age” and “breast density” have a cumulative effect on the breast cancer incidence in the main groups “fibroglandular” and “heterogeneously dense”. The increase in incidence between the 5-year age groups in the two most prevalent ACR groups ranged from 3.7 % to 8.9 % (ACR 2) and from 5.8 % to 11.7 % (ACR 3). In addition, the age-specific differences

► **Table 1** Number of women screened in subsequent rounds, number of women with screening (ScD)- and interval (ICD)-detected breast cancers (invasive breast cancers plus ductal carcinoma in situ (DCIS)); per 1000 women screened, by 5-year age group.

	total	5-year age groups				p-value ¹
	50–69 years	50–54 years	55–59 years	60–64 years	65–69 years	
number screened (%)	111 456 (100.0)	23 954 (21.49)	32 201 (28.89)	26 360 (23.65)	28 941 (25.96)	
number of women with ScD plus ICD breast cancer,	838	120	216	221	281	
ScD; ICD	651; 187	87; 33	160; 56	178; 43	226; 55	
2-year breast cancer incidence (per 1,000 women screened)	7.5	5.0	6.7	8.5	9.7	
program sensitivity (ScD/[ScD + ICD])	77.7 %	72.5 %	74.2 %	80.7 %	80.4 %	
adjusted ² odds ratio (95 % confidence interval)		Ref.	1.34 (1.07–1.68)	1.68 (1.34–2.10)	1.93 (1.55–2.39)	P < 0.0001

¹ p-value for one age variable with four levels (1 = '50–54', 2 = '55–59', 3 = '60–64', 4 = '65–69') as obtained from the multivariable logistic regression model, adjusted for screening unit.

² from a multivariable logistic regression model, with dummy variables for each age group (reference 50–54 years), adjusted for screening unit.

► **Table 2** Number of women screened in subsequent rounds, number of women with screening (ScD)- and interval (ICD)-detected breast cancers (invasive breast cancers plus ductal carcinoma in situ (DCIS)); per 1000 women screened, by ACR categories.

	total	ACR 1	ACR 2	ACR 3	ACR 4	p-value ¹
number screened (%)	111 456 (100.0)	6512 (5.84)	49 348 (44.28)	49 819 (44.70)	5777 (5.18)	
number of women with ScD plus ICD breast cancer,	838	17	287	478	56	
ScD; ICD	651; 187	13; 4	241; 46	357; 121	40; 11	
2-year breast cancer incidence (per 1,000 women screened)	7.5	2.6	5.8	9.6	9.7	
program sensitivity (ScD/[ScD + ICD])	77.7 %	76.5 %	84.0 %	74.7 %	71.4 %	
adjusted ² odds ratio (95 % confidence interval)		Ref.	2.17 (1.33–3.55)	3.65 (2.25–5.92)	3.76 (2.18–6.48)	< 0.0001

¹ p-value for one ACR variable with four levels as obtained from multivariable logistic regression model, adjusted for screening unit.

² from a multivariable logistic regression model, with dummy variables for ACR category (reference ACR 1), adjusted for screening unit.

between the ACR groups increased with age. The breast cancer incidence in density category ACR 3 among 50–54-year-olds was 2.1 % higher than in ACR 2, 5.3 % higher among 55–59-year-olds and 6.3 % higher among 60–64-year-olds. However, the difference decreased to 2.8 % among 65–69-year-olds. The highest incidence was 13.3 % for women between 60 and 64 years old in category ACR 4.

The median of the 2-year incidence was 7.5 % in the study population. Increased incidences above this median value were seen in women in breast density categories ACR 3 and ACR 4 except in the youngest age group and in the case of a combination of ACR 2 and the oldest age group. In contrast, a lower 2-year risk was seen for all age groups with density category ACR 1, for wom-

en in the youngest age group regardless of the density category and for ACR 2 except in the oldest age group.

The observations on the singular strongest risk factor for breast cancer “age”, the singular risk factor “breast density”, and their combination show consistent risk stratifications, published as an absolute 5-year risk for breast cancer [13]. The results of the multicenter approach regarding breast density are in agreement with the previously published results of an individual unit that is integrated in the present evaluation: There is a significant increase in the 2-year incidence from ACR 1 to ACR 3. ACR 4 is comparable to ACR 3 and does not result in a further increase [8]. In the multicenter approach, the program sensitivities differed from those of the published single-center approach for

► **Table 3** Number and 2-year breast cancer incidences of women with screening (ScD)- and interval (ICD)-detected breast cancers (invasive breast cancers plus ductal carcinoma in situ (DCIS)); per 1000 women screened for ACR categories, by age group.

	5-year age groups				p-value ¹
	50–54 years	55–59 years	60–64 years	65–69 years	
number of women in ACR 1	919	1829	1718	2046	
number with ScD+ICD breast cancer	4	3	3	7	
2-year breast cancer incidence (per 1000 women screened)	4.4	1.6	1.7	3.4	
adjusted ² odds ratio (95 % confidence interval)	ref.	0.37 (0.08–1.67)	0.39 (0.09–1.77)	0.78 (0.23–2.67)	0.45
number of women in ACR 2	8766	13 788	12 261	14 533	
number with ScD+ICD breast cancer	32	57	69	129	
2-year breast cancer incidence (per 1000 women screened)	3.7	4.1	5.6	8.9	
adjusted ² odds ratio (95 % confidence interval)	ref.	1.13 (0.73–1.74)	1.53 (1.01–2.33)	2.42 (1.64–3.57)	<0.001
number of women in ACR 3	12 302	14 829	11 255	11 433	
number with ScD+ICD breast cancer	71	139	134	134	
2-year breast cancer incidence (per 1000 women screened)	5.8	9.4	11.9	11.7	
adjusted ² odds ratio (95 % confidence interval)	ref.	1.63 (1.22–2.17)	2.07 (1.55–2.76)	2.03 (1.52–2.72)	<0.001
number of women in ACR 4	1967	1755	1126	929	
number with ScD+ICD breast cancer	13	17	15	11	
2-year breast cancer incidence (per 1000 women screened)	6.6	9.7	13.3	11.8	
adjusted ² odds ratio (95 % confidence interval)	ref.	1.43 (0.69–2.95)	1.93 (0.92–4.08)	1.71 (0.76–3.85)	0.35

¹ p-values for one age variable with four levels as obtained from the multivariable logistic regression model, adjusted for screening unit.

² from multivariable logistic regression model, using dummy variables for each age group (reference 50–54 years), adjusted for screening unit.

the less common breast density groups ACR 1 and ACR 4 (ACR 1: 77 % (current study) vs. 100 % [8]; ACR 4: 71 % (current study) vs. 50 % [8]). Given the lower number of cases of cancer, these groups are more susceptible to outliers than ACR 2 and ACR 3 as the main groups. The density groups ACR 1 (5.8 %) and ACR 4 (5.2 %) were seen with a considerably lower prevalence than ACR 2 (44.3 %) and ACR 3 (44.7 %) among 50–69-year-old screening participants in the multicenter approach. The number of cases of cancer (17 (ACR 1) and 56 (ACR 4)) was fundamentally less than the number of cases for ACR 2 (n = 287) and ACR 3 (n = 478) so that the results for individual incidences and program sensitivities in the two density groups with low prevalence must be interpreted with caution. At the same time there was no clear association between age and an increase in the 2-year breast cancer incidence in the less common groups (ACR 1 and ACR 4).

While the age of an individual participant is clearly known, the reproducibility of the visual evaluation of BI-RADS density categories by an interpreting physician is considered limited. Automated computer-assisted breast density determination is more consistent than visual evaluation in studies [14, 15] but is not currently established in the clinical routine.

To avoid underestimation, the higher density category was used for analysis in the present study in the case of a discrepancy in ACR category between the first and second independent reading. On average, there was high agreement regarding categorization between the first reading and the second reading. A discrepancy in ACR categorization per examination of 17 % corresponds with the data provided in the literature [16]. There is currently no gold standard regarding the determination of breast density. It is estimated that one in four women can be expected to be assigned to a different density category in the next examination evaluated by the same interpreting physician and one in three women in the case of a different interpreting physician [16]. A change from density category “not dense” (lipomatous, fibroglandular) to “dense” (heterogeneously dense, extremely dense) or vice versa occurs in 13–19 % of women [16]. The interobserver agreement for categorization is described as moderate (kappa: 0.49–0.56) to highly varied (kappa 0.02–0.72) [16]. On the whole, the results of the present study are based on a plausible distribution of the ACR categories without a disproportionate distortion to higher categories compared to published data (ACR 1: 10 %, ACR 2: 40 %, ACR 3: 40 %, ACR 4: 10 %) [17]. In the future, analyses of the texture of digital mammograms could provide additional

► **Fig. 1** 2-year breast cancer incidence (per 1000 women screened) for screening- plus interval-detected breast cancers (invasive breast cancers plus ductal carcinoma in situ (DCIS)), by ACR category and five-year age group, for women attending subsequent screening rounds.

information for the stratification of breast cancer risk [18]. In principle, analysis of the relationship between the known singular and independent risk factors “age” and “breast density” and the occurrence of breast cancer is of interest [10]. For women older than 50 years in whom a breast cancer-relevant gene mutation has been detected, intensified surveillance including an annual MRI examination is recommended [4]. The significance of MRI for the early detection of a high-risk collective has been shown in a recent study. Among women with the BRCA1 and BRCA2 mutation, 21 % and 36 %, respectively, of all cancers diagnosed in the early detection program were detected by MRI [19].

Based on the risk factor “age”, women 50 years of age or older, even without a family risk factor, are eligible to participate in early detection imaging in the form of mammography screening every two years in Germany [20]. Decreasing rates of advanced tumor stages among participants in subsequent screening rounds [21] and incidences of aggressive subtypes [22] have been reported. The use of substitute or supplementary methods for dense breasts with the goal of improving cancer detection is discussed in the literature. However, there is no evidence of a reduction in breast cancer mortality for any other imaging examination, such as tomosynthesis, sonography, or MRI [23]. Outside the high-risk situation, sonography is currently the recommended method for supplementing mammography in the dense breast (additional cancer detection 4.4 per 1000). However, in early detection it is

associated with a higher rate of additional diagnostic methods and with a low positive predictive value of 3 % to 8.3 % [16]. In the future it would be of interest to identify risk factors for breast cancer or for diagnosis in advanced stages with the goal of modifying risk-adapted screening strategies [14].

A strength of the present scientific analysis is that the selected method allowed combined risk stratification by age and breast density category and that the risk of cancers detected during screening as well as those occurring in the 2-year interval was included. The results are based on a large number of cases using the multicenter approach after program implementation. The results are exclusively from the subsequent screening rounds and are not contaminated by a high percentage of breast cancer cases from initial examinations. The interval cancer data from the screening period 2009 to 2010 were the most current validated data at the time of the evaluation.

A limitation of the study is that the reliability and reproducibility of subjective quantitative breast density determination is limited. Automated validation methods were not available. The aspect of how the diagnosis of breast cancer is distributed in screening and the subsequent interval, which is expressed as the program sensitivity, could not be precisely evaluated for the 16 subgroups resulting from the four 5-year age groups and four ACR categories due to the resulting case number limitations. However, this would be of further interest. As a result of the multi-

► **Fig. 2** Absolute frequency of newly diagnosed breast cancers by screening and during the following interval of two years (n = 838) among 111,456 participants of subsequent screening rounds, by breast density category (ACR 1–4) and five-year age group.

center approach and the small number of cases in the less common density categories, the results do not relate to individual screening units.

Conclusion

In the density categories ACR 2 and ACR 3, the incidences increase significantly up to double as the 5-year age group increases. In contrast, higher breast densities in the youngest age group and higher age in the ACR 1 group do not represent a risk constellation.

CLINICAL RELEVANCE

- The higher breast cancer incidence associated with breast density is modified by the age of the woman.
- Stratification by age and breast density on mammography identifies constellations for a high or low 2-year breast cancer risk.
- Knowledge of risk constellations is important for early detection concepts.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- [1] Robert Koch Institute (ed.) and the Association of Population-based Cancer Registries in Germany (ed.). Cancer in Germany 2011/2012. 10th edition Berlin, 2015
- [2] Broeders M, Moss S, Nystrom L et al. The impact of mammographic screening on breast cancer mortality in Europe: a review of observational studies. *J Med Screen* 2012; 19: 14–25
- [3] Lauby-Secretan B, Scoccianti C, Loomis D et al. Breast cancer screening-Viewpoint of the IARC working group. *N Engl J Med* 2015; 373: 2353–2358
- [4] Bick U. Intensified surveillance for early detection of breast cancer in high-risk patients. *Breast Care* 2015; 10: 13–20
- [5] www.krebsinformationsdienst.de/tumorarten/brustkrebs/brustkrebsrisiken-uebersicht.php
- [6] Boyd NF, Guo H, Martin LJ et al. Mammographic density and the risk and detection of breast cancer. *N Engl J Med* 2007; 356: 227–236
- [7] Pinsky RW, Helvie MA. Mammographic breast density: effect on imaging and breast cancer risk. *J Natl Compr Canc Netw* 2010; 8: 1157–1164
- [8] Weigel S, Heindel W, Heidrich J et al. Digital mammography screening: sensitivity of the programme dependent on breast density. *Eur Radiol* 2017; 27: 2744–2751
- [9] Checka CM, Chun JE, Schnabel FR et al. The relationship of mammographic density and age: implications for breast cancer screening. *Am J Roentgenol* 2012; 198: W292–W295
- [10] Wengert GJ, Helbich TH, Kapetas P et al. Density and tailored breast cancer screening: practice and prediction – an overview. *Acta Radiol Open* 2018; 7: doi:2058460118791212
- [11] Anlage 9.2 BMV-Ä/EKV Bundesmantelvertrag Anlage 9.2. Versorgung im Rahmen des Programmes zur Früherkennung von Brustkrebs durch

Mammografie-Screening. 2014 www.kbv.de/html/bundesmantelvertrag.php

- [12] American College of Radiology. Breast imaging reporting and data system: BI-RADS atlas. 4th edn Reston: American College of Radiology, VA 2003
- [13] Lee CI, Chen LE, Elmore JG. Risk-based Breast Cancer Screening: Implications of Breast Density. *Med Clin North Am* 2017; 101: 725–741
- [14] Cappello NM, Richetelli D, Lee CI. The Impact of Breast Density Reporting Laws on Women's Awareness of Density-Associated Risks and Conversations Regarding Supplemental Screening With Providers. *J Am Coll Radiol* 2019; 16: 139–146
- [15] Holland K, van Zelst J, den Heeten GJ et al. Consistency of breast density categories in serial screening mammograms: a comparison between automated and human assessment. *Breast* 2016; 29: 49–54
- [16] Melnikow J, Fenton JJ, Whitlock EP et al. Supplemental Screening for Breast Cancer in Women With Dense Breasts: A Systematic Review for the U.S. Preventive Service Task Force. [Internet]. Rockville (MD): Agency for Healthcare Research and Quality (US) 2016
- [17] Sickles EA, D'Orsi CJ, Bassett LW et al. ACR BI-RADS® Mammography. In: ACR BI-RADS® Atlas, Breast Imaging Reporting and Data System. Reston: American College of Radiology; 2013
- [18] Wanders JOP, van Gils CH, Karssemeijer N et al. The combined effect of mammographic texture and density on breast cancer risk: a cohort study. *Breast Cancer Res* 2018; 20: 36
- [19] Bick U, Engel C, Krug B et al. High-risk breast cancer surveillance with MRI: 10-year experience from the German consortium for hereditary breast and ovarian cancer. *Breast Cancer Res Treat* 2019; 175: 217–228
- [20] Bundesministerium für Umwelt, Naturschutz und nukleare Sicherheit Bekanntmachung der wissenschaftlichen Bewertung des Bundesamtes für Strahlenschutz gemäß § 84 Absatz 3 des Strahlenschutzgesetzes: Brustkrebsfrüherkennung mittels Mammographie vom 16. Dezember 2018. <https://www.bundesanzeiger.de>
- [21] Weigel S, Heindel W, Heidrich J et al. Reduction of Advanced Breast Cancer Stages at Subsequent Participation in Mammography Screening. *Fortschr Röntgenstr* 2016; 188: 33–37
- [22] Prange A, Bokhof B, Polzer P et al. Higher Detection Rates of Biologically Aggressive Breast Cancers in Mammography Screening than in the Biennial Interval. *Fortschr Röntgenstr* 2019; 191: 130–136
- [23] Leitlinienprogramm Onkologie (Deutsche Krebsgesellschaft, Deutsche Krebshilfe, AWMF). S3-Leitlinie Früherkennung, Diagnose, Therapie und Nachsorge des Mammakarzinoms. Version 4.0, 2017 AWMF Registernummer: 032–045OL, <http://www.leitlinienprogramm-onkologie.de/leitlinien/mammakarzinom/>