Rendezvous technique assisted by anchor device: a new trick for endoscopic treatment of fistulas in the digestive tract

► **Fig. 1** A hydrophilic guidewire passed through the fistula from the vagina to the colon and grasped with forceps.

► **Fig. 2** A colonoscope with an over-the-scope clip mounted on it was inserted into the colon and the guidewire was grasped with an anchor device.

► **Fig. 3** The wire was extracted from the vaginal side while the colonoscope was advanced up to the colonic orifice of the fistula using a rendezvous technique.

philic guidewire was therefore passed through the fistula from the vagina to the colon and grasped with a forceps (► **Fig. 1**). A colonoscope with an over-the-scope clip (OTSC) mounted onto it was used to grasp the guidewire with an anchor device (► **Fig. 2**) and, using a rendezvous technique, we extracted the guidewire from the vaginal side as the colonoscope was advanced up to the colonic orifice of the fistula (► **Fig. 3**). Finally, the OTSC clip was released (► **Video 1**). Following this procedure, the patient had complete remission of her symptoms. The OTSC device was initially designed as a mechanical method of achieving hemostasis and for the closure of small continuity solutions within the digestive tract [1]. This device provides a more lasting effect than a conventional clip owing to its higher compression force [2]. The OTSC is a useful tool in the management of digestive tract continuity solutions, although its effectiveness is greater in acute perforations and leaks [3]. On occasions, it can be difficult to identify a fistulous orifice within the digestive tract, meaning it is not possible to place an OTSC. In such cases, the instillation of methylene blue can serve as a guide to locate the colonic orifice of the fistulous tract [4]; however, this technique is not always effective. We describe an interesting alternative technique for the location of a fistulous orifice in the colonic lu-

A 50-year-old woman with a past history of total abdominal hysterectomy presented with pneumaturia. Barium enema showed a fistulous tract between the vagina and the sigmoid colon. It was not possible to identify the fistulous tract

during a colonoscopy; however, the tract of the fistula could be seen by placing a gastroscope into the vagina. Methylene blue was instilled through the tract but the entrance to the tract in the colon was still not clearly identified. A hydro-

Video 1 Steps in the closure of a colonic fistula to the vagina: the fistulous tract in the sigmoid colon is difficult to identify; a guidewire is placed through the fistula from the vagina; the guidewire is grasped with an anchor device; the colonoscope is positioned at the fistulous orifice using a rendezvous technique; an over-the-scope clip is placed.

men when the fistula runs between two structures that can be endoscopically instrumented.

Endoscopy_UCTN_Code_TTT_1AQ_2AG

Competing interests

The authors declare that they have no conflict of interest.

The authors

Victor Abecia¹, Manuel Alejandro Mahler¹, Sebastian Duran¹, Dante Manazzoni¹, Juan Pizzala¹, Guido Villa-Gómez Roig², Mariano Marcolongo¹

- 1 Gastroenterology Department, Hospital Italiano de Buenos Aires, Buenos Aires Argentina
- 2 Unidad de endoscopia digestiva, Unigastro, La Paz, Bolivia

Corresponding author

Manuel Alejandro Mahler, MD
Gastroenterology Department, Hospital Italiano de Buenos Aires, Perón 4190, Zip code: C1199ABB, Buenos Aires, Argentina
manuel.mahler@hospitalitaliano.org.ar

References

- [1] Kirschniak A, Kratt T, Stüker D et al. A new endoscopic over-the-scope clip system for treatment of lesions and bleeding in the GI tract: first clinical experiences. *Gastrointest Endosc* 2007; 66: 162–167
- [2] Von Renteln D, Vassiliou MC, Rothstein R et al. Randomized controlled trial comparing endoscopic clips and over-the-scope clips for closure of natural orifice transluminal endoscopic surgery gastrotomies. *Endoscopy* 2009; 41: 1056–1061
- [3] Haito-Chavez Y, Law JK, Kratt T et al. International multicenter experience with an over-the-scope clipping device for endoscopic management of GI defects (with video). *Gastrointest Endosc* 2014; 80: 610–622
- [4] Gaj F, Andreuccetti J, Trecca A et al. Identification of internal fistulous orifice: evolution of methylene blue technique with a mini-probe. *Clin Ter* 2012; 163: 57–60

Bibliography

DOI <https://doi.org/10.1055/a-1125-5703>
Published online: 27.3.2020
Endoscopy 2020; 52: E360–E361
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>