Recanalization of an obstructive pancreaticojejunal anastomosis with direct visualization by using antegrade peroral pancreatoscopy

► **Fig. 1** Contrast-enhanced computed tomography (coronal views) showing the dilated main pancreatic duct of the remnant pancreas (arrow).

► **Fig. 3** A 6-mm fully covered self-expandable metallic stent was inserted across the pancreaticogastrostomy.

A 60-year-old man came to our hospital complaining of upper abdominal pain possibly due to stenosis of a pancreaticojejunal anastomosis with upstream dilation of the main pancreatic duct (► **Fig. 1**). Because an endoscopic transluminal approach via the afferent loop failed, we performed endoscopic ultrasound (EUS)-guided pancreatic drainage with a 19-gauge needle (EZ Shot 3 Plus; Olympus Co., Tokyo, Japan). However, no contrast medium flowed out of the dilated main pancreatic duct to the jeju-

► **Fig. 2** Endoscopic ultrasound-guided pancreatic drainage. **a** The dilated main pancreatic duct (arrow) was punctured with a 19-gauge needle under endoscopic ultrasound guidance. **b** From fluoroscopy, no contrast medium flowed out of the dilated pancreatic duct (arrow-head).

► **Video 1** Recanalization of the stenosis of a pancreaticojejunal anastomosis under direct visualization by using antegrade peroral pancreatoscopy via endoscopic ultrasound-guided pancreaticogastrostomy.

num (► **Fig. 2**), and a 0.025-inch guidewire could not be inserted across the anastomosis. A 7-Fr plastic stent was exchanged, 1 month later, for a 6-mm fully covered self-expandable metallic stent (Niti-S Biliary S-type Stent, Century Medical Co., Ltd., Tokyo, Japan) across the pancreaticogastrostomy to perform per-

oral pancreatoscopy (POPS) (► **Fig. 3**). A SpyGlass DS system (Boston Scientific Co., Marlborough, Massachusetts, USA) was used to perform POPS to visualize the anastomosis from the inside of the main pancreatic duct (► **Video 1**). We found the duct completely obstructed at the anastomotic site and covered with

► **Fig. 4** Peroral pancreatoscopy (POPS) using a SpyGlass DS system. **a** POPS revealed complete obstruction of the main pancreatic duct at the anastomotic site, which was covered with fibrotic tissues. **b** The fibrotic tissues were partially broken by poking repeatedly with a guidewire. **c** A guidewire could be passed through the anastomosis. **d** After the anastomosis was dilated by use of a dilator and balloon catheter along the guidewire, POPS images confirmed that the anastomosis site was adequately dilated.

fibrotic tissues (► **Fig. 4**). It was difficult to break through this obstruction even with POPS guidance. However, repeated poking with a guidewire partially broke the fibrotic tissues and a guidewire could finally be passed through the anastomosis. After dilation of this anastomosis using a 7-Fr catheter and a 6-mm balloon catheter, contrast medium immediately flowed from the main pancreatic duct to the jejunum. No procedure-related adverse events were observed, and the abdominal symptoms improved after treatment.

Although the efficacy of EUS-guided pancreatic drainage for stenosis of the pancreaticojejunal anastomosis has been described [1,2], the procedure is still challenging. Recently, the usefulness of cholangioscopy for stenosis of the bilioenteric anastomosis has been reported [3,4]. Therefore, direct visualization using POPS via EUS-guided pancreatico-

gastrostomy appears to be a promising alternative method if fluoroscopic interventions have failed.

Endoscopy_UCTN_Code_TTT_1AR_2AG

Competing interests

The authors declare they have no conflict of interest.

The authors

Yujiro Kawakami^{1,2}, Shinsuke Koshita¹, Yoshihide Kanno¹, Takahisa Ogawa¹, Toji Murabayashi¹, Hiroshi Nakase², Kei Ito¹

¹ Department of Gastroenterology, Sendai City Medical Center, Sendai, Japan

² Department of Gastroenterology and Hepatology, Sapporo Medical University School of Medicine, Sapporo, Japan

Corresponding author

Yujiro Kawakami, MD

Department of Gastroenterology, Sendai City Medical Center, 5-22-1, Tsurugaya, Miyagino-ku, Sendai 9830824, Japan
Fax: +81-22-252-9431
yujiro.kawakami@gmail.com

References

- [1] Matsunami Y, Itoi T, Sofuni A et al. Evaluation of a new stent for EUS-guided pancreatic duct drainage: long-term follow-up outcome. *Endosc Int Open* 2018; 6: E505–E512
- [2] Ogura T, Nishioka N, Yamada M et al. Two-step endoscopic ultrasound-guided rendezvous technique combined with antegrade electrohydraulic lithotripsy for a huge pancreatic duct stone. *Endoscopy* 2019; 51: E149–E150
- [3] Fujii Y, Koshita S, Ito K. Percutaneous transhepatic cholangioscopy using SpyGlassDS for an anastomotic stenosis after choledochojunostomy. *Dig Endosc* 2018; 30: 806–807
- [4] Hakuta R, Kogure H, Nakai Y et al. Successful guidewire placement across hilar biliary stricture after decreased donor liver transplantation using new digital cholangioscopy. *Endoscopy* 2018; 50: E54–E56

Bibliography

DOI <https://doi.org/10.1055/a-1133-4304>
Published online: 27.3.2020
Endoscopy 2020; 52: E376–E377
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

ENDOSCOPY E-VIDEOS
<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>