Successful endoscopic treatment of an appendicocutaneous fistula using endoloops

► **Fig. 1** Abdominal wall radiograph showing the abdominal wall fistula connected with the ileocecal region.

► **Video 1** Appendicocutaneous fistula treated using endoscopic endoloops.

An appendicocutaneous fistula is a rare complication of appendectomy [1]. Moreover, the fistula can persist [2]. A 52-year-old man presented to our department with a 3-month history of purulent abdominal wall sinus that developed after appendectomy 3 months ago. Abdominal wall radiography revealed an abdominal wall fistula connecting with the ileocecal region (► **Fig. 1**). Under colonoscopy, we identified the swollen and purulent appendix stump with black stitching residue (► **Video 1**). After repeated washing of the appendix stump (► **Video 1**), milky pus remained around the stitching residue (► **Fig. 2**). Using grasping forceps, we easily removed the stitching residue (► **Video 1**). Saline solution with methylene blue was injected from the ostium of the abdominal wall fistula, and the inlet of the fistula was detected at the appendix stump (► **Fig. 3**). Normal saline was then injected repeatedly from the abdominal outlet of the sinus to wash the fistula (► **Video 1**). After washing, we released a nylon ring into the ileocecal region (► **Video 1**). Eight clips were used to fasten the nylon ring around the inlet of the fistula at the appendix stump (► **Video 1**). We then tightened

► **Fig. 2** Milky pus around the stitching residue.

► **Fig. 3** Inlet of the fistula at the appendix stump.

► **Fig. 4** Closing of the fistula inlet using endoloops.

► **Fig. 5** Inlet of the fistula healing 2 weeks after treatment.

and released the nylon ring (► **Fig. 4**). Methylene blue dye was again injected from the abdominal wall sinus outlet (► **Video 1**); the dye was refluxed back, and the ileocecal region did not show methylene blue. We found two outlets of the fistula (► **Video 1**). Two drainage tubes were placed at the outlets (► **Video 1**). Two weeks later, we noted that the inlet and the outlets of the fistula were healing (► **Fig. 5**). The use of endoscopic endoloops is a new strategy to promote healing of an appendicocutaneous fistula without surgery.

Endoscopy_UCTN_Code_CPL_1AJ_2AG

Acknowledgments

This article was supported by the Projects of the Bureau of Science and Technology, Nanchong City (grant nos. 18SXHZ0109, 18SXHZ0466).

Competing interests

The authors declare that they have no conflict of interest.

The authors

Zhang Tao¹, Dongbing Zhou², Xiaosan Hu¹, Mingyang Ren², Yiwen Yu¹, Xiaoqing Zhou¹

- 1 Department of Gastroenterology, Nanchong Central Hospital, Nanchong City, Sichuan, China
- 2 Department of Gastrointestinal Surgery, Nanchong Central Hospital, Nanchong City, Sichuan, China

Corresponding author

Xiaoqing Zhou, MD

Department of Gastroenterology, Nanchong Central Hospital, Nanchong City, Sichuan, 637000, China
Fax: +86-10-67788052
zhangtao195932@163.com

References

- [1] Brünagel G, Decker P, Hirner A. Delayed appendico-cutaneous fistula – a rare complication of simple abdominal drainage. *Zentralbl Chir* 1996; 121: 67–69
- [2] Shamim M, Haider SA, Iqbal SA. Persistent appendiceal faecal fistula following a complicated open appendicectomy. *J Pak Med Assoc* 2009; 59: 181–183

Bibliography

DOI <https://doi.org/10.1055/a-1134-4630>

Published online: 15.4.2020

Endoscopy 2020; 52: E383–E384

© Georg Thieme Verlag KG

Stuttgart · New York

ISSN 0013-726X

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>