Indeterminate biliary stricture treated by antegrade cholangioscopy through an endoscopic ultrasound-guided hepaticojejunostomy

► **Fig. 1** Abdominal computed tomography scan showing bile duct dilatation.

► **Fig. 2** Magnetic resonance cholangiopancreatography showing bile duct dilatation and a filling defect in the papillary area.

Indeterminate biliary strictures represent a clinical challenge [1,2]. Endoscopic retrograde cholangiopancreatography (ERCP) with brushings and transpapillary biopsies have limitations in terms of their sensitivity and specificity [3,4]. Cholangioscopy plays a relevant role in this field with a high sensitivity for the macroscopic appearance [5].

A 60-year-old woman, who had undergone total gastrectomy with Roux-en-Y reconstruction for gastric adenocarcinoma (pT2 N0 M0) 4 years previously, presented to our ambulatory unit with upper abdominal pain and alteration of liver

function tests. An abdominal computed tomography scan (► **Fig. 1**) and magnetic resonance cholangiopancreatography (MRCP) (► **Fig. 2**) showed bile duct dilatation and a distal stricture of the common bile duct. However, an endoscopic ultrasound (EUS) approach

was not feasible or effective because of her altered anatomy.

A first attempt to reach the papillary area and perform a subsequent ERCP using a pediatric colonoscope was unsuccessful owing to the length and angulations of the jejunal loop. Therefore, a hepatico-

► **Fig. 3** Fluoroscopic images showing: **a** an endoscopic ultrasound-guided hepaticojejunostomy being created; **b** the guidewire that had been passed through the transjejunal access and through the papillary area retrogradely being recovered with the pediatric colonoscope.

▶ Video 1 Antegrade cholangioscopy through an endoscopic ultrasound-guided hepaticojunostomy to treat a biliary indeterminate stricture in a patient with Roux-en-Y reconstruction.

References

- [1] Singh A, Gelrud A, Agarwal B. Biliary strictures: Diagnostic considerations and approach. *Gastroenterol Rep (Oxf)* 2015; 3: 22–31
- [2] Kato M, Onoyama T, Takeda Y et al. Peroral cholangioscopy-guided forceps biopsy and endoscopic scraper for the diagnosis of indeterminate extrahepatic biliary stricture. *J Clin Med* 2019; 8: 1–11
- [3] Ponchon T, Gagnon P, Berger F et al. Value of endobiliary brush cytology and biopsies for the diagnosis of malignant bile duct stenosis: results of a prospective study. *Gastrointest Endosc* 1995; 42: 565–572
- [4] Burnett AS, Calvert TJ, Chokshi RJ. Sensitivity of endoscopic retrograde cholangiopancreatography standard cytology: 10-y review of the literature. *J Surg Res* 2013; 184: 304–311
- [5] Navaneethan U, Njei B, Lourdasamy V et al. Comparative effectiveness of biliary brush cytology and intraductal biopsy for detection of malignant biliary strictures: A systematic review and meta-analysis. *Gastrointest Endosc* 2015; 81: 168–176

Bibliography

Endoscopy 2020; 52: 1143–1144
 DOI 10.1055/a-1149-8684
 ISSN 0013-726X
 published online 24.4.2020
 © 2020. Thieme. All rights reserved.
 Georg Thieme Verlag KG, Rüdigerstraße 14,
 70469 Stuttgart, Germany

jejunostomy was performed endoscopically (▶ Fig. 3a). First, a transjejunal hepatic EUS-guided puncture was done using a 19G needle. A guidewire was pushed through the transjejunal access through the papillary area retrogradely and was recovered through the pediatric colonoscope (▶ Fig. 3b). The papilla was cannulated from distance under fluoroscopic guidance (because of a tight angulation at Treitz's ligament) using a 12-mm Fogarty balloon. Cholangiography confirmed homogeneous dilatation of the bile ducts with a distal stricture. The stricture was dilated using a 10-mm pneumatic balloon; brushings and transpapillary biopsies were then performed. A 10-Fr 12-cm biliary plastic stent was left in place to maintain patency of the endoscopic hepaticojunostomy.

Because the cytology was non-diagnostic, antegrade cholangioscopy using the SpyGlassDS system (Boston Scientific Co.) was successfully performed through the endoscopic hepaticojunostomy 2 weeks later (▶ Video 1). The appearance of the stricture was non-malignant; several cholangioscopic biopsies were taken using a SpyBite (Boston Scientific Co.), which later confirmed there was no cellular atypia on histopathology. Finally, a 10-Fr 12-cm biliary plastic stent was positioned through the hepaticojunostomy.

No early adverse events occurred and the patient was discharged the day after the

procedure. At her 1-month follow-up visit, the patient had developed two liver abscesses (S4–S8), which were probably related to bile duct contamination during the cholangioscopy. Percutaneous drainage of the abscesses was performed, and the patient was found to be doing well at her 3-month follow-up visit.

Endoscopy_UCTN_Code_TTT_1AS_2AD

Competing interests

The authors declare that they have no conflict of interest.

The authors

Roberto Rosa, Lorenzo Dioscoridi, Edoardo Forti, Francesco Pugliese, Marcello Cintolo, Giulia Bonato, Massimiliano Mutignani
 Digestive Endoscopy Unit, ASST Niguarda, Milan, Italy

Corresponding author

Massimiliano Mutignani, MD
 Piazza dell'Ospedale Maggiore 3, 20158, Milan, Italy
 Fax: +39-02-64448565
 massimiliano.mutignani@ospedaleniguarda.it

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>