Rectal tonsil: a rare cause of rectal bleeding

► **Fig. 1** Rectal mucosa of a 21-year-old patient with a 4-month history of rectal bleeding. Lymphatic follicle with a prominent germinal center containing tingible body macrophages (H&E, ×10).

► **Fig. 2** Rectal mucosa of the same patient. **a** Lymphatic follicle positive for CD20 (×10); **b** lymphatic follicle with germinal center negative for Bcl2 (×10).

A rectal tonsil is a lymphoid follicular hyperplasia of the rectum and a rare cause of rectal bleeding. As it is a benign lesion of uncertain etiology [1, 2], treatment is reserved for symptomatic patients [2].

We present the case of a 21-year-old woman with normal bowel movements who presented with a history of rectal bleeding over a period of 4 months. Digital rectal examination revealed a rectal mass. Colonoscopy revealed circumferential granular involvement of the rectal mucosal surface (► **Video 1**). Histology showed a dense lymphoid infiltrate and lymphatic follicles (► **Fig. 1**).

Infection [3] and lymphoma were excluded (► **Fig. 2**) and a diagnosis of rectal tonsil was made [4]. Excellent clinical and endoscopic response was achieved after a 1-month course of rectal mesalazine administered twice daily (► **Fig. 3**). Because the lesion is benign in nature, with transformation into lymphoma being the exception [4], treatment was discontinued. The patient continued asymptomatic at 3-month clinical follow-up.

Endoscopy_UCTN_Code_CCL_1AD_2AC

► **Fig. 3** Treatment with rectal mesalazine: appearance of the rectal mucosa before treatment (**a**) and after treatment (**b**).

Acknowledgments

We acknowledge Prof. Dr. Carolina Olano (Hospital de Clínicas Uruguay) for her dedication and support in the publication of this E-Video.

Competing interests

The authors declare that they have no conflict of interest.

The authors

Ignacio Moratorio¹, M. Alejandra Arriola¹, Ximena Pazos¹, Patricia Dorfman², Marta Otero²

- 1 Department of Gastroenterology, Hospital de Clínicas, Facultad de Medicina, Universidad de la República, Montevideo, Uruguay
- 2 Department of Pathological Anatomy, Hospital de Clínicas, Facultad de Medicina, Universidad de la República, Montevideo, Uruguay

Video 1 Colonoscopic appearance of a rectal tonsil before and after treatment with mesalazine.

Bibliography

Endoscopy 2020; 52: E443–E444
DOI 10.1055/a-1158-8882
ISSN 0013-726X
published online 12.5.2020
 © 2020. Thieme. All rights reserved.
 Georg Thieme Verlag KG, Rüdigerstraße 14,
 70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>

Corresponding author

Ignacio Moratorio, MD
 Gregorio Suarez 2825/205, CP 11300
 Montevideo, Uruguay
ignaciomoratorio@gmail.com

References

- [1] Hong JB, Kim HW, Kang DH et al. Rectal tonsil: A case report and literature review. *World J Gastroenterol* 2015; 21: 2563–2567
- [2] Trillo L, Arias M, Iglesias A et al. Amígdala rectal o hiperplasia folicular linfoide del recto. *Radiología* 2014; 56: 370–373
- [3] Cramer SF, Romansky S, Hulbert B et al. The rectal tonsil: a reaction to chlamydial infection? *Am J Surg Pathol* 2009; 33: 483–485
- [4] Farris AB, Lauwers GY, Ferry JA et al. The rectal tonsil: a reactive lymphoid proliferation that may mimic lymphoma. *Am J Surg Pathol* 2008; 32: 1075–1079