Endoscopic submucosal dissection by transnasal endoscope for esophageal cancer with pharyngoesophageal anastomotic stricture after total pharyngo-laryngo-esophagectomy

Patients with squamous cell carcinoma (SCC) of the head and neck often develop secondary esophageal carcinomas. We describe a successful case of endoscopic submucosal dissection (ESD) using a transnasal endoscope to treat a patient with esophageal cancer with a pharyngoesophageal anastomotic stricture after total pharyngo-laryngo-esophagectomy (TPLE).

A 60-year-old man was referred to our division for treatment of esophageal cancer. The patient had undergone TPLE for advanced hypopharyngeal cancer 5 years previously, and endoscopy revealed a flat lesion in the lower third of the esophagus (►Fig. 1 a, b). The patient had a pharyngoesophageal anastomotic stricture after the TPLE that could not be passed using a conventional endoscope (►Fig. 1 c). Endoscopic balloon dilation was not possible because of the presence of a voice prosthesis (►Fig. 1 d). Recently, an improved transnasal endoscope, the EG-L580NW (Fujifilm, Tokyo, Japan), has become available, with a 5.8-mm outer diameter and a 2.4-mm forceps diameter (►Fig. 2 a) [1]. An ultrathin

►Fig. 1 a–d A 60-year-old man referred for treatment of esophageal cancer 5 years after undergoing total pharyngo-laryngo-esophagectomy (TPLE) for advanced hypopharyngeal cancer. **a** Blue laser imaging shows a brownish area in the lower third of the esophagus. **b** After chromoendoscopy using Lugol's iodine, markers were placed 2–3 mm outside of the lesion margins. **c** Pharyngoesophageal anastomotic stricture after the TPLE (arrow). **d** A voice prosthesis had been used to restore the voice (arrow).

►Fig. 2 **a** The EG-L580NW (Fujifilm, Tokyo, Japan) has an instrument channel with a 2.4-mm inner diameter. **b** The Souten ultrathin needle-type device has a diameter of 2.35 mm at the insertion sheath. **c** The Raicho hemostasis forceps has a diameter of 2.3 mm at the insertion sheath.

► **Fig. 3** **a** A caudal incision was performed. **b** Circumferential incisions were made using the Souten. **c** Submucosal dissection was performed. **d** En bloc resection was achieved without causing injury to the muscularis propria.

► **Fig. 4** **a** Resected specimen after iodine staining. **b** Histology shows esophageal squamous cell carcinoma limited to the epithelium.

needle-type device (Souten; Kaneka Medics, Tokyo, Japan) and hemostasis forceps (Raicho; Kaneka Medics, Tokyo, Japan) can be used with this endoscope (► **Fig. 2b,c**) [2]. Thus, we performed ESD using a transnasal endoscope to treat the esophageal cancer (► **Video 1**). Circumferential incisions were made using the Souten with Endo Cut I mode (VIO 300D; Erbe, Tübingen, Germany; effect 2, duration 2, interval 2) (► **Fig. 3a,b**). To control bleeding, the Raicho forceps was used in soft coagulation mode at 50 W. Submucosal dissection was performed using the Souten and swift coagulation mode (effect 3, 30 W) (► **Fig. 3c**). Finally, en bloc resection was achieved without injury to the muscularis propria (► **Fig. 3d**). Histological examination revealed R0 resection of the squamous cell carcinoma, which was limited to the epithelium (► **Fig. 4**).

Esophageal ESD using a transnasal endoscope was useful in treating esophageal cancer with an anastomotic stricture that could not be passed using a conventional endoscope.

Endoscopy_UCTN_Code_TTT_1AO_2AG

Competing interests

The authors declare that they have no conflict of interest.

The authors

Yoshiyasu Kitagawa¹, Takuto Suzuki¹, Kentaro Nakamura^{1,2}, Rino Nankinzan¹, Taketo Yamaguchi³

- 1 Endoscopy Division, Chiba Cancer Center, Chiba, Japan
- 2 Department of Gastroenterology, Toho University Sakura Medical Center, Sakura, Japan
- 3 Department of Gastroenterology, Chiba Cancer Center, Chiba, Japan

Video 1 Endoscopic submucosal dissection by transnasal endoscope for esophageal cancer with pharyngoesophageal anastomotic stricture after total pharyngo-laryngo-esophagectomy.

Corresponding author

Yoshiyasu Kitagawa, MD, PhD
Endoscopy Division, Chiba Cancer Center,
666-2 Nitonacho, Chuo-ku, Chiba, Japan
Fax: +81-43-2628680
ykitagawa@chiba-cc.jp

References

- [1] Suzuki T, Kitagawa Y, Nankinzan R et al. Early gastric cancer diagnostic ability of ultrathin endoscope loaded with laser light source. *World J Gastroenterol* 2019; 25: 1378–1386
- [2] Minato Y, Sakai E, Ohata K. Conscious transnasal hybrid endoscopic submucosal dissection enables safe and painless en bloc resection in elderly patients with early gastric cancer. *VideoGIE* 2019; 4: 157–158

Bibliography

Endoscopy 2020; 52: E445–E447
DOI 10.1055/a-1158-8948
ISSN 0013-726X
published online 12.5.2020
© 2020. Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>