Is artificial intelligence a superior diagnostician in ulcerative colitis?

► **Fig. 1** Colonoscopic image revealing a slightly elevated reddish lesion in the lower rectum (yellow arrows).

► **Fig. 2** Histologic appearance showing an atypical tubular gland with high grade dysplasia in the mucosa.

A 46-year-old woman with a 26-year history of pancolitis visited our hospital regularly. She had remained in remission for the past 5 years and was taking oral mesalamine. Annual endoscopy examination showed a slightly elevated reddish lesion (5 mm) in the lower rectum (► **Fig. 1**). Endocytoscopy (CF-H290ECI; Olympus, Tokyo, Japan) with narrow-band imaging revealed surface microvessels of uniform caliber and arrangement (► **Video 1**). An artificial intelligence-based system (EndoBRAIN; Cybernet Systems, Tokyo, Japan) diagnosed the lesion as a neoplasm. On the basis of biopsy findings, we diagnosed the lesion as being high grade dysplasia (► **Fig. 2**) and performed

► **Video 1** View during endocytoscopy with narrow-band imaging showing surface microvessels of uniform caliber and arrangement, which is subjected to computer diagnosis using a two-class diagnosis (non-neoplastic or neoplastic) with probability.

endoscopic submucosal dissection. After treatment, she was discharged on post-operative day 7, without any complications.

The risk of developing colorectal cancer (CRC) is significantly elevated in patients with ulcerative colitis. Surveillance colonoscopy is usually performed to identify colitis-associated CRC. Endocytoscopy is a novel type of ultramagnification endoscopy that enables microscopic cellular observation, and evaluation of the endocytoscopic microvasculature has been used for the diagnosis of colorectal lesions [1]. The EndoBRAIN system accurately differentiates neoplastic from non-neoplastic lesions in endocytoscopic narrow-band images, when pathological findings are used as the standard [2]. However, its application in the diagnosis of colitis-associated CRC has not previously been reported. Our findings suggest that endocytoscopy can be used to obtain information regarding the endocytoscopic vascular pattern and that EndoBRAIN can help detect colitis-asso-

ciated CRC. Furthermore, colitis-associated CRC is often difficult to diagnose because of the effects of inflammation and the use of EndoBRAIN can help non-expert endoscopists to identify colitis-associated CRC, thereby helping to avoid unnecessary biopsies.

Endoscopy_UCTN_Code_TTT_1AQ_2AB

Competing interests

The authors declare that they have no conflict of interest.

The authors

Shuhei Fukunaga, Yoshio Kusaba, Akihiro Ohuchi, Tsutomu Nagata, Keiichi Mitsuyama, Osamu Tsuruta, Takuji Torimura

Division of Gastroenterology, Department of Medicine, Kurume University School of Medicine, Fukuoka, Japan

Corresponding author

Shuhei Fukunaga, MD

Division of Gastroenterology, Department of Medicine, Kurume University School of Medicine, 67 Asahi-machi, Kurume, Fukuoka 830-0011, Japan

Fax: +81-942-342623

fukunaga_shyuhei@med.kurume-u.ac.jp

References

- [1] Kudo SE, Misawa M, Wada Y et al. Endocytoscopic microvasculature evaluation is a reliable new diagnostic method for colorectal lesions. *Gastrointest Endosc* 2015; 82: 912–923
- [2] Kudo SE, Misawa M, Mori Y et al. Artificial intelligence-assisted system improves endoscopic identification of colorectal neoplasms. *Clin Gastroenterol Hepatol* 2019. doi:10.1016/j.cgh.2019.09.009

Bibliography

Endoscopy 2021; 53: E75–E76

DOI 10.1055/a-1195-1986

ISSN 0013-726X

published online 26.6.2020

© 2020. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>