Endoscopic ultrasound-guided gallbladder drainage by transjejunal lumen-apposing metal stent placement in a patient with mini-gastric bypass

► **Fig. 1** Computed tomography showing a distended gallbladder adherent to the jejunal loop.

Endoscopic ultrasound-guided gallbladder drainage (EUS-GBD) using lumen-apposing electrocautery-enhanced metal stent (EC-LAMS) is a safe and effective treatment option in patients with acute cholecystitis who are unfit for surgery [1]. We present the case of a 64-year-old woman who presented to the emergency department with a grade III cholecystitis [2] (► **Fig. 1**). The patient had dilated cardiomyopathy with an implanted cardioverter-defibrillator, decompensated type II diabetes, and had undergone mini-gastric bypass surgery for obesity control 10 years ago. She was deemed unfit for surgery at this time and endoscopic drainage was proposed [3] (► **Video 1**). EUS revealed an over-distended gallbladder attached to the efferent loop of the gastrojejunal anastomosis. EUS-GBD was therefore performed by transmural placement of an EC-LAMS (Hot Axios, 10×10 mm; Boston Scientific, Marlborough, Massachusetts, USA) (► **Fig. 2**), allowing the release of a significant amount of purulent bile from the gallbladder into the jejunal efferent loop (► **Video 1**). After the procedure, the patient's general status progressively improved (► **Fig. 3**); however, 2 weeks after the procedure, fever persisted and a second endoscopic procedure was planned. Using a duodenoscope, the efferent limb was reached and the LAMS was still in

► **Fig. 2** Graphical representation of transjejunal endoscopic ultrasound-guided gallbladder drainage in a patient with a mini-gastric bypass.

place draining purulent bile; however, the stent lumen was partially clogged by biliary debris. Under endoscopic guidance, a stone extraction balloon catheter was placed into the stent and the debris was removed. A 7-Fr nasogallbladder drainage tube was then placed through the stent (► **Fig. 4**).

In the following days, the fever resolved and inflammatory markers returned to normal levels. The nasogallbladder drainage tube was removed and the patient

was discharged home. At 3-month follow-up, the patient remained asymptomatic with improvement in general status, allowing surgery to proceed. The EC-LAMS was removed using a gastroscope and snare without complications, and an elective cholecystectomy was planned.

Endoscopy_UCTN_Code_TTT_1AS_2AD

► **Fig. 3** Computed tomography demonstrating the correct positioning of the electrocautery-enhanced lumen-apposing metal stent between the jejunal effluent loop and the gallbladder lumen.

► **Fig. 4** Computed tomography documenting nasogallbladder drainage tube placed through the metal stent into the gallbladder lumen.

References

- [1] Krishnamoorthi R, Irani S. Endoscopic ultrasound-guided gallbladder drainage using lumen-apposing metal stents. *Tech Innov Gastrointest Endosc* 2020; 22: 19–23
- [2] Yokoe M, Hata J, Tadahiro T et al. Tokyo Guidelines 2018: diagnostic criteria and severity grading of acute cholecystitis (with videos). *J Hepatobiliary Pancreat Sci* 2018; 25: 41–54
- [3] Kohji O, Kenji S, Tadahiro T et al. Tokyo Guidelines 2018: flowchart for the management of acute cholecystitis. *J Hepatobiliary Pancreat Sci* 2018; 25: 55–72

Bibliography

Endoscopy 2021; 53: E81–E82
 DOI 10.1055/a-1196-1201
 ISSN 0013-726X
 published online 26.6.2020
 © 2020. Thieme. All rights reserved.
 Georg Thieme Verlag KG, Rüdigerstraße 14,
 70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>

► **Video 1** Endoscopic ultrasound-guided gallbladder drainage by transjejunal placement of a lumen-apposing metal stent.

Competing interests

The authors declare that they have no conflict of interest.

The authors

Annalisa Cappello¹, Alessia Santini², Ivano Biviano², Elena Gianni¹, Natalia Simonetti¹, Sandro Giannotti¹, Raffaele Macchiarelli²

¹ Gastroenterology Unit, A.O.U.S. Policlinico S. Maria alle Scotte, Siena, Italy

² Biliopancreatic Unit, A.O.U.S. Policlinico S. Maria alle Scotte, Siena, Italy

Corresponding author

Annalisa Cappello, MD

Gastroenterology Unit, A.O.U.S. Policlinico S. Maria alle Scotte, Viale Bracci, 53100, Siena, Italy
annalisacappello@yahoo.it