Endoscopic ultrasound-guided angiotherapy in bleeding gastrointestinal stromal tumors with coil deployment and cyanoacrylate injection

► **Fig. 1** Endoscopic view of the ulcerated bleeding GI stromal tumor (GIST).

Hemorrhage is the most frequent (30–40%) and dangerous complication of gastrointestinal stromal tumors (GISTs), with surgery being the treatment of choice and emergency surgery having a significantly higher risk than elective procedures [1]. Conventional endoscopic treatment of tumoral lesions is challenging and efficacy limited because of the complex neoangiogenesis of these lesions, which are beyond the reach of endoscopic contact procedures [2]. Cases of endoscopic ultrasound (EUS)-guided therapy of bleeding GISTs with injection of cyanoacrylate (CYA) [3,4] or ethanol [5] have been published. To the best of our knowledge, EUS-guided coil deployment has not been reported in this setting. We present two cases of bleeding GISTs successfully treated with EUS-guided combined coil deployment and further injection of CYA targeting the feeding artery (► **Video 1**).

Case 1: a 72-year-old man was admitted for melena and hemodynamic instability. An ulcerated fundic mass with a necrotic center and a 5-mm intratumoral artery was observed (► **Fig. 1**). EUS-guided injection of CYA (Histoacryl; B. Braun, Melsungen, Germany) into the feeding artery (► **Fig. 2**) with a 22-gauge needle was the initial approach. However, a spurting

► **Video 1** Endoscopic ultrasound (EUS)-guided angiotherapy in two patients with bleeding GI stromal tumors with EUS-guided coil deployment and cyanoacrylate (CYA) injection targeting the feeding artery.

► **Fig. 2** Endoscopic ultrasound (EUS) image displaying the feeding artery of the bleeding GI stromal tumor (GIST) (arrows) shown in ► **Fig. 1**.

bleed into the tumor cavity after puncture was observed (► **Fig. 3**). Hemostasis was achieved by deploying a 6-mm × 20-

cm coil (Nester; Cook Medical, Limerick, Ireland) and injecting 2 mL of CYA with a 19-gauge needle (► **Fig. 4**).

► **Fig. 3** Spurting bleed (arrows) after the endoscopic ultrasound (EUS)-guided puncture with a 22-gauge needle.

► **Fig. 4** Endoscopic ultrasound (EUS)-guided deployed coil (arrows) with a 19-gauge needle.

Case 2: a 79-year-old woman was admitted for hematemesis and hypovolemic shock. An ulcerated fundic mass with a 4-mm artery was observed. EUS-guided therapy with a 19-gauge needle was performed employing a 6-mm × 14-cm coil, followed by injection of 2 mL of CYA. The clinical outcome in both patients was uneventful, and scheduled surgery took place 4 weeks later. EUS-guided combined angiotherapy targeting the culprit artery of bleeding GISTs maintained hemostasis until scheduled surgery. Moreover, although the follow-up in these two patients was short, EUS-guided angiotherapy may be a therapeutic option in patients with bleeding GISTs deemed unfit for surgery.

Endoscopy_UCTN_Code_TTT_1AS_2AG

Competing interests

The authors declare they have no conflict of interest.

The authors

Rafael Romero-Castro^{1,2}, Victoria Alejandra Jimenez-Garcia^{1,2}, Atsushi Irisawa³, Pedro Hergueta-Delgado¹, Antonio Garrido-Serrano¹, Reyes Aparcero-Lopez¹, Angel Caunedo-Alvarez¹

- 1 Virgen Macarena University Hospital, Endoscopy Unit, Gastroenterology Division, Seville, Spain
- 2 Vithas Nisa Seville Hospital, Endoscopy Unit, Seville, Spain
- 3 Department of Gastroenterology, Dokkyo Medical University School of Medicine, Tochigi, Japan

Corresponding author

Rafael Romero Castro, MD
Virgen Macarena University Hospital,
Endoscopy Unit, Gastroenterology Division,
Calle Dr. Fedriani, 3, 41009 Seville, Spain
Fax: +34 955 00 88 05
rafaromecas@hotmail.com

References

- [1] Liu Q, Kong F, Zhou J et al. Management of hemorrhage in gastrointestinal stromal tumors: a review. *Cancer Manag Res* 2018; 10: 735–743
- [2] Venezia G, Asnaghi G, Manca A et al. Successful use of Hemospray as a bridge to surgery in life-threatening bleeding from a gastrointestinal stromal tumor of the stomach. *Endoscopy* 2016; 48: E232–E233
- [3] Levy MJ, Wong Kee Song LM, Farnell MB et al. Endoscopic ultrasound (EUS)-guided angiotherapy of refractory gastrointestinal bleeding. *Am J Gastroenterol* 2008; 103: 352–359

- [4] Kumbhari V, Gondal B, Okolo I et al. Endoscopic ultrasound-guided angiotherapy of a large bleeding gastrointestinal stromal tumor. *Endoscopy* 2013; 45: E326–E327
- [5] Valdivielso Cortazar E, Fernandez-Urien I, Vila Costas JJ et al. Endoscopic ultrasound-guided therapy of a gastrointestinal stromal tumor (GIST). *Endoscopy* 2015; 47: E262–E263

Bibliography

Endoscopy 2021; 53: E124–E125
DOI 10.1055/a-1216-0634
ISSN 0013-726X
published online 24.7.2020
© 2020. Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>