Transoral endoscopic ultrasound-guided fine-needle biopsy of a tumor of the parapharyngeal space

The parapharyngeal space is a pyramid-shaped space located between the base of the skull and the hyoid bone, lateral to the naso-oropharynx and medial to the jaw [1,2]. It contains the deep lobe of the parotid gland, cranial nerves IX–XII, the internal jugular vein, and the carotid artery [1]. A variety of benign and malignant tumors arise in the parapharyngeal space, the most common being of salivary gland and neurogenic origin [1]. Because of its deep location and concerns about damaging adjacent structures, the parapharyngeal space is difficult to access for biopsy [1–3]. Percutaneous, transoral, or transnasal approaches have been used; however, sampling may be challenging even under imaging guidance [2–5]. Here we report for the first time a technique of transoral biopsy using a flexible gastrointestinal echoendoscope.

A 41-year-old man with a remote history of mucoepidermoid carcinoma of the left parotid gland presented with a tumor of the right parapharyngeal space (►Fig. 1). A multidisciplinary tumor board recommended biopsy; however, this was deemed difficult due to the tumor location. After discussion with a gastroenterologist experienced in endoscopic ultrasound (EUS), a decision was made for biopsy under EUS guidance. With the patient in the left lateral position and under intravenous sedation, a flexible echoendoscope (Olympus GF-UCT180) was introduced in a standard manner into the oral cavity and torqued clockwise. After passing the palatopharyngeal fold, the EUS transducer was gently wedged against the right lateral wall of the pharynx (►Fig. 2). In this position the tumor was easily identified on the EUS image. Three passes with a 22-gauge Acquire needle (Boston Scientific) were performed (►Fig. 3, ►Video 1). The specimen was processed for histological evaluation, which revealed pleomorphic adenoma of the deep lobe of the right parotid gland (►Fig. 4). The patient was dis-

► **Fig. 1** Magnetic resonance images showing a tumor (TU) of the right parapharyngeal space. **a** Unenhanced T1-weighted coronal image. The tumor is located between the wall of the nasopharynx, lateral (lpm) and medial pterygoid muscles (mpm), and ramus of the mandible. It is 34 mm × 25 mm in size, has well-delineated borders, and a homogeneous low-intensity T1 signal. **b** Gadolinium-enhanced transverse image. The tumor has an inhomogeneous enhancement pattern. A connection to the deep lobe of the right parotid gland is visible (asterisk).

► **Video 1** Successful transoral endoscopic ultrasound-guided biopsy of a tumor of the parapharyngeal space (pleomorphic adenoma of the deep lobe of the parotid gland).

charged 48 h later after an uneventful course. The tumor was resected 4 weeks later. Surgical pathology confirmed the diagnosis of pleomorphic adenoma of the deep lobe of the right parotid gland.

Endoscopy_UCTN_Code_TTT_1AS_2AC

► **Fig. 2** Schematic drawing of the anatomy of the mouth and throat (midsagittal plane), showing the position of the echoendoscope during endoscopic ultrasound (EUS) examination and fine-needle biopsy of the tumor.

► **Fig. 3** EUS image of a tumor of the right parapharyngeal space located between the pharyngeal wall, the medial pterygoid muscle (mpm), and the ramus of the mandible. The tumor (TU) is 34 mm × 28 mm in size, has well-delineated borders and a hypoechoic, homogeneous echo pattern. A needle passing through the pharyngeal wall into the tumor is visible.

Competing interests

The authors declare that they have no conflict of interest.

The authors

Marcin Polkowski^{1,2}, Jacek Lenartowicz³, Jakub Zwoliński³, Jakub Pałucki⁴, Andrzej Mróz^{5,6}, Kamil Sokół⁶, Jarosław Reguła^{1,2}

- 1 Department of Gastroenterological Oncology, Maria Skłodowska-Curie National Research Institute of Oncology, Warsaw, Poland
- 2 Department of Gastroenterology, Hepatology and Clinical Oncology, Center of Postgraduate Medical Education, Warsaw, Poland
- 3 Department of Head and Neck Cancer, Maria Skłodowska-Curie National Research Institute of Oncology, Warsaw, Poland
- 4 Department of Radiology, Maria Skłodowska-Curie National Research Institute of Oncology, Warsaw, Poland
- 5 Department of Pathology, Center of Postgraduate Medical Education, Warsaw, Poland
- 6 Department of Pathology and Laboratory Medicine, Maria Skłodowska-Curie National Research Institute of Oncology, Warsaw, Poland

► **Fig. 4** Specimen obtained by EUS-guided fine-needle biopsy under 100× magnification. Typical appearance of a pleomorphic adenoma of the parotid gland with epithelial, myoepithelial, and stromal components. Hematoxylin and eosin stain.

Corresponding author

Marcin Polkowski, MD, PhD

Department of Gastroenterological
Oncology, Maria Skłodowska-Curie
National Research Institute of Oncology,
Roentgena 5, 02-781 Warsaw, Poland
mp.polkowski@gmail.com

References

- [1] Lopez F, Suarez C, Vander Poorten V et al. Contemporary management of primary parapharyngeal space tumors. *Head Neck* 2019; 41: 522–535
- [2] Riskalla A, Arora A, Vaz F et al. Novel use of ultrasound-guided endo-cavitary probe to evaluate an impalpable parapharyngeal mass. *J Laryngol Otol* 2010; 124: 328–329
- [3] Abbas JR, Hamlett KEL, de Carpentier J. Image-guided transnasal endoscopic fine nee-

dle aspiration or biopsy of parapharyngeal space tumours. *J Laryngol Otol* 2018; 132: 1026–1028

- [4] Chen R, Cai Q, Liang F et al. Oral core-needle biopsy in the diagnosis of malignant parapharyngeal space tumors. *Am J Otolaryngol* 2019; 40: 233–235
- [5] Arnason T, Hart RD, Taylor SM et al. Diagnostic accuracy and safety of fine-needle aspiration biopsy of the parapharyngeal space. *Diagn Cytopathol* 2012; 40: 118–123

Bibliography

Endoscopy 2021; 53: E145–E147

DOI 10.1055/a-1216-9928

ISSN 0013-726X

published online 5.8.2020

© 2020. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>