Establishment of Clinical and Lab Algorithms for the Identification of Carriers of Mutations in *CYP21A2* – A Study of 365 Children and Adolescents

Authors

J. Meinel¹, T. Haverkamp¹, F. Wünsche¹, A. Richter-Unruh^{1,2}

Affiliations

- 1 MVZ Dr. Eberhard & Partner Dortmund (ÜBAG), Dortmund Germany
- 2 Ruhr-University Bochum, University Children's Hospital Bochum, St. Josef- Hospital, Department of Pediatric Endocrinology, Bochum Germany

Key words

CYP21A2 gene, ACTH-test, Congenital adrenal hyperplasia, 17OHP values

received 12.03.2020

revised 16.06.2020

accepted 08.07.2020

published online 24.08.2020

Bibliography

Exp Clin Endocrinol Diabetes 2021; 129: 492–499

DOI 10.1055/a-1217-7169

ISSN 0947-7349

© 2020. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany

Correspondence

Prof. Annette Richter-Unruh
Childrens Hospital, Ruhr University Bochum,
Alexandrinenstr. 5
44801 Bochum
Germany
Tel.: 01737280863
richter-unruh@web.de

Supplementary Material for this article is available online at <http://www.thieme-connect.de/products>.

ABSTRACT

Background Mutations of *CYP21A2* encoding 21-hydroxylase are the most frequent cause of congenital adrenal hyperplasia (CAH) and are associated either with elevated basal or ACTH-stimulated levels of 17-hydroxyprogesterone (17OHP) in blood. **Objective** The study objective was to identify the most suitable of 12 different test algorithms and appropriate cut-off levels for that test to recognize patients with non-classical congenital adrenal hyperplasia (NCCAH) and carriers of clinically relevant mutations in *CYP21A2*.

Method and Patients Between July 2006 and July 2015 ACTH-tests were conducted in 365 children and adolescents (Age 1–20y) suspected to have NCCAH. As a reference, results from subsequent gene sequencing of *CYP21A2* was used. Inclusion criteria that were used were premature pubarche with accelerated bone age, hyperandrogenism, hirsutism, or menstrual irregularities. Receiver operating characteristics (ROC) were plotted. Evaluated test algorithms were composed around 17OHP measurements by radioimmunoassays. The most suitable test was identified by the greatest area under the curve (AUC).

Results Among the 12 tested algorithms, the sum of 30 min and 60 min stimulated 17OHP values (sum17OHPstim) showed the highest AUC of 0.774 for identifying heterozygous and bi-allelic mutations. A cut-off of 10.1 µg/l was advisable. Bi-allelic mutations only were best identified calculating the difference between 30 min and basal 17OHP values (Δ17OHP30). A cut-off of 9.4 µg/l was most effective.

Conclusion Alternatively to the above mentioned cut-offs the difference of 60 min after stimulation to basal 17OHP (Δ17OHP60) can be used for the benefit of a combined test to identify both heterozygotes and bi-allelic patients. There are minimal decreases in sensitivity and specificity compared to an approach that applies two tests. However, it denotes a simpler approach in the clinical routine.

Introduction

Congenital Adrenal Hyperplasia (CAH) is one of the most common autosomal recessive disorders. In 95 % of patients it is caused by a defect in the 21-hydroxylase enzyme encoding gene (*CYP21A2*) [1]. However, other enzyme defects are also possible as a cause of CAH. As CAH is generally regarded as a recessive disorder, both parental alleles must be affected [2]. 21-hydroxylase is important in the pro-

duction of aldosterone and cortisol from progesterone and 17α-hydroxyprogesterone (17OHP). As a consequence in 21-hydroxylase deficiency, adrenal androgens and above all the steroid and intermediate of androgen synthesis 17OHP are produced in excess. Among other effects, elevated levels of androgens can lead to hyperandrogenism and virilisation in females.

It is generally differentiated between classical and non-classical congenital adrenal hyperplasia (NCCAH). The classical form is caused by more severe enzyme defects, which lead to genital ambiguity at birth, and is further divided into the salt-wasting and non-salt-wasting form. The salt-wasting form leads to a potentially life-threatening loss of electrolytes and fluids (a salt-wasting crisis) in the first weeks after birth. NCCAH, also known as late-onset CAH, often remains unrecognised at birth and only becomes more apparent later on in childhood or puberty. This happens due to accelerated bone age, premature pubarche or severe acne.

New-born screening exists in Germany and many other countries around the world to identify patients with classical CAH. Also a third of NCCAH patients that are identified in this way [3, 4]. Nevertheless, this leaves a significant proportion of NCCAH patients undiagnosed. To identify patients with NCCAH or just heterozygous carriers, different methods exist. These either involve determining urine steroid profiles (USPs), conducting an ACTH-test and determining basal and post-stimulation blood steroid levels or performing molecular genetic analysis. USPs are favourable due to their non-invasiveness, especially in younger patients, but the ACTH-test poses an advantage over USPs as it may be conducted within about an hour's time during a single visit to the medical practice. In order to determine the USP the collection of urine over the course of 24 h is necessary. Molecular genetic tests are the only way of reliably identifying all NCCAH and heterozygous carriers, but currently involve significant costs and thus an effective pre-selection process is desirable.

Different approaches to ACTH-tests and their evaluation have been mentioned in previous publications [5–10]. Some studies have reported on the use of ACTH stimulated 21-deoxycortisol to identify NCCAH patients and heterozygotes [5, 11–13]. The results are especially promising when employing tandem mass spectrometry after high performance liquid chromatography separation (LC-MS/MS) [14] rather than radioimmunoassays [5, 12]. LC-MS/MS is becoming increasingly available but publications are still limited and non-decisive as to which calculation to use for carrier identification [14, 15]. Thus 21-deoxycortisol values are not ubiquitously accessible in the clinical routine and require further research. 17OHP presents a more common steroid used to distinguish NCCAH and heterozygotes but a medley of different approaches have been discussed. These include the use of basal 17OHP levels [16], conducting ACTH-tests and using stimulated 17OHP levels after 30 min [7] and 60 min [8, 17]; the difference (delta) between basal and stimulated (60 min) 17OHP levels [9, 18, 19] and the sum of stimulated (60 min) and basal 17OHP levels [10]. This study aims at identifying the most suitable test as well as proposing cut-off values to identify heterozygous or homozygous patients using that specific test.

Method

In this retrospective study, anonymised clinical data collected between July 2006 and July 2015 in the Ruhr area of Germany was used. In total, 365 patients with symptoms indicating NCCAH were included in this study. NCCAH symptoms for inclusion were premature pubarche, accelerated bone age, hyperandrogenism as well as signs of androgenisation such as hirsutism and menstrual irregu-

larities. The patients were above the age of one and up to and including the age of 20. All patients underwent an ACTH-Test and molecular genetic analysis of *CYP21A2*. When patients underwent two or more ACTH-Tests during the course of this study, only the most recent results were included. The ACTH-Test was conducted using 250 µg Synacthen, Novartis Pharma. Immediately before, 30 min after and 60 min after ACTH was administered, blood samples were collected and analysed. Tests were carried out in the morning and patients non-fasting. As an inclusion criterion for girls was irregular menstrual cycle, tests could not be performed during a specific known phase of the cycle. However, in patients with high levels of luteinizing hormone (LH), the ACTH test was repeated two weeks later to exclude measurements during the luteal phase. None of the patients was taking oral contraceptives.

The used data for this study included basal ACTH levels, basal 17OHP levels as well as 17OHP levels 30 and 60 min after ACTH was administered. Patients were only included when values of all three 17OHP measurements were known and the information was not pooled. Thus, associations between individual values remained to be known.

The result of the molecular genetic testing was regarded as gold standard to diagnose the presence or absence of *CYP21A2* mutation status and furthermore used as such when calculating sensitivity and specificity.

The following twelve tests will be compared:

- Basal 17OHP levels (17OHPBasal)
- Stimulated 17OHP levels 30 min after ACTH was given (17OHP30)
- Stimulated 17OHP levels 60 min after ACTH was given (17OHP60)
- Difference between 30 min stimulated and basal 17OHP levels (Δ 17OHP30)
- Difference between 60 min stimulated and basal 17OHP levels (Δ 17OHP60)
- Difference between 60 min and 30 min stimulated 17OHP levels (Δ 17OHPstim)
- Sum of basal and 30 min stimulated 17OHP levels (sum17OHP30)
- Sum of basal and 60 min stimulated 17OHP levels (sum17OHP60)
- Sum of 30 min and 60 min stimulated 17OHP levels (sum17OHPstim)
- Quotient: 30 min stimulated 17OHP divided by basal 17OHP (ratio17OHP30)
- Quotient: 60 min stimulated 17OHP divided by basal 17OHP (ratio17OHP60)
- Quotient: 60 min stimulated 17OHP divided by 30 min stimulated 17OHP levels (ratio17OHPstim)

Ratios of 17OHP values have thus far only been proposed in the form of a nomogram [20]. This paper investigates a wide range of tests to find the most appropriate. Not all of them have been mentioned in previous publications or used in practice.

17-OHP Measurements

Serum samples were analysed using radioimmunoassay (RIA) 17-alpha-Hydroxyprogesteron RIA-CT (DIA-Source). The intra- and in-

terassay coefficients of variation are 4.6 % (at 1.94 ng/ml) and 7.7 % (at 2.08 ng/ml) respectively [21]. During 2006 and 2015, the assay manufacturer and thus test characteristics did not change.

Molecular genetic analysis

The complete gene sequence of *CYP21A2* from c.-400 to c. * 179, including coding exons 1–10, all introns and relevant functional promoter elements starting with -400 and 3'UTR to c. * 179, were amplified by 3 allele specific PCR amplicons (E1–3, E2–6, E3–10) employing sequence differences (8bp deletion, cluster mutation in E6) to the highly similar pseudogene *CYP21A1*, and screened for mutations by Sanger sequencing. A search for deletions/duplications of *CYP21A2* and surrounding loci were added by MLPA methodology (Multiplex-Ligation-Dependent Probe Amplification; MRC-Holland, KIT P050, Lots B3 or C1). In patients presenting two different mutations, parents and siblings were offered genetic testing to exclude an allele carries mutations in *cis*. This mutation segregation was performed in 6 of the 8 patients with two differing mutations. All detected mutations were classified as mild or severe based on the database of the *Pharmacogene Variation Consortium* (PharmVar) [22]. A mutation was regarded as severe when causing salt-wasting or simple virilising CAH in homozygous state. Consequently, patients with bi-allelic mutations were divided into two groups (mild/mild; mild/severe) to assess, if harbouring a severe mutation correlated with more distinctive ACTH-Test results.

Statistics

Statistical analysis was conducted using SPSS v25. Differences in characteristics of patients with and without *CYP21A2* mutations were evaluated using the *t*-test for continuous variables and the χ^2 test for categorical variables. P-Values < 0.05 were considered significant. For the comparison of different tests accuracies for identifying patients with mutations in the *CYP21A2* gene, Receiver Operating Characteristics (ROC) were plotted. The calculation of an optimal cut-off point was performed using Pythagoras' theorem as described by Froud et al. [23].

Results

The studied patients were predominantly female, with females making up 88.8 % of the cohort. No difference in mean age and gender distribution between the group with and without a *CYP21A2* mutation could be observed (► **Table 1**). In contrast, mean 17OHP values before, 30 min and 60 min after the application of ACTH were significantly higher in the group carrying at least one mutated allele (► **Table 1**).

Molecular genetic analysis revealed one or more mutations in 83 (22.7 %) of the 365 patients. 8 patients had two or more different mild mutations and segregation analysis confirmed that mutations were located in *trans* on the *CYP21A2* alleles of 5 patients and in *cis* in 1 patient. The 2 patients for where mutation segregation was not performed were assigned to the heterozygous group. 4 patients harboured mild homozygous mutations, thus 9 patients with bi-allelic mutations were found. These can be regarded as NCCAH patients. 4 patients were compound heterozygous for a classical (severe) and a non-classical (mild) mutation and 1 patient carried a combination of two mild mutations (mild/mild).

Of the 74 carriers of mono-allelic mutations 66 had single heterozygous point mutations, 4 had duplications, 3 had gene conversions and 1 person had a gene deletion as illustrated by ► **Fig. 1**. By far the most often detected mutation was p.Val281Leu in 33 patients, 3 of them homozygous. The second and third most frequently detected mutations were p.Gln318Stop and p.Pro453Ser with 14 and 8 patients respectively. A complete count of detected mutations and their attributed severity is shown in ► **Table 2**.

Of the diagnostic tests investigated to identify patients with mutations affecting either one or both parental alleles of the *CYP21A2* gene, the sum of 30 min and 60 min stimulated values (sum 17OHPstim) showed the highest AUC of 0.774 (► **Table 3** and ► **Fig. 2**). The test was closely followed by 17OHP30 values and the difference between 60 min stimulated and basal values (Δ 17OHP60) with AUCs of 0.772 and 0.771 respectively. Basal 17OHP and all tests composed around calculating quotients were poor tests in this investigation, with AUC values below 0.6.

The optimal sum 17OHPstim cut-off found using Pythagoras' theorem was identified to be > 10.1 μ g/l. The > 10.1 μ g/l cut-off had a sensitivity 71.1 % and a specificity 71.6 %. In the studied cohort 138 of the patients had a Sum 17OHPstim > 10.1 μ g/l. 24 patients with mutations were not identified, which gave rise to a false negative rate of 0.289. The cut-off shows a positive likelihood ratio (LR+) of 2.50 and a negative likelihood ratio (LR-) of 0.40. Thus, a patient above this cut-off is 2.5 times more likely to be positive for at least one *CYP21A2* mutation than to be negative for the same, when genotyped.

Previously it has been suggested that one should use the Δ 17OHP60 test with a cut-off of 2.5 μ g/l for this purpose [19]. In our investigation this cut-off showed a sensitivity and specificity of 92.8 and 25.2 % respectively, therefore the LR+ was just 1.24 and the LR- 0.29.

To identify patients with bi-allelic mutations a number of tests that were conducted showed very high AUCs of 0.95 and above. In decreasing order these were Δ 17OHP30, Δ 17OHP60 and 17OHP30. The AUC of the most effective test, the difference between 30 min

► **Table 1** Patient characteristics by mutation status.

	No <i>CYP21A2</i> Mutation	All <i>CYP21A2</i> Mutation	Only heterozygous <i>CYP21A2</i> Mutations	Only bi-allelic <i>CYP21A2</i> Mutation
n	282	83	74	9
Females (n, %)	253 (89.7)	71 (85.5)	62 (83.8)	9 (100)
Age in years (mean $\pm$ SD)	10.99 $\pm$ 4.14	10.08 $\pm$ 3.87	10.13 $\pm$ 3.79	9.69 $\pm$ 4.71
17OHPBasal in μ g/l (mean $\pm$ SD)	1.45 $\pm$ 0.99	3.20 $\pm$ 7.73	1.77 $\pm$ 1.49	14.89 $\pm$ 20.51
17OHP30 in μ g/l (mean $\pm$ SD)	4.05 $\pm$ 1.47	10.17 $\pm$ 15.71	5.90 $\pm$ 2.32	45.29 $\pm$ 30.39
17OHP60 in μ g/l (mean $\pm$ SD)	4.76 $\pm$ 1.79	12.18 $\pm$ 19.67	6.82 $\pm$ 2.65	56.25 $\pm$ 38.11

stimulated and basal 17OHP ($\Delta 17\text{OHP}_{30}$), was 0.971. Again, calculating quotients of different 17OHP values showed to be a very weak test of which none refuted the null hypothesis (► **Table 3**).

As one might expect, the AUCs for the respective tests are even higher when used for only identifying patients with compound heterozygous mutations that show a combination of a mild and a severe mutation (► **Table 1S**, Supplementary Material).

A 60 min after stimulation 17OHP (17OHP₆₀) value of above 10 µg/l is described as diagnostic for a homozygous condition [24]. However, it is of note that this cut-off was established before the introduction of molecular genetic analysis of the *CYP21* gene. Previous publications have noted an overlap with heterozygous carriers for 17OHP₆₀ > 10 µg/l. This can also be observed in our cohort of the genotyped patients. 18 patients fulfilled the criterion 17OHP₆₀ > 10 µg/l. Of these, 8 (44%) showed biallelic mutations, 7 (39%) were heterozygous and in 3 (17%) genotyping revealed no

mutation. The three patients which showed no mutations had 17OHP₆₀ values of 11.6, 10.6 and 10.5 µg/l but no further genotyping for other genes was performed. Previously it has been reported, that the 17OHP₆₀ > 10 µg/l cut-off yielded no false negatives [25]. In our cohort one homozygous patient had 17OHP values 60 min after stimulation below 10 µg/l.

As an ideal test we recommend calculating the difference between 30 min stimulated and basal 17OHP values ($\Delta 17\text{OHP}_{30}$) and to use a cut-off of 9.4 µg/l. This criterion applied to 9 patients. 8 had confirmed bi-allelic mutations and 1 carried a p.Arg339His and p.Pro453Ser mutation. Due to lack of segregation analysis they could not be confirmed to be located in *trans*. Sensitivity is equal for the two cut-offs (88.9%) whereas specificity is slightly higher for $\Delta 17\text{OHP}_{30}$ with 99.7% compared to 99.2% for the 17OHP₆₀ cut-off. The LR+ is 296.3 for the $\Delta 17\text{OHP}_{30}$ cut-off compared to 111.1 for the 17OHP₆₀ cut-off. The LR- is equal for the two with 0.111.

► **Table 2** Detected Mutations and respective frequencies.

Mutation	Count	Classification (Mild/Severe)
p.Val281Leu (CYP21A2 * 15)	36	Mild
p.Gln318Stop (CYP21A2 * 17)	17	Severe
p.Pro453Ser (CYP21A2 * 19)	8	Mild
IVS2-13 A/C>G (CYP21A2 * 9)	5	Severe
p.Pro482Ser (CYP21A2 * 61)	3	Mild
p.Ile172Asn (CYP21A2 * 11)	3	Severe
-103A>G (CYP21A2 * 20D without c.-126C>T, c.-113G>A, c.-110T>C,)	3	Mild
p.Arg479Leu (CYP21A2 * 100)	2	Mild
8bp-Deletion (CYP21A2 * 10)	2	Severe
Gene deletion	2	Severe
-126 C>T (CYP21A2 * 20D without , c.-113G>A, c.-110T>C, c.-103A>G)	2	Mild
c.-126C>T, c.-113G>A, c.-110T>C, c.-103A>G (CYP21A2 * 20D)	1	Mild
-110T>C; -113G>A; -126C>T (CYP21A2 * 20D without -103A>G)	1	Mild
p.Ile172Asn, p.Ile236Asn, p.Val237Glu, p.Met239Lys (CYP21A2 * 20F without V281Leu))	1	Severe
p.Pro30Leu (CYP21A2 * 15)	1	Mild
p.Arg356Trp (CYP21A2 * 18)	1	Severe
* IVS2-13AG>A	1	Severe ¹
p.Ala434Val (CYP21A2 * 95)	1	Severe
8bp-Deletion, pGln318Stop	1	Severe
p.Arg339His, p.Pro453Ser (CYP21A2 * 24)	1	Mild

Count refers to number of alleles with respective mutation, not patients. Hence bi-allelic patients are counted twice. Unless stated otherwise, classification into mild and severe mutations is based on Data from the Pharmacogen Variation (PharmVar) Consortium.² 1. Neocleous V, Fanis P, Phylactou LA & Skordis N. Genotype Is Associated to the Degree of Virilization in Patients With Classic Congenital Adrenal Hyperplasia. *Front Endocrinol (Lausanne)* 2018 9 733. 2. Gaedigk A, Ingelman-Sundberg M, Miller NA, Leeder JS, Whirl-Carrillo M, Klein TE & PharmVar Steering C. The Pharmacogene Variation (PharmVar) Consortium: Incorporation of the Human Cytochrome P450 (CYP) Allele Nomenclature Database. *Clin Pharmacol Ther* 2018 103 399–401.

In both tests one patient with mutations affecting both parental alleles remained unrecognised. This patient showed a $\Delta 17\text{OHP}30$ of $3.6 \mu\text{g/l}$ and a homozygous $-126\text{C}>\text{T}$ promoter mutation in the *CYP21A2* gene. The $-126\text{C}>\text{T}$ mutation is commonly associated with $-103\text{A}>\text{G}$, $-110\text{T}>\text{C}$ and $-113\text{G}>\text{A}$ mutations, which were not found in this patient. However, in vitro experiments showed that the $-126\text{C}>\text{T}$ mutation itself can reduce *CYP21A2* transcriptional activity to 52 % [26], which suggests a pathogenic effect correlating to NCCAH.

For the identification of patients with at least one mutated allele or only bi-allelic mutations we therefore recommend the calculation of two different values id est for $\text{sum}17\text{OHPstim}$ and $\Delta 17\text{OHP}30$. For simplicity a physician might favour an approach with a single calculation or test with two separate cut-off values instead. In this case we propose the calculation of $\Delta 17\text{OHP}60$ values.

► **Fig. 2** ROC with the three best performing test for identification of patients with at least one mutated *CYP21A2* allele.

For identifying patients with at least one mutated parental allele the best cut-off point is $3.6 \mu\text{g/l}$. Compared to the $\text{sum}17\text{OHPstim}$ cut-off of $10.1 \mu\text{g/l}$ this cut-off has a higher sensitivity (75.9 %) and a lower specificity (66.0 %). Due to the higher sensitivity the number of false negatives is smaller with just 19 compared to 24. The false negative rate is 0.229 for the $\Delta 17\text{OHP}60$ cut-off. This obviously comes with the trade-off of more FPs. These were 97 and 79 for $\Delta 17\text{OHP}60$ and $\text{sum}17\text{OHPstim}$ respectively.

When using the $\Delta 17\text{OHP}60$ test to identify patients with bi-allelic mutations we recommend a cut-off of $10 \mu\text{g/l}$. It resulted in values for sensitivity and specificity of 88.9 % and 99.2 % respectively. Sensitivity is equal to that of the appropriate $\Delta 17\text{OHP}30$ cut-off. Specificity was 99.7 % and therefore a bit higher in the $\Delta 17\text{OHP}30$ test.

Discussion

In this study we assessed 12 different tests to identify individuals with heterozygous and bi-allelic *CYP21A2* mutations, based on ACTH-tests and molecular genetic analysis of 365 patients with symptoms of hyperandrogenism. For identifying carriers of at least one *CYP21A2* mutation we advocate calculating the sum of 30 min and 60 min post-ACTH stimulated values ($\text{sum}17\text{OHPstim}$) and to apply a cut-off of $> 10.1 \mu\text{g/l}$. The difference between 17OHP 30 min after stimulation and basal values ($\Delta 17\text{OHP}30$) above $9.4 \mu\text{g/l}$ proved to be the most effective for identifying patients with bi-allelic mutations. As a single test with two different cut-offs for detecting either carriers or NCCAH patients can be desirable in practice due to the simpler workflow we propose calculating the difference between basal and 17OHP values 60 min after ACTH was given ($\Delta 17\text{OHP}60$). As a cut-off to identify carriers of heterozygous mutations we propose elevating the pre-existing $\Delta 17\text{OHP}60$ cut-off from $2.5 \mu\text{g/l}$ [19] to $3.6 \mu\text{g/l}$. We ob-

► **Table 3** Information from Receiver operating characteristic for 12 investigative tests for identifying patients with at least one or two mutated alleles.

	Test Variable	Area under the curve (AUC)	Asymptotic Significance	Asymptotic 95 % Confidence Interval	
				Lower Bound	Upper Bound
At least one mutated parental allele	17OHPBasal	0.574	0.040	0.498	0.650
	17OHP30	0.772	0.000	0.707	0.836
	17OHP60	0.768	0.000	0.704	0.831
	Δ17OHP30	0.760	0.000	0.698	0.823
	Δ17OHP60	0.771	0.000	0.712	0.831
	Δ17OHPstim	0.604	0.004	0.529	0.679
	Sum17OHP30	0.739	0.000	0.673	0.806
	Sum17OHP60	0.737	0.000	0.671	0.803
	Sum 17OHPstim	0.774	0.000	0.710	0.838
	Ratio 17OHP30	0.600	0.006	0.527	0.673
	Ratio 17OHP60	0.604	0.004	0.531	0.677
	Ratio 17OHPstim	0.495	0.885	0.425	0.565
Only one mutated parental allele (heterozygous)	17OHPBasal	0.529	0.039	0.435	0.452
	17OHP30	0.728	0.035	0.000	0.658
	17OHP60	0.726	0.035	0.000	0.658
	Δ 17OHP30	0.713	0.034	0.000	0.646
	Δ17OHP60	0.727	0.033	0.000	0.663
	Δ17OHPstim	0.567	0.039	0.076	0.491
	Sum 17OHP30	0.694	0.036	0.000	0.623
	Sum 17OHP60	0.694	0.036	0.000	0.624
	Sum 17OHPstim	0.731	0.035	0.000	0.662
	Ratio 17OHP30	0.592	0.039	0.015	0.516
	Ratio 17OHP60	0.595	0.038	0.012	0.519
	Ratio 17OHPstim	0.489	0.036	0.778	0.418
Both parental alleles mutated (bi-allelic)	17OHPBasal	0.843	0.000	0.669	1.000
	17OHP30	0.955	0.000	0.871	1.000
	17OHP60	0.939	0.000	0.827	1.000
	Δ 17OHP30	0.971	0.000	0.919	1.000
	Δ17OHP60	0.956	0.000	0.877	1.000
	Δ17OHPstim	0.808	0.002	0.588	1.000
	Sum 17OHP30	0.943	0.000	0.837	1.000
	Sum 17OHP60	0.929	0.000	0.798	1.000
	Sum 17OHPstim	0.947	0.000	0.850	1.000
	Ratio 17OHP30	0.613	0.244	0.397	0.83
	Ratio 17OHP60	0.621	0.214	0.394	0.848
	Ratio 17OHPstim	0.533	0.735	0.311	0.755
The above AUC-calculations are based on the data of all 365 genotyped Patients. n (positive for at least one mutated parental allele)= 83; n (positive for both parental alleles mutated)= 9.					

served the Δ17OHP60 cut-off for bi-allelic patients to be optimal at 10 µg/l.

One of the major strengths of this study is the relatively large number of ACTH-tested and genotyped patients of a paediatric cohort (1–20 years) with clinical signs of hyperandrogenism. However, several limitations must be acknowledged. 17OHP measurements were performed using RIA, which have been shown to deliver measurements that tend to be substantially higher compared to those retrieved by LC-MS/MS [27], the gold standard for 17OHP measurements [1]. Nevertheless, the results can be of guidance in

the clinical routine, as LC-MS/MS has a higher cost and longer turn-around times, and thus the use of RIA still reflects the diagnostic routine in many clinical settings. In addition, the current Endocrine Society guidelines do not offer recommendations on thresholds for 17OHP values measured through RIA in the context of NCCAH diagnostics and carrier identification [1].

Considering an intra-assay variation of 4.6 % for the RIA used to determine 17OHP levels, one should consider to what extent the variation between the compared tests can be attributed to variance in laboratory results. 17OHP measurements may also have

been influenced by patients' meals prior to ACTH-testing as patients were non-fasting and meals are known to influence cortisol production [28–30]. The non-fasting interventions reflect data retrieved in the clinical routine and the effect is most likely minor as reported by Klose et al. [31] but this influence should be acknowledged.

Taken together the intra- and interassay variations of RIAs, as well as possible effects due to food intake and possible menstrual cycle confounders may reduce the practical difference between the investigated tests. The AUCs of the three best performing tests for identifying those patients with at least one mutated parental allele (Sum17OHPstim, 17OHP30, Δ 17OHP60) are very similar (► **Fig. 2**) with 0.774, 0.772 and 0.771, respectively. In the clinical routine they may indeed all be equivalent in their diagnostic capacity.

The AUCs may also be slanted by cases with p.Gln318Stop mutations attributed to the group of heterozygous patients, as localization of the mutation on an allele carrying two *CYP21A2* copies reduces the pathogenicity of this mutation and older cases have not been tested by MLPA to determine the co-occurrence of a duplicated haplotype.

The patients in this study were selectively analysed for mutations in *CYP21A2*, thus other mutations causing CAH such as in *CYP11B1* or *CYP17A1* are not accounted for. Mutations in these genes generally do not cause NCCAH [32] but it should be kept in mind that when aiming to identify carriers, the thresholds only apply for screening of 21-hydroxylase deficiency.

The CAH causing mutations besides *CYP21A2* are also a possible reason for the occurrence of elevated ACTH-test responses and negative molecular genetic analysis in this study. However, it is unlikely that this is the only explanation for negative genetic tests and high Δ 17OHP values. 132 Patients had Δ 17OHP60 > 4 µg/l and of these 79 (60 %) had negative genetic tests. Cinar et al. reported that PCOS patients had significantly higher ACTH stimulated 17OHP levels after 60 min than controls [33]. Though this is not supported by an investigation by Luboshitzky et al. [34], it demonstrates the necessity to identify possible reasons for increased 17OHP response during ACTH-testing besides CAH.

Though generally elevated, the significant overlap of responses to ACTH stimulation between individuals with and without *CYP21A2* mutations demonstrates the superiority of molecular genetic analysis in identifying heterozygotes. Similarly, also for bi-allelic patients genotyping remains necessary for making the final diagnosis of NCCAH. In the future, with continuously declining costs of molecular genetic analysis, it may become more resourceful to perform molecular genetic analysis initially, should symptoms apply. Considering especially that ACTH-tests may need to be repeated, additional costs can arise. To evaluate the adrenal reserve an ACTH-test could nonetheless be of relevance after genetic diagnosis in confirmed bi-allelic patients.

In individual cases, sequencing without prior ACTH-testing can already be adequate today when definitive molecular genetic results are needed, especially in the context of family planning and genetic counselling. For example, when the wish to have children exists and the patient presents with hyperandrogenemia, there is a notable family anamnesis or to determine carrier state of a patients' spouses. This is especially important as half of NCCAH males are asymptomatic and can otherwise remain unrecognized [25]. On the basis of genetic counselling prenatal therapy or, in the fu-

ture, genotyping using cell-free fetal DNA may be offered when the pregnancy is established and the neonate is at risk of CAH [35]. Therefore, it remains essential to critically reevaluate diagnostic schemas with evolving research and changing costs and access to diagnostic methods such as LC-MS/MS and sequencing technologies to ensure optimal patient care and clinical decisionmaking.

Ethics

In 2006 we made a request at the Ruhr-Universität Bochum, if an approval of an ethics committee will be needed for the envisaged clinical trial. Since we conducted an evaluation of inevitable clinical routine tests, we received a negative answer in this regard. After the introduction of the German Act of Gene Diagnostics ("Gendiagnostikgesetz") in Germany in 2009, we have not requested an ethics vote. Consent has been obtained from all parents after full explanation of the purpose and nature of all procedures used.

Funding

A. Richter-Unruh received honoraria and funding from Ferring, Hexal/Sandoz/Novartis, NovoNordisk, Ipsen, Kyowa Kirin and Merck/Serono.

Specific grant numbers were not assigned.

Author Statement

J. Meinel designed and performed the analysis and prepared the manuscript with assistance of A. Richter-Unruh. T. Haverkamp gathered the data, contributed the molecular genetic expertise and commented on the manuscript. F. Wünsche supported in the evaluation of the laboratory chemical data on the course and measurement procedures during the study and commented on the manuscript. A. Richter-Unruh conceptualized the idea and supervised the work's progression.

Declaration of Interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

References

- [1] Speiser PW, Arlt W, Auchus RJ et al. Congenital adrenal hyperplasia due to steroid 21-hydroxylase deficiency: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab* 2018; 103: 4043–4088. DOI: 10.1210/clinem.2018-01865
- [2] Merke DP. Approach to the adult with congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *J Clin Endocrinol Metab* 2008; 93: 653–660. DOI: 10.1210/jc.2007-2417
- [3] Gidlof S, Falhammar H, Thilen A et al. One hundred years of congenital adrenal hyperplasia in Sweden: a retrospective, population-based cohort study. *Lancet Diabetes Endocrinol* 2013; 1: 35–42. DOI: 10.1016/S2213-8587(13)70007-X

- [4] Gidlof S, Wedell A, Guthenberg C et al. Nationwide neonatal screening for congenital adrenal hyperplasia in Sweden: A 26-year longitudinal prospective population-based study. *JAMA Pediatr* 2014; 168: 567–574. DOI: 10.1001/jamapediatrics.2013.5321
- [5] Fiet J, Gueux B, Gourmelen M et al. Comparison of basal and adrenocorticotropin-stimulated plasma 21-deoxycortisol and 17-hydroxyprogesterone values as biological markers of late-onset adrenal hyperplasia. *J Clin Endocrinol Metab* 1988; 66: 659–667. DOI: 10.1210/jcem-66-4-659
- [6] Witchel SF, Lee PA. Identification of heterozygotic carriers of 21-hydroxylase deficiency: Sensitivity of ACTH stimulation tests. *Am J Med Genet* 1998; 76: 337–342
- [7] Kreutzmann DJ, Cowell CT, Howard NJ et al. Congenital adrenal hyperplasia family studies using the short ACTH test. *Australian Paediatric Journal* 1989; 25: 340–345
- [8] Bachega TA, Brenha EM, Billerbeck AE et al. Variable ACTH-stimulated 17-hydroxyprogesterone values in 21-hydroxylase deficiency carriers are not related to the different CYP21 gene mutations. *J Clin Endocrinol Metab* 2002; 87: 786–790. DOI: 10.1210/jcem.87.2.8247
- [9] Knochenhauer ES, Cortet-Rudelli C, Cunnigham RD et al. Carriers of 21-hydroxylase deficiency are not at increased risk for hyperandrogenism. *J Clin Endocrinol Metab* 1997; 82: 479–485. DOI: 10.1210/jcem.82.2.3759
- [10] Dacou-Voutetakis C, Dracopoulou M. High incidence of molecular defects of the CYP21 gene in patients with premature adrenarche. *J Clin Endocrinol Metab* 1999; 84: 1570–1574. DOI: 10.1210/jcem.84.5.5683
- [11] Kulle AE, Riepe FG, Hedderich J et al. LC-MS/MS based determination of basal- and ACTH-stimulated plasma concentrations of 11 steroid hormones: implications for detecting heterozygote CYP21A2 mutation carriers. *Eur J Endocrinol* 2015; 173: 517–524. DOI: 10.1530/EJE-14-1084
- [12] Bidet M, Bellanne-Chantelot C, Galand-Portier MB et al. Clinical and molecular characterization of a cohort of 161 unrelated women with nonclassical congenital adrenal hyperplasia due to 21-hydroxylase deficiency and 330 family members. *J Clin Endocrinol Metab* 2009; 94: 1570–1578. DOI: 10.1210/jc.2008-1582
- [13] Tardy V, Forest M, De Rougemont A et al. Detection of heterozygotes for 21-hydroxylase deficiency: validation of 21-deoxycortisol after ACTH test by sequencing of the entire CYP21 gene. *Horm Res Paediatr* 2005; 64: 41–42
- [14] Costa-Barbosa FA, Tonetto-Fernandes VF, Carvalho VM et al. Superior discriminating value of ACTH-stimulated serum 21-deoxycortisol in identifying heterozygote carriers for 21-hydroxylase deficiency. *Clin Endocrinol (Oxf)* 2010; 73: 700–706. DOI: 10.1111/j.1365-2265.2010.03871.x
- [15] Kulle AE, Riepe FG, Melchior D et al. A novel ultrahigh pressure liquid chromatography tandem mass spectrometry method for the simultaneous determination of androstenedione, testosterone, and dihydrotestosterone in pediatric blood samples: Age- and sex-specific reference data. *J Clin Endocrinol Metab* 2010; 95: 2399–2409
- [16] Escobar-Morreale HF, Sanchon R, San Millan JL. A prospective study of the prevalence of nonclassical congenital adrenal hyperplasia among women presenting with hyperandrogenic symptoms and signs. *J Clin Endocrinol Metab* 2008; 93: 527–533. DOI: 10.1210/jc.2007-2053
- [17] Nimkarn S, Gangishetti PK, Yau M et al. 21-Hydroxylase-Deficient Congenital Adrenal Hyperplasia. In: Pagon RA, Adam MP, Ardinger HH, Eds. *GeneReviews*(R). Seattle (WA): University of Washington, Seattle University of Washington, Seattle. All rights reserved. 1993
- [18] Schenk D. Prä-mature Pubarche - ACTH-Test und Mutationsanalyse des 21-Hydroxylasegens; Dissertation, LMU München. 2005
- [19] Weil J, Bidlingmaier F, Sippell WG et al. Comparison of two tests for heterozygosity in congenital adrenal hyperplasia (CAH). *Acta endocrinologica* 1979; 91: 109–121
- [20] New MI, Lorenzen F, Lerner AJ et al. Genotyping steroid 21-hydroxylase deficiency: hormonal reference data. *J Clin Endocrinol Metab* 1983; 57: 320–326. DOI: 10.1210/jcem-57-2-320
- [21] [Anonymous] In: *DIAsource ImmunoAssays S.A.*
- [22] Gaedigk A, Ingelman-Sundberg M, Miller NA et al. The Pharmacogene Variation (PharmVar) Consortium: Incorporation of the Human Cytochrome P450 (CYP) Allele Nomenclature Database. *Clin Pharmacol Ther* 2018; 103: 399–401. DOI: 10.1002/cpt.910
- [23] Froud R, Abel G. Using ROC curves to choose minimally important change thresholds when sensitivity and specificity are valued equally: the forgotten lesson of pythagoras. theoretical considerations and an example application of change in health status. *PLoS One* 2014; 9: e114468. DOI: 10.1371/journal.pone.0114468
- [24] Moran C, Azziz R. 21-hydroxylase-deficient nonclassical adrenal hyperplasia: the great pretender. *Seminars in reproductive medicine* 2003; 21: 295–300. DOI: 10.1055/s-2003-43307
- [25] Livadas S, Dracopoulou M, Dastamani A et al. The spectrum of clinical, hormonal and molecular findings in 280 individuals with nonclassical congenital adrenal hyperplasia caused by mutations of the CYP21A2 gene. *Clin Endocrinol (Oxf)* 2015; 82: 543–549. DOI: 10.1111/cen.12543
- [26] Araujo RS, Mendonca BB, Barbosa AS et al. Microconversion between CYP21A2 and CYP21A1P promoter regions causes the nonclassical form of 21-hydroxylase deficiency. *J Clin Endocrinol Metab* 2007; 92: 4028–4034. DOI: 10.1210/jc.2006-2163
- [27] Chesover AD, Millar H, Sepiashvili L et al. Screening for nonclassical congenital adrenal hyperplasia in the era of liquid chromatography-tandem mass spectrometry. *J Endocr Soc* 2020; 4: bvz030. DOI: 10.1210/jendso/bvz030
- [28] Brandenberger G, Follenius M, Hietter B. Feedback from meal-related peaks determines diurnal changes in cortisol response to exercise. *J Clin Endocrinol Metab* 1982; 54: 592–596. DOI: 10.1210/jcem-54-3-592
- [29] Pasquali R, Biscotti D, Spinucci G et al. Pulsatile secretion of ACTH and cortisol in premenopausal women: effect of obesity and body fat distribution. *Clin Endocrinol (Oxf)* 1998; 48: 603–612. DOI: 10.1046/j.1365-2265.1998.00458.x
- [30] Quigley ME, Yen SS. A mid-day surge in cortisol levels. *J Clin Endocrinol Metab* 1979; 49: 945–947. DOI: 10.1210/jcem-49-6-945
- [31] Klose M, Lange M, Rasmussen AK et al. Factors influencing the adrenocorticotropin test: role of contemporary cortisol assays, body composition, and oral contraceptive agents. *J Clin Endocrinol Metab* 2007; 92: 1326–1333. DOI: 10.1210/jc.2006-1791
- [32] Krone N, Arlt W. Genetics of congenital adrenal hyperplasia. *Best Pract Res Clin Endocrinol Metab* 2009; 23: 181–192. DOI: 10.1016/j.beem.2008.10.014
- [33] Cinar N, Harmanci A, Aksoy DY et al. Adrenocortical steroid response to ACTH in different phenotypes of non-obese polycystic ovary syndrome. *J Ovarian Res* 2012; 5: 42. DOI: 10.1186/1757-2215-5-42
- [34] Luboshitzky R, Ishai A, Shen-Or Z et al. Evaluation of the pituitary-adrenal axis in hyperandrogenic women with polycystic ovary syndrome. *Neuro endocrinology letters* 2003; 24: 249–254
- [35] Dorr HG, Binder G, Reisch N et al. Experts' Opinion on the Prenatal Therapy of Congenital Adrenal Hyperplasia (CAH) Due to 21-Hydroxylase Deficiency - Guideline of DGKED in cooperation with DGGG (S1-Level, AWMF Registry No. 174/013, July 2015). *Geburtshilfe Frauenheilkd* 2015; 75: 1232–1238. DOI: 10.1055/s-0041-109717