A case of severe hepatolithiasis after hepaticojejunostomy with Roux-en-Y reconstruction treated by endoscopic ultrasound-guided transhepatic antegrade stone removal

Balloon enteroscopy-assisted endoscopic retrograde cholangiopancreatography (ERCP) is useful for the treatment of bile duct stones in patients with surgically altered anatomy [1–3]; however, the procedure cannot always successfully remove stones when the anastomotic site or papilla of Vater cannot be reached. Recently, endoscopic ultrasound (EUS)-guided antegrade intervention has been developed for benign biliary diseases, including hepatolithiasis in patients with altered anatomy [4, 5]. We report a case of EUS-guided transhepatic antegrade stone removal (EUS-TASR) in a patient with altered anatomy (► **Video 1**).

The 59-year-old man underwent hepaticojejunostomy with Roux-en-Y reconstruction for hepatolithiasis several years previously. He was referred to our hospital for treatment of severe hepatolithiasis. Abdominal computed tomography showed huge impacted stones in the hilum and a dilated left intrahepatic bile duct (► **Fig. 1 a**).

We chose to perform EUS-TASR because accessing the anastomotic site by balloon-assisted ERCP was judged to be dif-

► **Video 1** Endoscopic ultrasound-guided transhepatic antegrade stone removal was useful for severe hepatolithiasis after hepaticojejunostomy with Roux-en-Y reconstruction.

ficult owing to the long afferent loop and because multiple sessions would be required for stone removal.

We first performed EUS-guided hepaticogastrostomy (EUS-HGS). The dilated left intrahepatic bile duct was punctured using a 19-gauge fine-needle aspiration

needle from the residual stomach side. Cholangiography showed multiple impacted stones from the hilum to the bilateral intrahepatic bile ducts. A 0.025-inch guidewire was inserted into the intrahepatic bile duct and the needle tract was dilated using an ultra-tapered mechani-

► **Fig. 1** Computed tomography. **a** Before treatment, showing huge impacted stones in the hilum and a dilated left intrahepatic bile duct. **b** After the procedure, showing complete clearance.

► **Fig. 2** Endoscopic ultrasound-guided hepaticogastrostomy.

► **Fig. 3** Bile duct stones were crushed using electrohydraulic lithotripsy under direct visualization.

► **Fig. 4** Stone fragments were removed using a basket catheter through the hepaticogastrostomy route under direct visualization.

cal dilator. A fully covered metal stent was then placed over the guidewire (► **Fig. 2**). EUS-TASR was performed 1 month later. After dilation of the anastomotic stricture using a balloon catheter, a cholangioscope was inserted into the intrahepatic bile duct through the metal stent. Bile duct stones were crushed using electrohydraulic lithotripsy under direct visualization (► **Fig. 3**). Subsequently, stone fragments were removed using a basket catheter and balloon catheter through the anastomotic route and HGS route (► **Fig. 4**). Finally, the huge hepatic bile duct stone was completely removed (► **Fig. 1 b**).

Endoscopy_UCTN_Code_TTT_1AS_2AD

Competing interests

The authors declare that they have no conflict of interest.

The authors

Toshihiro Homma¹, Kenjiro Yamamoto¹, Takayoshi Tsuchiya¹, Ryosuke Tonoizuka¹, Hirohito Minami¹, Eri Joyama², Takao Itoi¹

- 1 Department of Gastroenterology and Hepatology, Tokyo Medical University Hospital, Tokyo, Japan
- 2 Department of International medicine, Tokyo Medical University Hospital, Tokyo, Japan

Corresponding author

Kenjiro Yamamoto, MD

Department of Gastroenterology and Hepatology, Tokyo Medical University, 6-7-1, Nishishinjuku, Shinjuku-ku, Tokyo 160-0023, Japan
Fax: +81-3-53816654
ken.yamamoto5544@gmail.com

References

- [1] Itoi T, Ishii K, Sofuni A et al. Long- and short-type double-balloon enteroscopy-assisted therapeutic ERCP for intact papilla in patients with a Roux-en-Y anastomosis. *Surg Endosc* 2011; 25: 713–721
- [2] Itokawa F, Itoi T, Ishii K et al. Single- and double-balloon enteroscopy-assisted endoscopic retrograde cholangiopancreatography in patients with Roux-en-Y plus hepaticojejunostomy anastomosis and Whipple resection. *Dig Endosc* 2014; 26: 136–143
- [3] Ishii K, Itoi T, Tonoizuka R et al. Balloon enteroscopy-assisted ERCP in patients with Roux-en-Y gastrectomy and intact papillae (with videos). *Gastrointest Endosc* 2016; 83: 377–386
- [4] Itoi T, Sofuni A, Tsuchiya T et al. Endoscopic ultrasonography-guided transhepatic antegrade stone removal in patients with surgically altered anatomy: case series and technical review (with videos). *J Hepatobiliary Pancreat Sci* 2014; 21: E86–E93
- [5] Mukai S, Itoi T, Sofuni A et al. EUS-guided antegrade intervention for benign biliary diseases in patients with surgically altered anatomy (with videos). *Gastrointest Endosc* 2019; 89: 399–407

Bibliography

Endoscopy 2021; 53: E230–E231

DOI 10.1055/a-1244-9556

ISSN 0013-726X

published online 23.9.2020

© 2020. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

CORRECTION

A case of severe hepatolithiasis after hepaticojejunostomy with Roux-en-Y reconstruction treated by endoscopic ultrasound-guided transhepatic antegrade stone removal

Homma T, Yamamoto K, Tsuchiya T et al. A case of severe hepatolithiasis after hepaticojejunostomy with Roux-en-Y reconstruction treated by endoscopic ultrasound-guided transhepatic antegrade stone removal.

Endoscopy 2020, 52:
10.1055/a-1244-9556

In the above-mentioned article, the institution of Takao Itoi has been corrected. This was corrected in the online version on September 30, 2020.