Obstructive jaundice with a biliary clot post-endoscopic sphincterotomy treated with clipping and endoscopic biliary stenting

A 62-year-old man was admitted with epigastralgia. He had a history of laparoscopic cholecystectomy for cholecystolithiasis. He had no coagulopathy and was not taking anticoagulants. Abdominal computed tomography (CT) showed a common bile duct (CBD) stone (► **Fig. 1 a**). Endoscopic retrograde cholangiography (ERC) and intraductal ultrasonography (IDUS) also showed a 2.8-mm CBD stone (► **Fig. 1 b**, ► **Fig. 2 a, b**). Endoscopic sphincterotomy (EST) was performed (► **Fig. 2 c**) and the CBD stone was removed using a wire basket (► **Fig. 2 d**).

The patient complained of epigastralgia again after 4 days. Laboratory investigations demonstrated elevated cholestatic parameters: total bilirubin 2.8 mg/dL (normal range 0.4–1.5 mg/dL), aspartate aminotransferase 176 U/L (13–30 U/L), alanine aminotransferase 146 U/L (10–42 U/L), alkaline phosphatase 233 U/L (38–113 U/L), and gamma-glutamyl transpeptidase 695 U/L (9–32 U/L); hemoglobin was within the normal limit. CT showed a diffuse high-density structure in the CBD, with the bile duct mildly dilated (► **Fig. 3 a**). ERC revealed post-EST bleeding and a biliary clot in the CBD (► **Fig. 3 b**, ► **Fig. 4 a**). The clot was removed using a grasping forceps and wire basket (► **Fig. 3 c**, ► **Fig. 4 b**), and an endoscopic biliary stent (EBS) was inserted into the CBD for biliary drainage. Clipping was applied to stop the bleeding (► **Fig. 3 d**, ► **Fig. 4 c, d**, ► **Video 1**). The patient progressed well after the procedures. The EBS was removed 8 days postoperatively and the patient was discharged 10 days postoperatively.

► **Video 1** The biliary clot, caused by delayed bleeding after endoscopic sphincterotomy, was removed using a grasping forceps and wire basket, and clips were applied to stop the bleeding.

► **Fig. 1** Common bile duct stone (arrow). **a** Abdominal computed tomography. **b** Endoscopic retrograde cholangiography.

► **Fig. 2** Removal of the common bile duct (CBD) stone. Endoscopic views (**a, c, d**) and intraductal ultrasonography (IDUS) view (**b**). **a** The ampulla of Vater was intact (arrow). **b** IDUS showed a CBD stone, approximately 2.8 mm in size (arrow). **c** Immediately after endoscopic sphincterotomy (arrow). **d** The CBD stone was removed using a wire basket (arrow).

The incidence of post-EST delayed hemorrhage is 1.62% [1], and biliary obstruction with a biliary clot caused by post-EST bleeding is extremely rare [2–5]. Endoscopic hemostasis is currently the first treatment choice for post-ERC bleeding, with balloon dilation and biliary stent placement used for treatment [2–5]. To the best of our knowledge,

this is the first English case report of obstructive jaundice with a biliary clot caused by post-EST bleeding, treated with clipping and an EBS, which may be an effective endoscopic technique for treating such cases.

Endoscopy_UCTN_Code_CPL_1AK_2AC

Competing interests

The authors declare that they have no conflict of interest.

The authors

Takashi Abe¹, Takehiko Nariyasu¹, Takayuki Nagai¹, Marina Hamamoto¹, Masato Hanzawa¹, Yasuhisa Hiroshima¹, Kazunari Murakami²

- 1 Department of Gastroenterology, Oita Kouseiren Tsurumi Hospital, Beppu, Japan
- 2 Department of Gastroenterology, Faculty of Medicine, Oita University, Yufu, Japan

Corresponding author

Takashi Abe, MD, PhD

Department of Gastroenterology, Oita Kouseiren Tsurumi Hospital, Tsurumi 4333, Beppu City, Oita 874-8585, Japan
Fax: +81-977-237884
takashi0315@oita-u.ac.jp

► **Fig. 3** Imaging after endoscopic sphincterotomy (EST) and common bile duct (CBD) stone removal. **a** Reconstructed coronal image of abdominal computed tomography on day 4 after EST showed a diffuse high-density structure in the CBD with the bile duct being mildly dilated (arrows). **b** Endoscopic retrograde cholangiography revealed a diffuse filling defect in the CBD with the bile duct mildly dilated (arrows). **c** The CBD was cleaned up by removing the biliary clot using a grasping forceps and wire basket. **d** An endoscopic biliary stent was inserted into the CBD and clipping was applied for endoscopic hemostasis.

References

- [1] Yan J, Zhou CX, Wang C et al. Risk factors for delayed hemorrhage after endoscopic sphincterotomy. *Hepatobiliary Pancreat Dis Int* 2020. doi:10.1016/j.hbpd.2019.12.010
- [2] Mosenkis BN, Brandt LJ. Bleeding causing biliary obstruction after endoscopic sphincterotomy. *Am J Gastroenterol* 1997; 92: 708–709
- [3] Ala A, Khin CC, van Someren N. Common bile duct thrombus: a cause of persisting obstructive jaundice after endoscopic sphincterotomy. *Gastrointest Endosc* 1999; 50: 285–286
- [4] Ergül B, Koçak E, Köklü S. An unusual complication of ERCP: obstructive jaundice due to a blood clot. *Clin Res Hepatol Gastroenterol* 2012; 36: e40–e41
- [5] Zhu Y, Wang S, Zhao S et al. Obstructive jaundice due to a blood clot after ERCP: a case report and review of the literature. *BMC Gastroenterol* 2018; 18: 163

Bibliography

Endoscopy 2021; 53: E297–E300
 DOI 10.1055/a-1260-2903
 ISSN 0013-726X
 published online 8.10.2020
 © 2020. Thieme. All rights reserved.
 Georg Thieme Verlag KG, Rüdigerstraße 14,
 70469 Stuttgart, Germany

► **Fig. 4** Treatment of the biliary clot and delayed bleeding after endoscopic sphincterotomy (EST). Endoscopic views. **a** Post-EST delayed bleeding and the clot at the orifice of the common bile duct (CBD). **b** The biliary clot was removed using a grasping forceps. **c** The bleeding point was revealed (arrow). **d** An endoscopic biliary stent was inserted into the CBD and clipping was applied to stop the bleeding.

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>