Endoscopic ultrasound-guided recanalization of complete pharyngoesophageal stenosis

► **Fig. 1** Esophagogram showing complete obstruction of the esophagus and passage of contrast into the airway.

A 74-year-old man was referred because of complete esophageal obstruction. He had a diagnosis of T3N1 hypopharyngeal squamous cell carcinoma and had undergone chemoradiotherapy with a complete response. He had a percutaneous endoscopic gastrostomy (PEG) for nutrition and had had aphagia for 18 months. Assessment of the stricture using contrast swallow demonstrated complete esophageal obstruction (► **Fig. 1**), and combined anterograde (peroral) and retrograde endoscopy (through the feeding gastrostomy tract) revealed complete

esophageal obstruction at the level of the upper esophageal sphincter (► **Fig. 2**, ► **Fig. 3**). An attempt at rendezvous was unsuccessful [1].

We decided to perform the procedure guided by endoscopic ultrasound (EUS) (► **Video 1**). Through the PEG, a guidewire was advanced into the upper esophagus and an endoscopic retrograde cholangiopancreatography (ERCP) extractor balloon inserted over the guidewire. The balloon was filled with contrast to give a visible target on EUS. However, the balloon was not adequately identified with

► **Fig. 2** Peroral endoscopy. Complete obstruction at the upper esophageal sphincter.

► **Fig. 3** Retrograde endoscopy through the feeding gastrostomy tract. Complete obstruction in the upper esophagus.

► **Fig. 4** The guidewire passing into a suitable position in the esophagus.

Video 1 Endoscopic ultrasound-guided recanalization of complete esophageal obstruction.

Corresponding author

José R. Aparicio, MD, PhD

Unidad de Endoscopia, Servicio de Medicina Digestiva, ISABIAL Hospital General Universitario de Alicante, Avda. Pintor Baeza s/n, 03013 Alicante, Spain
Fax: +34-96-5913629
japariciot@gmail.com

Reference

- [1] Fusco S, Kratt T, Gani C et al. Rendezvous endoscopic recanalization for complete esophageal obstruction. *Surg Endosc* 2018; 32: 4256–4262

Bibliography

Endoscopy 2021; 53: E309–E310

DOI 10.1055/a-1270-6704

ISSN 0013-726X

published online 8.10.2020

© 2020. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>

the echoendoscope positioned in the hypopharynx.

On fluoroscopy, a separation between the balloon and the echoendoscope was observed, by which contrast was introduced through the lumen of the balloon. EUS now showed a good target in the esophageal lumen. With a 19-G needle, the esophageal lumen was punctured and a 0.025-inch Visiglide guidewire advanced. The echoendoscope was removed, and adequate positioning of the guidewire was verified with a gastro-scope (►Fig. 4). A 6-Fr cystotome and a 6-mm dilation balloon were used to create a passage, followed by placement of a 12-Fr jejunal probe to keep the passage patent.

After 7 days, regular dilation sessions were started with Savary bougienage to 16-mm in diameter. In two of the sessions, mitomycin was injected at the level of the stenosis. After 13 dilations, ade-

quate tolerance of a normal diet was achieved and the PEG was withdrawn. At 3 years' follow-up, no recurrence of stenosis was seen.

Endoscopy_UCTN_Code_TTT_1AO_2AH

Competing interests

Dr. J. R. Aparicio is a consultant for Boston Scientific.

The authors

Belén Martínez-Moreno, Lucia Medina-Prado, Sandra Baile-Maxía, Carolina Mangas-Sanjuan, Juan A. Casellas, José R. Aparicio

Unidad de Endoscopia, Servicio de Medicina Digestiva, ISABIAL Hospital General Universitario de Alicante, Spain