Salvage endoscopic submucosal dissection with a small-caliber endoscope for recurrent esophageal cancer on distal side of esophageal stricture

► **Fig. 1** Endoscopy showed circumferential ulceration and stricture in the cervical esophagus that had developed due to chemoradiotherapy.

► **Fig. 2** Endoscopy showed a half-circumferential flat cancer on the distal side of the esophageal stricture.

Salvage endoscopic submucosal dissection (ESD) is effective for local recurrent esophageal cancer after chemoradiotherapy [1, 2]. However, severe late toxicity, mostly in the form of esophageal strictures and ulcerations, can occur in patients who undergo chemoradiotherapy in the definitive treatment of esophageal cancer [3]. Endoscopic treatment for esophageal neoplasia in patients with esophageal stricture is challenging [4]. A 76-year-old woman with recurrent esophageal cancer after chemoradiotherapy was referred to our department for endoscopic treatment. Endoscopy showed a cervical esophageal stricture

► **Video 1** Endoscopic submucosal dissection with a small-caliber endoscope for recurrent esophageal cancer on the distal side of the esophageal stricture.

► **Fig. 3** Markings, mucosal incision, and submucosal dissection were all performed using an endo-knife.

► **Fig. 4** The lesion was completely resected en bloc in 55 minutes without any complications.

due to chemoradiotherapy (► **Fig. 1**) and a half-circumferential flat cancer on the distal side of the stricture (► **Fig. 2**; ► **Video 1**). Although balloon dilation was performed, the therapeutic endoscope could not be passed through the stricture. We performed ESD with a small-caliber endoscope (EG-L580NW7; Fujifilm, Tokyo, Japan; diameter 5.8 mm, working channel 2.4 mm). Small-caliber ESD devices such as an endo-knife, clip-line traction device, and hemostatic for-

ceps were used (Souten, Ichigan, and Raicho, respectively; Kaneka Medics, Tokyo, Japan). A distal attachment was used to stabilize the endoscope. Markings, mucosal incision, and submucosal dissection were all performed using the endo-knife (► **Fig. 3**). After circumferential mucosal incision, a traction device was applied to obtain a good field of view for dissection. A small-caliber hemostatic forceps was used to arrest bleeding. Finally, the lesion was comple-

► **Fig. 5** Histopathological examination revealed squamous cell carcinoma with slight submucosal invasion and negative horizontal and vertical margins. Upper: macroscopic view of the resected specimen. Lower: hematoxylin–eosin stain, $\times 100$ magnification.

tely resected en bloc (► **Fig. 4**) in 55 minutes without any complications. Histopathology revealed squamous cell carcinoma with slight submucosal invasion and negative horizontal and vertical margins (► **Fig. 5**). No other treatment for the recurrent esophageal cancer was performed because the patient refused additional surgical treatment.

In conclusion, salvage ESD with a small-caliber endoscope can be a treatment option for recurrent superficial esophageal cancer after chemoradiotherapy

where a conventional endoscope cannot be used because of proximal esophageal stricture.

Endoscopy_UCTN_Code_TTT_1AO_2AG

Competing interests

The authors declare that they have no conflict of interest.

The authors

Kyosuke Tanaka¹, Yuhei Umeda^{1,2}, Hiroki Asakawa², Hiroshi Miura¹, Misaki Nakamura², Masaki Katsurahara¹, Yasuhiko Hamada²

¹ Department of Endoscopy, Mie University Hospital, Tsu, Japan

² Department of Gastroenterology and Hepatology, Mie University Hospital, Tsu, Japan

Corresponding author

Kyosuke Tanaka, MD, PhD

Department of Endoscopy, Mie University Hospital, 2-174 Edobashi, Tsu, Mie 514-8507, Japan

Fax: +81-59-231-5200

kyosuket@clin.medic.mie-u.ac.jp

References

- [1] Hatogai K, Yano T, Kojima T et al. Local efficacy and survival outcome of salvage endoscopic therapy for local recurrent lesions after definitive chemoradiotherapy for esophageal cancer. *Radiat Oncol* 2016; 11: 31
- [2] Saito Y, Takisawa H, Suzuki H et al. Endoscopic submucosal dissection of recurrent or residual superficial esophageal cancer after chemoradiotherapy. *Gastrointest Endosc* 2008; 67: 355–359
- [3] Roeder F, Nicolay NH, Nguyen T et al. Intensity modulated radiotherapy (IMRT) with concurrent chemotherapy as definitive treatment of locally advanced esophageal cancer. *Radiat Oncol* 2014; 9: 191
- [4] Nishiyama N, Kobara H, Yachida T et al. Strategy of small-caliber endoscopic submucosal dissection for esophageal neoplasia distal to severe stricture. *Endoscopy* 2020; 52: E94–E95

Bibliography

Endoscopy 2021; 53: E363–E364

DOI 10.1055/a-1296-7632

ISSN 0013-726X

published online 19.11.2020

© 2020. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany