Management of fistula between neobladder and rectum using an atrial septal defect closure device

► **Fig. 1** Contrast-enhanced computed tomography of the abdomen showing a fistulous opening measuring 1.4 cm × 1.9 cm between neobladder and rectum.

Radiation therapy is widely used to treat malignancies in the pelvic cavity such as cancers of the colorectum, urinary tract, uterus, adnexa, and prostate. Studies have reported pelvic radiotherapy to have been closely associated with gastrointestinal complications such as mucosal ulcers, stenosis, perforation, and fistula formation [1].

These fistulous tracts are life-threatening since the patient is in a perpetual state of urosepsis. Surgical interventions are risky due to underlying advanced disease. Therefore, the management of an enterovesical fistula between a surgical neobladder and the rectum in patients previously operated on for bladder malignancy is challenging [2]. We present the case of a 58-year-old man who underwent surgery for bladder carcinoma with construction of a neobladder. The surgery was followed by chemoradiation therapy.

The patient presented in the emergency department with a 15-day history of fever, hypotension, pneumaturia, and fecaluria. Contrast-enhanced computed tomography of the abdomen revealed a fistulous opening measuring 1.4 cm × 1.9 cm between neobladder and rectum (► **Fig. 1**). Successful endoscopic closure

► **Video 1** Management of fistula between neobladder and rectum using an atrial septal defect closure device.

of the rectovesical fistula was achieved using an atrial septal defect closure device (► **Video 1**). Following the procedure, the urosepsis and hypotension improved and the patient was catheterized for 3 days. The patient had clear urine on the 3rd day and a culture after 72 hours was sterile. He was discharged on the 7th day. He remained well for one and a half months before succumbing to his underlying advanced malignancy.

Endoscopy_UCTN_Code_TTT_1AQ_2AJ

Competing interests

The authors declare that they have no conflict of interest.

The authors

Ashlee Sidhu, Nagoor Basha Shaik, Malay Sharma

Aryavart Hospital, Daurala Sugar Mills, Meerut, Uttar Pradesh, India

Corresponding author

Ashlee Sidhu, MD

Aryavart Hospital, NH334, Daurala Sugar Mills, Meerut, Uttar Pradesh – 250110, India
ashlee2907@gmail.com

References

- [1] Iwamuro M, Hasegawa K, Hanayama Y et al. Enterovaginal and colovesical fistulas as late complications of pelvic radiotherapy. *J Gen Fam Med* 2018; 19: 166–169
- [2] Golabek T, Szymanska A, Szopinski T et al. Enterovesical fistulae: aetiology, imaging, and management. *Gastroenterol Res Pract* 2013; 2013: 617967

Bibliography

Endoscopy 2021; 53: 1090
DOI 10.1055/a-1298-3906
ISSN 0013-726X
published online 26.11.2020
© 2021. Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany