Electrohydraulic lithotripsy for the treatment of stone impacted in a lumen-apposing metal stent in a patient with endoscopic cholecystoduodenostomy

► **Fig. 1** Lumen-apposing metal stent (LAMS) obstructed by a biliary stone impacted in the stent lumen.

► **Fig. 2** Fragmentation of biliary stone by means of electrohydraulic lithotripsy.

An 89-year-old woman presented to our department having experienced acute cholecystitis a few weeks previously. Her comorbidities, chronic kidney disease, and congestive heart failure meant she was not a suitable candidate for surgery. She underwent EUS-guided gallbladder drainage, which was performed without complications. During this procedure, a 10-mm×10-mm lumen-apposing metal stent (LAMS) (Hot Axios; Boston Scientific) was implanted for gallbladder drainage. A few days later, the patient complained of right-upper quadrant abdominal pain and fever, and a new acute cholecystitis episode was diagnosed.

► **Video 1** Biliary stone occluding a lumen-apposing metal stent treated with electrohydraulic lithotripsy in a patient who had previously undergone endoscopic ultrasound-guided cholecystoduodenostomy.

► **Fig. 3** A 10-mm fistula in the gallbladder wall with fragmented stones in the cavity.

► **Fig. 4** Placement of a plastic double-pigtail stent to prevent renewed occlusion.

Upper endoscopy with a therapeutic endoscope (GIF-1TH190; Olympus) was performed. The endoscope was introduced as far as the duodenum, where it became evident that the LAMS was obstructed by a biliary stone impacted in the stent lumen (► **Fig. 1**; ► **Video 1**). Electrohydraulic lithotripsy (EHL) (1.9-Fr,

375-cm Biliary EHL Probe Autolith; Boston Scientific) was performed to fragment the obstructive biliary stone. The EHL probe was introduced through the endoscope, and fragmentation with the infusion of saline was started (► **Fig. 2**). The gallbladder stone was broken into multiple fragments that were removed with a Dormia basket. During the procedure, which lasted about 60 min, purulent secretion was seen draining from the

gallbladder. The gallbladder wall showed a 10 mm-fistula with some fragmented stones within the cavity (►Fig. 3); these were removed with a Roth Net retriever. To prevent renewed LAMS occlusion, placement of a plastic double-pigtail stent was decided on (►Fig. 4). No complications were observed in the patient; she tolerated the procedure and was discharged home a few days later. EUS-guided gallbladder drainage in patients who are not candidates for surgery is a safe technique and has a low complication rate. In patients with large stones, recurrent cholecystitis, impaction, and LAMS obstruction can occur [1,2]. EHL and laser lithotripsy are two useful techniques that allow fragmentation of stones; EHL generates high-amplitude hydraulic pressure waves, while laser lithotripsy uses a laser beam with repetitive pulses of laser energy to create a mechanical shockwave [3,4]. EHL is a safe option to resolve LAMS occlusion in cholecystoduodenostomy.

Endoscopy_UCTN_Code_CPL_1AL_2AD

Competing interests

The authors declare that they have no conflict of interest.

The authors

Edson Guzmán-Calderón^{1,2,3}, **Belén Martínez-Moreno⁴**, **Juan A. Casellas⁴**, **José Ramón Aparicio⁴**

- 1 Gastroenterology Unit, Hospital Nacional Edgardo Rebagliati Martins, Lima, Peru
- 2 Universidad Peruana de Ciencias Aplicadas (UPC), Lima, Peru
- 3 Gastroenterology Unit, Angloamericana Clinic, Lima, Peru
- 4 Gastroenterology Unit, Hospital General Universitario de Alicante, Alicante, Spain

Corresponding author

Gerly Edson Guzmán-Calderón, MD
Av. Edgardo Rebagliati 490, Jesús María,
Lima, Peru
edson_guzman@hotmail.com

References

- [1] Anderloni A, Buda A, Vieceli F et al. Endoscopic ultrasound-guided transmural stenting for gallbladder drainage in high-risk patients with acute cholecystitis: a systematic review and pooled analysis. *Surg Endosc* 2016; 30: 5200–5208
- [2] Nasser A, Bill K, Barawi M. Lumen-apposing metal stent–assisted electrohydraulic lithotripsy and mechanical lithotripsy for cholelithiasis in a nonsurgical patient. *VideoGIE* 2019; 4: 159–160

- [3] Sievert C, Silvis S. Evaluation of electrohydraulic lithotripsy as a means of gallstone fragmentation in a canine model. *Gastrointest Endosc* 1987; 33: 233–235
- [4] Hochberger J, Gruber E, Wirtz P et al. Lithotripsy of gallstones by means of a quality-switched giant-pulse neodymium: yttrium-aluminum-garnet laser. Basic in vitro studies using a highly flexible fiber system. *Gastroenterology* 1991; 101: 1391–1398

Bibliography

Endoscopy 2021; 53: E415–E416

DOI 10.1055/a-1334-3970

ISSN 0013-726X

published online 14.1.2021

© 2021. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>