Endoscopic submucosal dissection for polyps in the anal transitional zone after ileoanal anastomosis in a patient with familial adenomatous polyposis

► **Fig. 1** Endoscopic views showing multiple coalescent polypoid lesions: **a** in the anal transitional zone; **b** at the ileoanal anastomosis.

► **Fig. 2** Endoscopic views showing: **a** severe fibrosis and surgical staples visible at the anastomotic site; **b** the post-endoscopic submucosal dissection ulcer in the anal transitional zone.

A 48-year-old man with a history of familial adenomatous polyposis (FAP) underwent prophylactic restorative proctocolectomy and ileal-pouch anal anastomosis 16 years ago and was subsequently lost to follow-up. He re-presented with rectal bleeding and endoscopy revealed multiple coalescent polypoid lesions (T0-Is + IIa), circumferentially occupying the anal transitional zone (► **Fig. 1 a**) and the area of the surgical anastomosis (► **Fig. 1 b**). Biopsy specimens showed a tubulovillous adenoma with low grade

dysplasia and focal high grade dysplasia. The patient was referred for treatment by endoscopic submucosal dissection (ESD). The procedure was performed in the endoscopy unit, with anesthetic support, using a gastroscope (GIF-HQ190; Olympus), FlushKnife BTS (Fujifilm), and IT Knife nano (Olympus). Severe fibrosis was found at the ileoanal anastomosis and multiple staples were identified beneath the tumor, some of which were removed during the procedure (► **Fig. 2 a**).

► **Fig. 3** Macroscopic appearance of the resected specimen, which measured 75 × 60 × 10 mm and showed coalescent polyps.

En bloc excision of the remaining rectal mucosa was successfully achieved (► **Fig. 2 b**; ► **Video 1**), producing a specimen that measured 75 × 60 × 10 mm (► **Fig. 3**). Pathology confirmed a tubulovillous adenoma with low and high grade dysplasia, with free tumor margins. No recurrence, stricture, or changes in bowel habit were noted at follow-up.

The ESD technique can remove gastrointestinal neoplasias, even when they are associated with severe fibrosis [1]. Although technically difficult, the authors suggest that ESD is a safe and effective treatment for complete resection of lesions associated with severe fibrosis, as found at the perianastomotic site. Restorative proctocolectomy and ileal-pouch anal anastomosis is the treatment of choice for patients with FAP and profuse adenomas, but may leave residual anal transitional zone mucosa that is prone to adenoma formation (incidence 10%–51%) [2]. Long-term annual endoscopic surveillance needs to be emphasized [3]. To the best of the authors' knowledge, this is the first case of ESD for coalescent polyps located in the anal

Video 1 Endoscopic submucosal dissection for coalescent polyps in the anal transitional zone after restorative proctocolectomy and ileal-pouch anal anastomosis in a patient with familial adenomatous polyposis.

transitional zone after restorative proctocolectomy and ileal-pouch anal anastomosis in a patient with FAP.

Endoscopy_UCTN_Code_TTT_1AQ_2AD

Competing interests

The authors declare that they have no conflict of interest.

The authors

Catarina O'Neill¹, **Pedro Barreiro^{1,2}**, **Rui Mendo¹**, **Catarina Félix¹**, **Iala Costa¹**, **Cristina Chagas¹**

- 1 Department of Gastroenterology, Centro Hospitalar de Lisboa Ocidental, Hospital de Egas Moniz, Lisbon, Portugal
- 2 Department of Gastroenterology, Hospital dos Lusíadas, Lisbon, Portugal

Corresponding author

Catarina O'Neill, MD

Department of Gastroenterology, Centro Hospitalar Lisboa Ocidental, Hospital de Egas Moniz, Lisbon, Portugal
oneill.catarina@gmail.com

References

- [1] Pimentel-Nunes P, Dinis-Ribeiro M, Ponchon T et al. Endoscopic submucosal dissection: European Society of Gastrointestinal Endoscopy (ESGE) Guideline. *Endoscopy* 2015; 47: 829–854
- [2] Ozdemir Y, Kalady MF, Aytac E et al. Anal transitional zone neoplasia in patients with familial adenomatous polyposis after restorative proctocolectomy and IPAA: incidence, management, and oncologic and functional outcomes. *Dis Colon Rectum* 2013; 56: 808–814

- [3] Syngal S, Brand RE, Church JM et al. ACG clinical guideline: Genetic testing and management of hereditary gastrointestinal cancer syndromes. *Am J Gastroenterol* 2015; 110: 223–263

Bibliography

Endoscopy 2022; 54: E11–E12

DOI 10.1055/a-1346-8903

ISSN 0013-726X

published online 16.2.2021

© 2021. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>