Endoscopic ultrasonography-guided pancreaticoduodenostomy with a lumen-apposing metal stent to treat main pancreatic duct dilatation

A 60-year-old woman was referred to our hospital for work-up of a mass in the pancreas with main pancreatic duct (MPD) dilatation. She had no symptoms, but had a history of abnormal glucose tolerance for 2 years. A contrast-enhanced computed tomography (CT) scan showed a low density mass with scattered contrast enhancement in the pancreatic head and dilatation of the distal MPD (► **Fig. 1**). Endoscopic ultrasonography (EUS) revealed a multilocular cyst with microcysts in the area, which was compressing the MPD and causing the MPD dilatation (► **Fig. 2**). The lesion was eventually diagnosed as a serous cystadenoma and was observed with regular imaging tests.

Over a period of 4 years, the MPD dilatation and atrophy of the pancreatic body and tail gradually progressed (► **Fig. 3**) and her glucose tolerance tests worsened. Because pancreaticoduodenectomy was judged to be too invasive, endoscopic ultrasonography-guided pancreaticoduodenostomy (EUS-PDS) using a lumen-apposing metal stent (LAMS) was scheduled, with her written informed consent, as a minimally invasive alternative.

After inserting an echoendoscope to the bulb of the duodenum, we saw remarkable MPD dilatation, with the appearance of a cyst measuring 4.2×4.8 cm at the pancreatic head, adjacent to the duodenum; a safe puncture route was confirmed. Thereafter, a skilled endoscopist punctured the MPD with cautery assistance and a LAMS was placed within 80 seconds and without adverse events (► **Video 1**). CT images 2 months after the procedure confirmed decompression of the MPD had been achieved (► **Fig. 4**) and this was sustained without adverse events 3 months after endoscopic removal of the LAMS (► **Fig. 5**).

EUS-guided pancreatic duct drainage, including by EUS-PDS, is one of the methods of endoscopic pancreatic duct drain-

► **Fig. 1** Contrast-enhanced computed tomography image showing a low density mass with scattered contrast enhancement in the pancreatic head (arrowhead) and dilatation of the distal main pancreatic duct (arrows).

► **Fig. 2** Endoscopic ultrasonography image showing a multilocular cyst with microcysts in the pancreatic head (arrowhead) and dilatation of the distal main pancreatic duct (arrows).

► **Fig. 3** Magnetic resonance image showing cystic main pancreatic duct dilatation (arrow) and atrophy of the pancreatic body and tail 4 years after the initial diagnosis.

age and has been used for MPD dilatation due to pancreatitis, pancreatic fistulas, or postoperative anastomotic strictures [1–4]. This is the first case reported with a serous cystadenoma and cystic MPD dilatation for which EUS-PDS was feasible and available. This therapy is minimally

invasive and could provide a new option for patients with MPD dilatation before pancreatic resection.

Endoscopy_UCTN_Code_TTT_1AS_2AD

▶ Video 1 Endoscopic ultrasonography-guided pancreaticoduodenostomy using a lumen-apposing metal stent for the treatment of cystic main pancreatic duct dilatation.

▶ Fig. 4 Contrast-enhanced computed tomography image 2 months after the procedure showing decompression of the main pancreatic duct.

▶ Fig. 5 Magnetic resonance image 3 months after endoscopic removal of the lumen-apposing metal stent showing the sustained decompression of the main pancreatic duct (arrow).

Competing interests

The authors declare that they have no conflict of interest.

The authors

Masaki Kuwatani , **Kosuke Nagai**, **Yunosuke Takishin** , **Ryutaro Furukawa**, **Hajime Hirata**, **Kazumichi Kawakubo**, **Naoya Sakamoto**
Department of Gastroenterology and Hepatology, Hokkaido University Hospital, Sapporo, Japan

Corresponding author

Masaki Kuwatani, MD
Department of Gastroenterology and Hepatology, Hokkaido University Hospital, North 14, West 5, Kita-ku, Sapporo 060-8648, Japan
mkuwatan@med.hokudai.ac.jp

References

- [1] Krafft MR, Nasr JY. Anterograde endoscopic ultrasound-guided pancreatic duct drainage: a technical review. *Dig Dis Sci* 2019; 64: 1770–1781

- [2] Zein CO, Baron TH, Morgan DE. Endoscopic pancreaticoduodenostomy for treatment of pancreatic duct disconnection because of severe acute pancreatitis. *Gastrointest Endosc* 2003; 58: 130–134
- [3] Arvanitakis M, Delhaye M, Bali MA et al. Endoscopic treatment of external pancreatic fistulas: when draining the main pancreatic duct is not enough. *Am J Gastroenterol* 2007; 102: 516–524
- [4] Oh D, Park DH, Cho MK et al. Feasibility and safety of a fully covered self-expandable metal stent with antimigration properties for EUS-guided pancreatic duct drainage: early and midterm outcomes. *Gastrointest Endosc* 2016; 83: 366–373

Bibliography

Endoscopy 2022; 54: E113–E114

DOI 10.1055/a-1408-0148

ISSN 0013-726X

published online 30.3.2021

© 2021. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>