Long-term endoscopic follow-up after closure of a post-bariatric surgery fistula with a cardiac septal defect occluder

▶ **Video 1** Long-term follow-up of successful gastrocutaneous fistula closure using a cardiac septal defect occluder.

▶ **Fig. 2** Image from an upper gastrointestinal series showing no evidence of gastrointestinal leakage or a fistula, and no evidence of the previously placed cardiac septal defect occluder.

▶ **Fig. 1** Endoscopic images showing: **a** the well-positioned cardiac septal defect occluder, with closure of the gastrocutaneous fistula at 1-year follow-up; **b** no evidence of leakage or a fistulous tract, despite apparent migration of the cardiac septal defect occluder, at 3-year follow-up.

Bariatric surgery is currently the most effective treatment strategy for obesity; however, post-surgical fistulas may occur in up to 8.3% of patients following traditional Roux-en-Y gastric bypass (RYGB) [1]. Currently endoscopic treatment of these complications remains challenging, with unsuccessful fistula closure occurring in 20% of patients [1,2]. More recently, the use of a cardiac septal defect occluder (CSDO) device has been pro-

posed as a novel treatment for the closure of gastrointestinal fistulas. A CSDO is a double-disc self-expanding closure device made of nitinol and interwoven polyester. The successful use of CSDOs in the management of gastrointestinal surgical and bariatric leaks has been reported; however, there are limited data regarding long-term outcomes [3,4]. In this video, we describe the successful closure of a gastrocutaneous fistula using

a CSDO and demonstrate persistent closure at long-term follow-up (▶ **Video 1**). A 36-year-old man with a history of RYGB 3 years previously presented to our institution with a gastrocutaneous fistula. Initial treatment with a fully covered self-expandable metal stent (SEMS) was unsuccessful, and the patient subsequently underwent treatment with a CSDO [5]. The CSDO procedure was immediately successful with closure of the gastrocutaneous fistula and, at 1-year follow-up, endoscopy demonstrated an intact, well-positioned CSDO, with no evidence of a fistula (▶ **Fig. 1 a**). At 3-year follow-up, routine endoscopy did not identify the CSDO (▶ **Fig. 1 b**). An upper gastrointestinal contrast study was performed and confirmed that the CSDO was not present in the gastrointestinal lumen or intra-abdominal cavity (▶ **Fig. 2**). Despite the presumed migration of the device, the fistula remained closed with no recurrent fistula or leakage noted. In summary, the use of a CSDO device appears to be a safe and effective long-

term treatment for patients with post-surgical gastrocutaneous fistulas. While more data are needed to verify these results, CSDOs may be a feasible alternative for gastrointestinal leaks and fistulas that are refractory to traditional endoscopic techniques.

Endoscopy_UCTN_Code_TTT_1AO_2AI

Competing interests

The authors declare that they have no conflict of interest.

The authors

Diogo Turiani Hourneaux de Moura¹, **Mateus Bond Boghossian¹**, **Bruno Salomão Hirsch¹**, **Thomas R. McCarty²**, **Alberto Jose Baptista³**, **Eduardo Guimarães Hourneaux de Moura¹**

- 1 Gastrointestinal Endoscopy Unit, Hospital das Clínicas da Faculdade de Medicina da Universidade de São Paulo, São Paulo, Brazil
- 2 Gastroenterology, Hepatology, and Endoscopy Division, Harvard Medical School, Brigham and Women's Hospital, Boston, Massachusetts, USA
- 3 Department of Endoscopy, Hospital de Clínicas Caracas, San Bernardino, Venezuela

Corresponding author

Mateus Bond Boghossian, MD

Av. Dr Enéas de Carvalho Aguiar, 225,
6º andar, bloco 3, Cerqueira César,
São Paulo, SP, 05403-010, Brazil
mateus.boghossian@hc.fm.usp.br

References

- [1] Okazaki O, Bernardo WM, Brunaldi VO et al. Efficacy and safety of stents in the treatment of fistula after bariatric surgery: a systematic review and meta-analysis. *Obes Surg* 2018; 28: 1788–1796
- [2] de Moura DTH, Sachdev AH, Thompson CC. Endoscopic full-thickness defects and closure techniques. *Curr Treat Options Gastroenterol* 2018; 16: 386–405
- [3] Baptista A, de Moura DTH, Jirapinyo P et al. Efficacy of the cardiac septal occluder in the treatment of post-bariatric surgery leaks and fistulas. *Gastrointest Endosc* 2019; 89: 671–679.e1
- [4] De Moura DTH, Baptista A, Jirapinyo P et al. Role of cardiac septal occluders in the treatment of gastrointestinal fistulas: a systematic review. *Clin Endosc* 2020; 53: 37–48
- [5] de Moura DTH, Ribeiro IB, Funari MP et al. Novel use of a cardiac septal occluder to treat a chronic recalcitrant bariatric fistula after Roux-en-Y gastric bypass. *Endoscopy* 2019; 51: E111–E112

Bibliography

Endoscopy 2022; 54: E127–E128

DOI 10.1055/a-1422-2631

ISSN 0013-726X

published online 16.4.2021

© 2021. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>