Endoscopic retrieval of a migrated surgical clip in a choledochojejunal anastomosis using the rendezvous technique

It is a rare complication for a surgical clip to migrate into a bile duct [1, 2], and removing a migrated surgical clip endoscopically is technically challenging, particularly in patients with surgically altered anatomy. We report here the first successful endoscopic removal of a migrated surgical clip using the rendezvous technique of percutaneous transhepatic biliary drainage (PTBD) and double-balloon endoscope (DBE)-assisted endoscopic retrograde cholangiopancreatography (ERC) (► **Video 1**).

A 35-year-old woman with a history of congenital biliary dilatation who had undergone extrahepatic bile duct resection and choledochojejunostomy 8 months previously was admitted for obstructive jaundice. Computed tomography and magnetic resonance cholangiopancreatography showed intrahepatic biliary dilatation and anastomotic obstruction of the choledochojejunal anastomosis (► **Fig. 1 a, b**). We attempted ERC assisted by a short-type DBE (EI-580BT; Fujifilm, Tokyo, Japan). Insertion to the blind end and discovery of the choledochojejunal anastomosis were successful; however, endoscopic imaging showed obstruction with a migrated surgical clip in the choledochojejunal anastomosis (► **Fig. 1 c**).

It was difficult to gain access into the bile duct beyond the clip, and at this point we gave up the attempt at endoscopic treatment. Four days later, we tried PTBD and succeeded in guiding the tip of a PTBD catheter beyond the clip to the jejunal side of the obstruction. As the patient firmly refused to undergo further surgery, however, we reattempted DBE-as-

sisted ERC. The endoscopic catheter successfully accessed the bile duct using the PTBD catheter (► **Fig. 2**). After dilation using an 8-mm biliary dilation balloon (REN; Kaneka, Osaka, Japan) was successful, the migrated surgical clip was carefully removed with forceps, allowing placement of two endoscopic biliary drainage tubes consisting of 5-Fr plastic stents.

► **Video 1** Endoscopic retrieval of a migrated surgical clip in a choledochojejunal anastomosis using the rendezvous technique: percutaneous transhepatic bile duct drainage and double-balloon endoscope-assisted endoscopic retrograde cholangiopancreatography.

► **Fig. 1** Intrahepatic biliary dilation and anastomotic obstruction of a choledochojejunal anastomosis: **a** computed tomography, **b** magnetic resonance cholangiopancreatography. **c** Short-type double-balloon endoscopic imaging showed obstruction due to a migrated surgical clip in the choledochojejunal anastomosis.

► **Fig. 2 a, b** Endoscopic catheter successfully accessed the bile duct by means of the percutaneous transhepatic biliary drainage catheter, using the rendezvous technique.

The DBE-assisted ERC-PTBD rendezvous technique has been reported to be effective [3–5]. A combined PTBD procedure which exerts force on the tip of the catheter may also be effective, especially if endoscopic treatment for postoperative mechanical obstruction is unsuccessful.

Endoscopy_UCTN_Code_TTT_1AO_2AL

Competing interests

The authors declare that they have no conflict of interest.

The authors

Takashi Ito, Masaaki Shimatani, Masataka Masuda, Koh Nakamaru, Toshiyuki Mitsuyama, Makoto Takaoka, Makoto Naganuma

Kansai Medical University, The Third Department of Internal Medicine, Division of Gastroenterology and Hepatology, Hirakata, Japan

Corresponding author

Masaaki Shimatani, MD, PhD

The Third Department of Internal Medicine, Division of Gastroenterology and Hepatology, Kansai Medical University, 2-5-1 Shinmachi, Hirakata, Osaka, Japan
shimatam@takii.kmu.ac.jp

References

- [1] Chong VH, Chong CF. Biliary complications secondary to post-cholecystectomy clip migration: review of 69 cases. *J Gastrointest Surg* 2010; 14: 688–696
- [2] Daniel S, Oren G, Barch O et al. Cholangitis caused by surgical clip migration into the common bile duct: a rare complication of a daily practice. *BMJ Case Rep* 2020; 13: e231201
- [3] Mönkemüller K, Popa D, McGuire B et al. Double-balloon enteroscopy-ERCP rendezvous technique. *Endoscopy* 2013; 45: E333–E334
- [4] Bokemeyer A, Müller F, Niesert H et al. Percutaneous-transhepatic-endoscopic rendezvous procedures are effective and safe in patients with refractory bile duct obstruction. *United European Gastroenterol J* 2019; 3: 397–404

- [5] Yamauchi H, Kida M, Imaizumi H et al. Innovations and techniques for balloon-enteroscopy-assisted endoscopic retrograde cholangiopancreatography in patients with altered gastrointestinal anatomy. *World J Gastroenterol* 2015; 21: 6460–6469

Bibliography

Endoscopy 2022; 54: E182–E183

DOI 10.1055/a-1463-2618

ISSN 0013-726X

published online 12.5.2021

© 2021. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>