Double-pigtail plastic stent is effective as a rescue therapy after obstruction of choledochoduodenostomy using a lumen-apposing metal stent

► **Fig. 1** Gastroscopy revealed invasion of the duodenal stent and obstruction of the proximal flange of the lumen-apposing metal stent with bezoars.

► **Fig. 2** Double-pigtail plastic stent in the lumen-apposing metal stent (view from the gastric antrum with transpyloric duodenal stent).

Endoscopic ultrasound (EUS)-guided choledochoduodenostomy using a lumen-apposing metal stent (LAMS) is an effective alternative when endoscopic retrograde cholangiopancreatography fails or is impossible [1–3]. However, although the technique opens up new therapeutic possibilities, it is also associated with new complications, such as obstruction, for which an optimal treatment has not yet been clearly established [4].

We report the case of a 62-year-old patient with metastatic pancreatic adenocarcinoma of the liver exhibiting duode-

► **Video 1** Double-pigtail plastic stent placement is an effective rescue therapy after obstruction of a lumen-apposing metal stent placed during choledochoduodenostomy.

nal invasion. Initial management featured the insertion of a 6-mm LAMS (Hot AXIOS; Boston Scientific, Marlborough, Massachusetts, USA) to allow biliary drainage and a metallic duodenal stent (Wallflex; Boston Scientific) to treat the parietal and ampullary infiltration.

After 1 month, the patient presented with fever, abdominal pain, and jaundice suggestive of a LAMS obstruction. Gastroscopy revealed invasion of the duodenal stent and that the proximal flange of the LAMS was obstructed by food bezoars (► **Fig. 1**). After the obstruction was cleared using a lavage pump, we placed a 7-Fr, 10-cm double-pigtail biliary stent (Boston Scientific) inside the LAMS (► **Fig. 2**, ► **Video 1**).

Opacification evident at the end of the procedure confirmed permeability of the biliary and duodenal prostheses (► **Fig. 3**). Over the following days, the biological jaundice and hyperthermia decreased, allowing resumption of chemotherapy.

The addition of a double-pigtail, plastic biliary stent efficiently ensures LAMS patency, especially in cases of duodenal

► **Fig. 3** Opacification showing permeability of the biliary and duodenal prostheses.

stenosis associated with risks of recurrent obstruction caused by bezoar impaction.

Endoscopy_UCTN_Code_TTT_1AS_2AD

Competing interests

J. Albouys is a consultant for Biogen. R. Legros is a consultant for Olympus and Boston Scientific and has received (ESD Training) from Erbe. V. Loustaud-Ratti is a speaker for Gilead and Abbvie. M. Pioche is a consultant for Olympus, Boston Scientific and 3D matrix and has received ESD training from Cook Medical and Olympus. J. Jacques is a consultant for Olympus, Boston Scientific and has received ESD Training from Erbe.

The authors

Jérémie Albouys¹, Sophie Geyl¹, Benjamin Redelsperger¹, Véronique Loustaud-Ratti¹, Romain Legros¹, Mathieu Pioche², Jérémie Jacques¹

- 1 Gastroentérologie et endoscopie digestive, CHU Dupuytren, Limoges, France
- 2 Unité d'endoscopie digestive, Service de Gastroentérologie, Hôpital Edouard Herriot, Hospices Civils de Lyon, Lyon, France

Corresponding author

Jérémie Albouys, MD

Service d'Hépatogastro-entérologie, CHU Dupuytren, 2 Avenue Martin Luther King, 87042 Limoges, France
Fax: +33-5-55-05-87-33
jeremie.albouys@gmail.com

References

- [1] Jacques J, Privat J, Pinard F et al. EUS-guided choledochoduodenostomy by use of electrocautery-enhanced lumen-apposing metal stents: a French multicenter study after implementation of the technique (with video). *Gastrointest Endosc* 2020; 92: 134–141. doi:10.1016/j.gie.2020.01.055
- [2] Krishnamoorthi R, Dasari CS, Thoguluva Chandrasekar V et al. Effectiveness and safety of EUS-guided choledochoduodenostomy using lumen-apposing metal stents (LAMS): a systematic review and meta-analysis. *Surg Endosc* 2020; 34: 2866–2877. doi:10.1007/s00464-020-07484-w
- [3] Jain D, Shah M, Patel U et al. Endoscopic ultrasound-guided choledochenterostomy by using lumen-apposing metal stent in patients with failed endoscopic retrograde cholangiopancreatography: a literature review. *Digestion* 2018; 98: 1–10. doi:10.1159/000487185
- [4] Fugazza A, Gabbiadini R, Colombo M et al. Self-expandable metal stent placement as a rescue procedure for lumen-apposing metal stent misdeployment in biliary drainage. *Endoscopy* 2020; 52: E430–E431. doi:10.1055/a-1158-8749

Bibliography

Endoscopy 2022; 54: E184–E185

DOI 10.1055/a-1471-1444

ISSN 0013-726X

published online 12.5.2021

© 2021. Thieme. All rights reserved.

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>