Endoscopic submucosal dissection of a subepithelial neuroendocrine tumor located in the duodenum using a countertraction technique

Endoscopic submucosal dissection (ESD) is the technique of choice to remove large superficial neoplasms of the esophagus, stomach, colon, and rectum [1]. In the duodenum, for non-ampullary lesions, it has been associated with a high risk of perforation [2] owing to the relative thinness of the duodenal wall. Data with regard to ESD for subepithelial lesions are scarce. One study described a low rate of curative treatment for duodenal neuroendocrine tumors (NETs), but few patients were included [3]. Techniques such as countertraction using a rubber band [4] and anchoring the clips to close the scar [5] can be considered to secure the procedure.

We herein report the case of a 58-year-old man with a non-ampullary duodenal subepithelial NET of 9mm that was removed by ESD (► **Video 1**). First, a circumferential incision was performed within the mucosa. Countertraction with two clips and a rubber band was then used to facilitate dissection of the lesion (► **Fig. 1 a**). The lesion was removed en bloc. After the procedure, the scar was closed by anchoring the clips: small mucosal incisions were performed around the scar (► **Fig. 1 b**) to allow better grip for the clip at each edge of the scar with-

► **Video 1** Endoscopic submucosal dissection of a subepithelial neuroendocrine tumor located in the duodenum using a countertraction technique.

out their slipping (► **Fig. 1 c**). The patient was discharged the day after the procedure without experiencing any adverse events.

The pathology report revealed a well-differentiated NET, G2 (Ki-67 index 4%; mitotic count 1), without perineural invasion but with lymphatic emboli. The NET was completely resected by the procedure. Surveillance was determined to be

the appropriate further management by a NET multidisciplinary team.

This case highlights the feasibility of ESD for subepithelial NETs located in the duodenum. It also illustrates the countertraction technique and closure of the scar by anchoring the clips to secure the procedure. This must however only be done in highly selected patients and by skilled endoscopists.

► **Fig. 1** Endoscopic views showing: **a** countertraction with two clips and a rubber band being used to facilitate dissection of the lesion; **b** small mucosal incisions that were made around the scar to allow better grip for the clips at each edge of the scar without their slipping; **c** the appearance of the closed scar.

Competing interests

The authors declare that they have no conflict of interest.

The authors

Thomas Lambin^{1,2}, Clara Yzet³, Jérémie Albouys⁴, Pierre Lafeuille¹, Florian Rostain¹, Jérôme Rivory¹, Mathieu Pioche^{1,2}

- 1 Gastroenterology and Endoscopy Unit, Pavillon L, Edouard Herriot Hospital, Lyon, France
- 2 Inserm U1032, Labtau, Lyon, France
- 3 Department of Gastroenterology, Amiens University Hospital, Amiens, France
- 4 Gastroenterology and Endoscopy Unit, CHU Dupuytren, Limoges, France

Corresponding author

Thomas Lambin, MD
Endoscopy Unit – Digestive Disease
Department, Pavillon L – Hôpital Edouard
Herriot, 69437 Lyon, France
thomaslambin@hotmail.fr

References

- [1] Pimentel-Nunes P, Dinis-Ribeiro M, Ponchon T et al. Endoscopic submucosal dissection: European Society of Gastrointestinal Endoscopy (ESGE) Guideline. *Endoscopy* 2015; 47: 829–854
- [2] Watanabe D, Hayashi H, Kataoka Y et al. Efficacy and safety of endoscopic submucosal dissection for non-ampullary duodenal polyps: A systematic review and meta-analysis. *Dig Liver Dis* 2019; 51: 774–781
- [3] Kobara H, Miyaoka Y, Ikeda Y et al. Outcomes of endoscopic submucosal dissection for subepithelial lesions localized within the submucosa, including neuroendocrine tumors: a multicenter prospective study. *J Gastrointest Liver Dis* 2020; 29: 41–49
- [4] Jacques J, Charissoux A, Bordillon P et al. High proficiency of colonic endoscopic submucosal dissection in Europe thanks to countertraction strategy using a double clip and rubber band. *Endosc Int Open* 2019; 7: E1166–E1174
- [5] Otake Y, Saito Y, Sakamoto T et al. New closure technique for large mucosal defects after endoscopic submucosal dissection of colorectal tumors (with video). *Gastrointest Endosc* 2012; 75: 663–667

Bibliography

Endoscopy 2022; 54: E230–E231
DOI 10.1055/a-1492-2362
ISSN 0013-726X
published online 31.5.2021
© 2021. Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>