A Chiral Sulfoxide-Based C–H Acid

Denis Höfler^a

Karl Kaupmees^b

Ivo Leito^b

Benjamin List^{*a}

^a Max-Planck-Institut für Kohlenforschung, Kaiser-Wilhelm-Platz 1, 45470, Mülheim an der Ruhr, Germany
list@kofo.mpg.de

^b University of Tartu, Institute of Chemistry, Ravila 14a, 50411 Tartu, Estonia

Published as part of the Editorial Board Cluster

Received: 17.09.2021

Accepted after revision: 04.11.2021

Published online: 04.11.2021

DOI: 10.1055/a-1695-4516; Art ID: st-2021-u0339-c

License terms:

© 2021. The Author(s). This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Abstract We report the design and synthesis of a strong, chiral, enantiopure sulfoxide-based C–H acid. Single-crystal X-ray analysis confirms the proposed structure and its absolute configuration. The new motif shows a high acidity and activity in Brønsted and Lewis acid catalyzed transformations. So far, only little to no enantioselectivities were achieved.

Key words sulfoxide C–H acids, stereogenic sulfur, triflyl groups, strong acids, noncoordinating anions, Brønsted acids

Chiral binaphthyl-derived acids have shown great success in asymmetric Lewis and Brønsted acid catalysis,¹ especially confined variants.² However, their catalytic activity is inherently limited by the electron-rich binaphthyl system, which also limits their acidity and catalytic reactivity. With both enantiomers readily available, chiral sulfur-stereogenic sulfoxides are attractive ligands in transition-metal catalysis.³ In organocatalysis, a stereogenic sulfur has been either a contributing factor or exclusively responsible for high enantioselectivities when using weakly acidic chiral urea- or thiourea-derived catalysts.^{3,4} We envisioned a new, tris(triflyl)methane (**2**)-inspired motif with the acidic proton very close to the stereogenic sulfur atom, which we hypothesized could lead to efficient asymmetric induction. These considerations led to the design of **1**, expected to be a very strong C–H acid, with two triflyl (SO₂CF₃) groups⁶ and one chiral sulfoxide moiety (Scheme 1A). Indeed, a synthesis was developed, from commercially available iodide **3**,

Scheme 1 Design (A), synthesis (B), and application (C) of the chiral, enantiopure sulfoxide C–H acid. TMP = 2,2,6,6-tetramethylpiperidine.

which was converted into a diastereomeric mixture of two oxazolidinones **6** by following reported procedures.⁷ The major diastereomer (**6a**) was separated by flash chromatography and converted into the desired enantiopure sulfoxide acid **1** by treatment with bis(triflyl)methane in the presence of a strong base followed by H₂SO₄ acidification.⁸ With the desired C–H acid **1** in hand, we were able to assign its absolute configuration by X-ray single-crystal structure analysis of its hydroxonium hydrate (see Supporting Information).⁹

Further, an experimental p*K*_{ip} value of -12.5 ± 0.5 (in 1,2-dichloroethane, relative to picric acid) was determined for **1**. The corresponding free-ion p*K*_a value for a molecule of this size is expected to be essentially the same.¹⁰ This acidity corresponds to a p*K*_a of around 0 in acetonitrile.¹¹ Therefore, to the best of our knowledge, sulfoxide **1** can be considered to be the strongest enantiopure Brønsted acid that has been prepared so far. We applied acid **1** as a catalyst in a variety of different reactions including two Mukaiyama aldolizations and a Hosomi–Sakurai allylation (Scheme 1C). Although the catalytic activity was promising, little to no enantioselectivity was observed in all cases. In the future, further modifications of this easily accessible motif to increase its enantiodiscrimination are envisioned as well as its potential applications as an anionic ligand in transition-metal catalysis.

Conflict of Interest

The authors declare no conflict of interest.

Funding Information

Generous support by the European Research Council (Advanced Grant ‘C–H Acids for Organic Synthesis, CHAOS’) and the Deutsche Forschungsgemeinschaft (Leibniz Award to B.L. and Cluster of Excellence RESOLV, EXC 1069) is gratefully acknowledged. Work at Tartu was supported by the Estonian Research Council grant (PRG690), and by the EU through the European Regional Development Fund under project TK141 (2014–2020.4.01.15–0011).

Acknowledgment

We also thank Petra Wedemann, Diana X. Sun, Jonas Aronow, Lucas Schreyer, and Hyejin Kim for experimental assistance, as well as the members of our analytical departments for their excellent service, especially Dr. Christophe Farès, Dr. Richard Goddard, and Nils Nöthling. We acknowledge DESY (Hamburg, Germany), a member of the Helmholtz Association HGF, for the provision of experimental facilities; parts of this research were carried out at PETRA III, and we would like to thank Sofiane Saouane for excellent assistance in using the P11–High-Throughput Macromolecular Crystallography Beamline.

Supporting Information

Supporting information for this article is available online at <https://doi.org/10.1055/a-1695-4516>.

References and Notes

- (1) (a) Akiyama, T. *Chem. Rev.* **2007**, *107*, 5744. (b) Akiyama, T.; Mori, K. *Chem. Rev.* **2015**, *115*, 9277. (c) Terada, M.; Kanomata, K. *Synlett* **2011**, 1255.
- (2) (a) Mitschke, B.; Turberg, M.; List, B. *Chem* **2020**, *6*, 2515. (b) Schreyer, L.; Properzi, R.; List, B. *Angew. Chem. Int. Ed.* **2019**, *58*, 12761.
- (3) Trost, B. M.; Rao, M. *Angew. Chem. Int. Ed.* **2015**, *54*, 5026.
- (4) (a) Robak, M. T.; Trincado, M.; Ellman, J. A. *J. Am. Chem. Soc.* **2007**, *129*, 15110. (b) Kimmel, K. L.; Weaver, J. D.; Lee, M.; Ellman, J. A. *J. Am. Chem. Soc.* **2012**, *134*, 9058; corrigendum: *J. Am. Chem. Soc.* **2012**, *134*, 11828.
- (5) Turowsky, L.; Seppelt, K. *Inorg. Chem.* **1988**, *27*, 2135.
- (6) (a) Höfler, D.; van Gemmeren, M.; Wedemann, P.; Kaupmees, K.; Leito, I.; Leutzsch, M.; Lingnau, J. B.; List, B. *Angew. Chem. Int. Ed.* **2017**, *56*, 1411. (b) Höfler, D.; Goddard, R.; Nöthling, N.; List, B. *Synlett* **2019**, *30*, 433. (c) Gatzemeier, T.; van Gemmeren, M.; Xie, Y.; Höfler, D.; Leutzsch, M.; List, B. *Science* **2016**, *351*, 949. (d) Gheewala, C. D.; Collins, B. E.; Lambert, T. H. *Science* **2016**, *351*, 961.
- (7) (a) Wang, X.-J.; Liu, J.-T. *Tetrahedron* **2005**, *61*, 6982. (b) Liu, L.-J.; Chen, L.-J.; Li, P.; Li, X.-B.; Liu, J.-T. *J. Org. Chem.* **2011**, *76*, 4675.
- (8) **1-[[Bis(triflyl)methyl]sulfinyl]nonafluorobutane (1)**
A Schlenk flask was charged with bis(triflyl)methane (0.46 g, 1.6 mmol, 1.0 equiv) and THF (3.5 mL) to give a colorless clear solution that was cooled to -78°C . A 1.2 M solution of TMP-MgCl·LiCl in THF (3.1 mL, 3.8 mmol, 2.3 equiv) was added dropwise, followed by the diastereomerically pure sulfinyl oxazolidinone **6a** (4.3 g, 14 mmol, 1.0 equiv; dr >99:1) in THF (3.5 mL). The mixture was allowed to reach RT overnight. All volatiles were then removed under reduced pressure to give an orange solid that was dissolved in CH₂Cl₂ (30 mL) and washed with sat. aq. NaHCO₃ (3 × 10 mL). The pooled aqueous phases were extracted with CH₂Cl₂ (2 × 10 mL). The pooled organic phases were washed with concd aq HCl (3 × 30 mL), dried (MgSO₄), filtered, and concentrated under reduced pressure. The resulting yellowish viscous oil was dissolved in CH₂Cl₂ (30 mL), washed with concd H₂SO₄ (3 × 10 mL), stirred over dried BaCl₂ for 80 min, and filtered. All volatiles were removed under reduced pressure until 7 mL of liquid remained. This solution was stored at -29°C overnight, which led to the formation of a precipitate. The mother liquor was removed to give a colorless solid; yield: 0.51 g (57%, 0.93 mmol).
LC/MS (chiral): (150 mm Chiralpak IC-3, 4.6 mm i.d., 30:70 MeCN–0.2% TFA, 1.0 mL/min, 20.8 MPa, 298 K): *t*_{R(S)-enantiomer} = 21.06 min, *t*_{R(R)-enantiomer} = 22.69 min; *e.r.* = 1:99. ¹³C NMR (151 MHz, acetone-*d*₆): δ = 123.74 (t, *J* = 23.1 Hz, C₆), 120.22 (q, *J* = -322.0 Hz, C_{7,8}), 117.14 (tt, *J* = 322.7, 36.6 Hz, C₁), 114.51 (qt, *J* = 287.0, 32.5 Hz, C₄), 111.21 (tp, *J* = 266.2, 34.0 Hz, C₂), 108.76 (th, *J* = 271.0, -35 Hz, C₃). ¹⁹F NMR (471 MHz, acetone-*d*₆): δ = -179.16 (br, 3 F, *F*_{7,8}), -80.68 (br, 3 F, *F*_{7,8}), -81.37 to -82.51 (td, *J* = 9.6, 5.1 Hz, 3 F, *F*₄), -103.82 (br d, *J* = 220.6 Hz, 1 F, *F*₁), -121.02 (dt, *J* = 221.8, 10.3 Hz, 1 F, *F*₁), -123.63 [br d (AB system), *J* = -300.0 Hz, 1 F, *F*₂], -124.33 [br d (AB system), *J* = -300.0 Hz, 1 F, *F*₂], -126.39 [br d (AB system), *J* = 293.0 Hz, 1 F, *F*₃], -127.19 [br d (AB system), *J* = 293.0 Hz, 1 F, *F*₃]. MS (ESI[−]): *m/z* = 545 [M − H].

HRMS (ESI[−]): m/z [M − H] calcd for C₇F₁₅O₅S₃: 544.8674; found: 544.8674. Anal.: Calcd for C₇HF₁₅O₅S₃ (546.24 g/mol): C, 15.39; H, 0.18; F, 52.17; S, 17.61; Found: C, 15.42; H, 0.17; F, 52.14; S, 17.60.

- (9) CCDC 2106642 and 2106643 contain the supplementary crystallographic data for **1** hydroxonium hydrate and **6a**. The data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/structures

- (10) Paenurk, E.; Kaupmees, K.; Himmel, D.; Kütt, A.; Kaljurand, I.; Koppel, I. A.; Krossing, I.; Leito, I. *Chem. Sci.* **2017**, *8*, 6964.
 (11) Kütt, A.; Rodima, T.; Saame, J.; Raamat, E.; Mäemets, V.; Kaljurand, I.; Koppel, I. A.; Garlyauskayte, R. Y.; Yagupolskii, Y. L.; Yagupolskii, L. M.; Bernhardt, E.; Willner, H.; Leito, I. *J. Org. Chem.* **2011**, *76*, 391.