A homemade endoscopic guillotine strikes again: removal of a nasogastric tube mysteriously knotted in the proximal third of the esophagus

► **Fig. 1** Chest radiograph showing the nasogastric tube forming a knot (arrow) in the upper third of the esophagus.

A 63-year-old woman with a past history of gastric bypass a few years ago was referred to our hospital with a mechanical ileus. A 14-Fr Salem nasogastric tube (NGT) was already in place for abdominal decompression at the time of admission. An ileal resection including a Meckel's diverticulum was performed with an ileo-ileal laterolateral mechanical anastomosis. Postoperatively NGT removal was impossible owing to strong resistance and the patient experiencing thoracic pain. A chest radiograph (► **Fig. 1**) and an upper gastrointestinal endoscopy (► **Fig. 2a**) showed coiling of the distal end of the tube, which had formed a knot in the proximal esophagus. Endoscopic removal of the NGT was attempted with the patient under general anesthesia. The NGT knot was pushed down into the gastric pouch to give more space for endoscopic maneuvers but, despite this, the knot could not be untied.

It was therefore decided to cut the knot using an “endoscopic guillotine,” as has been previously described for plastic stents [1] and a 14-Fr NGT [2]. In order to avoid accidental interposition of gastric,

esophageal, or small-intestinal mucosal wall during the cutting phase, we opted for a technical variant that has previously been described for plastic stents [3] and migrated gastric band trimming [4, 5]. A sphincterotome (CleverCut3V; Olympus) was used to thread a 0.035-inch, 450-cm, straight-tip guidewire (Dreamwire; Boston Scientific) through the knot in the NGT (► **Fig. 2b**) and the distal end of the guidewire was retrieved using a standard biopsy forceps (Endojaw Large; Olympus). Both ends of the guidewire were then locked into a mechanical lithotripsy device (Olympus) (► **Fig. 3**) and the crank handle of the lithotripter was gradually turned until the NGT was completely transected (► **Video 1**). The proximal part of the NGT was gently removed by pulling it through the patient's nose. The three remaining sections of the NGT were retrieved using a raptor grasping device (Steris) (► **Fig. 4**). No relevant iatrogenic lesions were observed at endoscopy, and no peri- or post-procedural complications occurred.

Endoscopy_UCTN_Code_CPL_1AH_2AJ

► **Fig. 2** Endoscopic views showing: **a** the knot in the nasogastric tube; **b** the guidewire wrapped around the knot.

► **Fig. 3** Photograph of the crank handle of the lithotripter being turned during the procedure.

Competing interests

The authors declare that they have no conflict of interest.

Video 1 A sphincterotome was used to thread a guidewire through the knot in the nasogastric tube (NGT) and the end of the guidewire was retrieved, with both ends then being locked into a mechanical lithotripsy device. The crank handle was turned until the NGT was transected and the parts of the NGT were then removed through the patient's nose and mouth.

Fig. 4 Photograph showing two of the three pieces of the nasogastric tube that were removed through the patient's mouth.

The authors

Fabio Pinto do Couto¹, Aurélie Cavin¹, Olivier Pittet¹, Paraskevi Archanioti², Sébastien Godat³, Domenico Galasso^{2,3}

- 1 Department of Surgery, Riviera-Chablais Hospital, Rennaz, Switzerland
- 2 Gastroenterology Unit, Riviera-Chablais Hospital, Rennaz, Switzerland
- 3 Department of Gastroenterology and Hepatology, Vaud University Hospital Centre (CHUV), Lausanne, Switzerland

Corresponding author

Aurélie Cavin, MD

Department of Surgery, Riviera-Chablais Hospital, Route du Vieux Séquoia 20, Rennaz (Vaud), CH1847, Switzerland
aurelie.cavin@hopitalrivierachablais.ch
aurelie.cavin@chuv.ch

References

- [1] Mutignani M, Dokas S, Perri V et al. Post-insertion tailoring of plastic biliary stents: A novel technique. *Endoscopy* 2006; 38: 856
- [2] Galasso D, Tringali A, Riccioni ME et al. Use of a homemade endoscopic guillotine to cut through a nasogastric tube inadvertently sutured to a colon-gastric anastomosis. *Endoscopy* 2013; 45: E413–E414
- [3] Mutignani M, Salerno R, Perri V et al. Removal of a doubly impacted plastic stent: A new technique. *Endoscopy* 2008; 40: E215–E216
- [4] Gutiérrez JP, Acosta F, Geninazzi H et al. Removal of a migrated gastric band using the Soehendra bile stone lithotripter. *Endoscopy* 2013; 45: E371–E372
- [5] de Sousa Damião F, Noronha Ferreira C, Nogueira F et al. Endoscopic removal of a migrated gastric band. *Endoscopy* 2021; 53: S189

Bibliography

Endoscopy 2022; 54: E676–E677
DOI 10.1055/a-1738-9112
ISSN 0013-726X
published online 18.2.2022
© 2022. Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany

ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>

Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>