


Successful closure of a refractory gastrobronchial fistula using endoscopic mucosal ablation followed by single loop-and-clips technique


OPEN
ACCESS

A 78-year-old man developed a late fistula between the gastropasty pouch and right bronchus following Ivor Lewis esophagectomy for esophageal neoplasia (► **Fig. 1**). He was unable to ingest food orally, necessitating the placement of a permanent percutaneous jejunostomy. He had a persistent cough, and the fistula was refractory to various endoscopic interventions, including placement of clips, tissue adhesives, and stents.

In the first session, the fistula was treated using a combination of endoscopic submucosal dissection (ESD) and over-the-scope (OTS) clip. ESD was performed around and inside the fistula tract, aiming to create a 1-cm mucosal patch that was centrally positioned at the orifice (► **Video 1**). To extract the fistula from the wall, traction was applied to the mucosal flap using the clip-with-line traction technique, enabling deeper dissection of the fistula tract. Finally, an OTS clip was deployed to close the orifice. However, after a period of 7 days, the patient experienced a recurrence of symptoms due to the detachment of the OTS clip from the gastrointestinal wall. During the second session, a persistent 5-mm fibrotic fistula orifice was observed. To address this, a combination of mucosal ablation and the single loop-and-clips technique (“King” closure) was performed (► **Fig. 2**). Endoscopic mucosal ablation was performed using argon plasma coagulation targeting the intra-fistular mucosa and a surrounding circumferential area to eliminate the mucosal scar tissue and promote the healing of the fistular orifice (► **Fig. 1**). Next, the King closure technique was successfully applied using five clips and a coaxial polyloop. Immediate post-procedure capnography did not detect the presence of CO₂. The follow-up at 8 weeks after the procedure confirmed fistula resolution based on the absence of symptoms and no contrast leakage on




► **Fig. 1** Endoscopic images of the post-esophagectomy fistula **a** before and **b** after endoscopic mucosal ablation using argon plasma coagulation.



► **Video 1** Ultimately successful closure of a refractory gastrobronchial fistula using the combination of endoscopic submucosal dissection (ESD) and application of an over-the-scope (OTS) clip, followed later by argon ablation and single loop-and-clip technique.

imaging (► **Fig. 3**). At the 1-year follow-up, the patient remained asymptomatic. The management of esophagobronchial and gastrobronchial fistulas is challenging. Recently, some French investigators have reported the use of ESD followed by the application of an OTS clip for the treatment of severe fistulas. However, this method can fail in some cases [1–3].

Studies have reported that the King closure is a safe technique and provides superior long-term histological healing outcomes compared with OTS clipping [4–5]. In conclusion, peristitular mucosal ablation followed by the single loop-and-clips technique appears to be an effective approach for the treatment of complex fistulas.



► **Fig. 2** a, b Gastrobronchial fistula closure using the single-loop-and-clips technique.



► **Fig. 3** Esophagogram showing no signs of persistence of fistula.

Endoscopy_UCTN_Code_CPL_1AH_2AG


Acknowledgements

We thank the CERCA Programme/Generalitat de Catalunya for their support.

Competing interests

JB Gornals is a consultant for Boston Scientific.

The authors

Sandra Maisterra^{1,2,3,4}, **Sergi Quintana-Carbo**^{1,2,3}, **Humberto Aranda**^{2,3,4,5}, **Mariona Calvo**^{2,4,6}, **Leandre Farran**^{2,3,4,5}, **Nuria Virgili**^{2,3,4,7}, **Joan B. Gornals**^{1,2,3,4} 

- 1 Endoscopy Unit, Department of Digestive Diseases, Hospital Universitari de Bellvitge, Barcelona, Catalonia, Spain
- 2 Bellvitge Biomedical Research Institute (IDIBELL), Barcelona, Catalonia, Spain
- 3 Universitat de Barcelona, Barcelona, Catalonia, Spain
- 4 Gastroesophageal Tumours Functional Unit (UTEG), Hospital Universitari de Bellvitge, Institut Català d'Oncologia, Barcelona, Spain
- 5 General and Digestive Surgery Department, Hospital Universitari de Bellvitge, Barcelona, Catalonia, Spain
- 6 Medical Oncology Department, Institut Català d'Oncologia (ICO), Barcelona, Spain
- 7 Endocrinology and Clinical Nutrition department, Hospital Universitari de Bellvitge, Barcelona, Catalonia, Spain

Corresponding author

Joan B. Gornals, MD, PhD

Endoscopy Unit, Department of Digestive Diseases, Hospital Universitari de Bellvitge, Bellvitge Biomedical Research Institute (IDIBELL), Universitat de Barcelona, Feixa Llarga s/n, 08907 L'Hospitalet de Llobregat, Barcelona, Catalonia, Spain
Fax: +34-93-2607681
jgornals@bellvitgehospital.cat

References

- [1] Lafeuille P, Wallenhorst T, Lupu A et al. Endoscopic submucosal dissection combined with clips for closure of gastrointestinal fistulas including those refractory to previous therapy. *Endoscopy* 2022; 54: 700–705
- [2] Beoletto F, Rostain F, Lesne A et al. Refractory esophagopleural fistula post-lobectomy successfully treated by endoscopic submucosal dissection and over-the-scope clip. *Endoscopy* 2021; 53: E138–E139
- [3] Ciochina M, Cruceru M, Rivory J et al. Esophagobronchial fistula after sleeve gastrectomy successfully treated by endoscopic submucosal dissection and over-the-scope clip. *Endoscopy* 2020; 52: E100–E101
- [4] Ryska O, Martinek J, Filipkova T et al. Single loop-and-clips technique (King closure) for gastrectomy closure after transgastric ovariectomy: a survival experiment. *Wideochir Inne Tech Maloinwazyjne* 2012; 7: 233–239
- [5] Dolezel R, Ryska O, Kollar M et al. A comparison of two endoscopic closures: over-the-scope clip (OTSC) versus KING closure (endoloop + clips) in a randomized long-term experimental study. *Surg Endosc* 2016; 30: 4910–4916

Bibliography

Endoscopy 2023; 55: E944–E945

DOI 10.1055/a-2127-4663

ISSN 0013-726X

© 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited.
(<https://creativecommons.org/licenses/by/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



E-Videos is an open access online section of the journal *Endoscopy*, reporting on interesting cases

and new techniques in gastroenterological endoscopy. All papers include a high-quality video and are published with a Creative Commons CC-BY license. *Endoscopy* E-Videos qualify for HINARI discounts and waivers and eligibility is automatically checked during the submission process. We grant 100% waivers to articles whose corresponding authors are based in Group A countries and 50% waivers to those who are based in Group B countries as classified by Research4Life (see: <https://www.research4life.org/access/eligibility/>).

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>

CORRECTION**Correction: Successful closure of a refractory gastrobronchial fistula using endoscopic mucosal ablation followed by single loop-and-clips technique**

Maisterra S, Quintana-Carbo S, Aranda H et al. Successful closure of a refractory gastrobronchial fistula using endoscopic mucosal ablation followed by single loop-and-clips technique.

Endoscopy 2023, 55: E944–E945.

In the above-mentioned article, the video has been replaced.

This was corrected in the online version on December 19, 2024.