



“Pure Fat Flap”—Perforator-based Adiposal Layer Only Flap for Lateral Ankle Reconstruction

Seok Joon Lee, MD¹ Jeongmok Cho, MD, PhD¹ Changsik Pak, MD, PhD¹
Hyunsuk Suh, MD, PhD¹ Joon Pio Hong, MD, PhD, MMM¹

¹ Department of Plastic and Reconstructive Surgery, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Republic of Korea

Address for correspondence Joon Pio Hong, MD, PhD, MMM, Department of Plastic Surgery, Asan Medical Center, University of Ulsan, School of Medicine, 88, Olympic-ro 43-gil, Songpa-gu, Seoul 138-736, Republic of Korea (e-mail: joonphong@amc.seoul.kr).

Arch Plast Surg 2025;52:41–45.

Abstract

Lateral ankle soft tissue defects pose challenges, especially in cases due to chronic pressure from cross-legged sitting, which usually present with a large dead space, small skin opening that often accompanies an open joint. Traditional reconstruction methods using fasciocutaneous flaps may result in donor site morbidity such as delayed wound healing or nerve injury. In this article, we present a case of diabetes-related lateral ankle defect successfully treated using adiposal layer only flap, also known as pure fat flap. The anatomy and the surgical technique of adiposal layer only flap were reviewed. These flaps preserve the subdermal plexus and deep fascia while obliterating dead space and providing a gliding surface for proper ankle movement. A perforator-based adiposal layer only flap was elevated from the peroneal artery and used to cover the defect. Flap perfusion was confirmed using indocyanine green video angiography and color duplex ultrasound. Patient had a successful recovery with minimal donor site morbidity. The technique expands the reconstructive microsurgeon's options for complex ankle coverage, ensuring optimal wound healing and functional outcomes.

Keywords

- ▶ ankle
- ▶ perforator flap
- ▶ wound healing

Introduction

Trauma, tumor, diabetes mellitus, and peripheral vascular disease are responsible for majority of cases that present with soft tissue defect of lateral ankle.^{1,2} Defects related to diabetes mellitus especially presents a significant challenge in terms of reconstruction and healing.³ In culture where sitting cross-legged on the floor is common, the lateral ankle wound becomes further difficult as they are usually presented with a large dead space, small skin opening that often accompanies an open joint.

The first line for reconstruction can be a simple rotation of a local flap or a propeller flap based on peroneal artery perforator.⁴ These fasciocutaneous flaps are commonly

utilized and have provided several advantages in lateral ankle reconstruction including reduced operative time and improved wound healing. However, the use of fasciocutaneous local flaps or propeller flaps, in which a skin paddle is harvested from a tight donor site often can lead to morbidity such as delayed wound healing or nerve injury.

To address these issues for diabetes-related lateral ankle defects where the dead space is far greater issue than the skin defect, the authors aimed for a flap that can obliterate the dead space while allowing smooth gliding for underneath structures. Theoretically, obliterating the dead space with fat can be anatomically ideal. Thus, the adiposal layer only flap or so called the pure fat flap based on a single perforator was designed. The pure fat flap will preserve the subdermal

received

September 4, 2023

accepted after revision

January 30, 2024

accepted manuscript online

February 13, 2024

article published online

April 10, 2024

DOI <https://doi.org/10.1055/a-2267-4205>

10.1055/a-2267-4205.

eISSN 2234-6171.

© 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)

Thieme Medical Publishers, Inc., 333 Seventh Avenue, 18th Floor, New York, NY 10001, USA

plexus of the skin allowing the donor site skin to heal without the need for any skin grafts, preserve the deep fascia reducing the chance of muscle hernia or nerve injury. To our knowledge, this is the first case of utilizing a pure fat flap of moderate size which was performed to obliterate the dead space and provide coverage for the lateral ankle defect.

Case

A 46-year-old male patient was noted with bursitis of the left lateral malleolus. The orthopaedic team performed multiple incisions and drainage without success and ultimately referred to reconstruction for the defect. Debridement was performed and resulted in a 2.5×2 cm skin defect and a 5×5 cm dead space pocket with small opening of the joint leaking fluid (►Fig. 1A). A contrast-enhanced magnetic resonance imaging of the left ankle was performed, revealing no evidence of osteomyelitis. The patient had history of coronary intervention for 3-vessel disease, end-stage renal

disease on hemodialysis, and had diabetes for 20 years. The ankle brachial index was 1.35 on the left side (1.28 on the right side), and enhanced computed tomography lower extremity angiography revealed patency of the anterior tibial artery, posterior tibial artery, and peroneal artery, which was further confirmed by ultrasound examination. Prior to coverage, duplex ultrasonography was used preoperatively to find a perforator that would allow a perforator-based adiposal layer turnover flap (►Fig. 1B).⁵ A perforator was identified from the peroneal artery.

A 9×4 cm turnover pure fat flap was designed based on the perforator located 2 cm from the edge of the defect (►Fig. 1C). Dissection was performed above the deep fascia, perforator was identified, and adiposal layer only flap was elevated preserving subdermal plexus. The sural nerve was also saved. The perforator based pure fat flap was elevated and turned over to cover lateral malleolus. The indocyanine green angiography confirmed the flap perfusion. Primary closure with quilting suture was performed at the donor site



Fig. 1 (A) (Left) Preoperative photo of patient 1. (Right) Postoperative photo of patient 1 after debridement. (B) Ultrasonographic finding of patient 1, showing peroneal artery (shown by arrowheads) and accompanying perforator from peroneal artery (shown by arrow). It was traced before surgery using color duplex ultrasound. (C) Adiposal layer only flap design of patient 1 according to perforator course. (D) (Above, Left) After adiposal layer only flap elevation. (Above, Right) After flap rotation. (Below, Left) ICG video angiography after flap rotation. Note that flap is well perfused including distal portion after rotation. (Below, Right) Immediate postoperative photo. (E) Postoperative ultrasonographic finding showing intact pedicle. Peroneal artery perforator is seen (shown by arrow) and continued in axial pattern (shown by arrowheads). (F) Postoperative 8 months photo of patient 1. ICG, indocyanine green.

to eliminate dead space. The pure fat flap obliterated and covered the defect, and the skin was closed primarily. A silastic drain was placed at the lateral ankle with a splint to immobilize the joint (►Fig. 1D).

Postoperative process was uneventful, and the drain was removed on day 4. The follow-up duplex ultrasound on day 5 showed good flow (►Fig. 1E). The follow-up at 8 months showed good coverage and healing with minimal donor site morbidity (►Fig. 1F).

Discussion

For diabetes-related lateral ankle defects with specific features such as dead space and a relatively small skin defect, several factors need to be considered. The presence of dead space in the lateral ankle defect refers to the gap or void that may exist above the lateral malleolus, which is easily seen during debridement for wound preparation. Dead space can pose a challenge to wound healing and increase the risk of complications such as fluid accumulation and infection. It is, therefore, critical to effectively manage this dead space to promote optimal healing and functional outcomes. The dynamic nature of the ankle also requires the provision of a gliding surface within the defect reconstruction.

In addressing these specific requirements, surgeons may employ various approaches. Since lateral ankle is a watershed area between two angiosomes, it has a relatively poor blood supply compared with other areas of the body leading to potential poor secondary healing.⁶ Patients with diabetes mellitus or peripheral vascular disease are prone to malleolus defect to occur and they may not be a good candidate for free tissue transfer.⁷ For local flap coverage, such as reverse sural artery flap,⁸ it is common for distal foot wounds to become

larger due to donor site morbidity. Adipofascial flap, composed of perforator, deep fascia, and fat layer spares subdermal plexus and skin for donor and this makes donor site healing more effective. They were previously employed in head and neck reconstruction⁹ and lower extremity reconstruction,⁴ utilizing various methods such as the adipofascial fold-down flap¹⁰ or the venoadipofascial pedicled fasciocutaneous flap.¹¹ However, with deep fascia harvested together, it has possibility of muscle hernia and injuring the nerve leading to persistent pain, altered sensation, or numbness in the donor site. The pure fat flap, with proper dissection, can preserve deep fascia which poses no risk for sural nerve damage.

Perforator-based propeller flaps have become a valuable tool in reconstructive surgery,¹² allowing for the transfer of tissue from a donor site to a recipient site while preserving major blood vessels. These flaps are designed based on the concept of utilizing specific perforating blood vessels that supply the overlying tissue. With duplex ultrasound, a noninvasive imaging technique, we can identify the pedicle of interest and design and elevate flap according to pedicle course looking at actual vessels. However, we commonly encounter fat necrosis without skin necrosis after proper flap elevation according to perforator^{13,14} and thus circulation to subcutaneous fat (adiposal layer) could be of question.

Blood circulation in the skin is indeed facilitated by capillaries, which are tiny blood vessels that form a dense network throughout the dermis, the middle layer of the skin. As for fat (adipose cell), it does not have its own circulatory system like capillaries.^{15,16} Instead, the blood supply to adipose tissue comes from the surrounding capillaries. In other words, adipose tissue does not have terminal branches of blood vessels like capillaries but relies on the adjacent capillaries to maintain its metabolic functions (►Fig. 2).




Fig. 2 Schematic illustration of fat circulation. At subcutaneous fat level, there are no terminal capillary and cells are supplied by adjacent capillaries.

The capillaries in the skin, which are part of the overall circulatory system, serve not only the skin itself but also support the functions of underlying fat tissue. Therefore, we suggest that adiposal layer only flap should be designed more conservatively in smaller dimension than we would normally design a perforator-based propeller flap.

Gold standard of checking blood flow of adipose tissue is ¹³³Xenon washout technique,¹⁷ however it is hard to use it in clinical setting. Alternative techniques such as Doppler ultrasound was studied, and recent study¹⁸ comparing ¹³³Xenon washout technique and Doppler ultrasound by Lempesis et al showed ultrasound can be effectively used as alternative to check and quantify adipose tissue blood flow. Thus the circulation of adiposal layer only flap was checked with color duplex ultrasound both preoperatively and postoperatively, showing intact pedicle. Moreover, intraoperative indocyanine fluorescent angiography helps evaluate the perfusion of flaps,¹⁹ and all flap perfusion was checked with indocyanine fluorescent angiography after elevation and rotation of the flap to the lateral ankle defect. Postoperative care is also important because stretching of flap can result in damage in pedicle such as thrombosis or ischemia. Ankle immobilization in early postoperative period will be important, and we used short leg cast for protection.

There have been previous reports of adiposal flap in finger reconstruction, which used pedicled adipose tissues based on digital artery.²⁰ They identified digital artery and elevated preserving pedicle, but they did not check the perfusion afterwards. With aid of indocyanine fluorescent angiography and color duplex ultrasound, we ensured the perfusion of flap. Also, digital artery-based adiposal flaps were much smaller compared with our flap being long as 7 to 9 cm.

Limitation of adiposal layer only flap exists. Further understanding of the vascularity for the adiposal layer is needed. Nevertheless, this report demonstrates the possibility and potential of using the fat only as a flap.

Conclusion

Pure fat flap or the adiposal layer only flap may be an alternative for reconstruction in areas with large dead space, small skin defect, and joint movement. The advantages of the pure fat flap are preserving the donor site skin, allow vascular bulk tissue to obliterate dead space, and the fat to preserve good gliding function of the underlying structures. By carefully addressing the specific requirements of the defect and utilizing advanced imaging techniques, we can enhance the chances of optimal wound healing using this approach. This technique adds to the reconstructive microsurgeon's armamentarium for complex coverage of the ankle region.

Authors' Contributions

Conceptualization: S.J.L., J.M.C., C.S.P., H.S.S., J.P.H. Data curation: S.J.L. Formal analysis: S.J.L. Methodology: J.P.H. Project administration: J.P.H. Visualization: S.J.L. Writing - original draft: S.J.L. Writing - review & editing: J.M.C., J.P.H. All authors read and approved the final manuscript.

Patient Consent

Written informed consent was obtained from the patient.

Funding

None.

Conflict of Interest

H.S. and J.P.H. are editorial board members of the journal but were not involved in the peer reviewer selection, evaluation, or decision process of this article. No other potential conflicts of interest relevant to this article were reported.

References

- Bhandari PS, Bath AS, Sadhotra LP, Singh M, Mukherjee MK. Management of soft tissue defects of the ankle and foot. *Med J Armed Forces India* 2005;61(03):253–255
- Chen H, Yin G, Hou C, Zhao L, Lin H. Repair of a lateral malleolus defect with a composite pedicled second metatarsal flap. *J Int Med Res* 2018;46(12):5291–5296
- Schaper NC, van Netten JJ, Apelqvist J, et al; IWGDF Editorial Board. Practical guidelines on the prevention and management of diabetes-related foot disease (IWGDF 2023 update). *Diabetes Metab Res Rev* 2023:e3657
- Suliman MT. Distally based adipofascial flaps for dorsal foot and ankle soft tissue defects. *J Foot Ankle Surg* 2007;46(06):464–469
- Cho MJ, Kwon JG, Pak CJ, Suh HP, Hong JP. The role of duplex ultrasound in microsurgical reconstruction: review and technical considerations. *J Reconstr Microsurg* 2020;36(07):514–521
- Attinger CE, Evans KK, Bulan E, Blume P, Cooper P. Angiosomes of the foot and ankle and clinical implications for limb salvage: reconstruction, incisions, and revascularization. *Plast Reconstr Surg* 2006;117(7 Suppl):261S–293S
- Lese I, Biedermann R, Constantinescu M, Grobelaar AO, Olariu R. Predicting risk factors that lead to free flap failure and vascular compromise: a single unit experience with 565 free tissue transfers. *J Plast Reconstr Aesthet Surg* 2021;74(03):512–522
- Daar DA, Abdou SA, David JA, Kirby DJ, Wilson SC, Saadeh PB. Revisiting the reverse sural artery flap in distal lower extremity reconstruction: a systematic review and risk analysis. *Ann Plast Surg* 2020;84(04):463–470
- Mohindra A, Parmar S, Praveen P, Martin T. The fat-fascia paddle only with a composite fibula flap: marked reduction in donor site morbidity. *Int J Oral Maxillofac Implants* 2016;45(08):964–968
- Lee KJ, Lee SH, Kim MB, Lee YH. Adipofascial fold-down flaps based on the posterior tibial artery perforator to cover the medial foot and ankle defects. *J Plast Reconstr Aesthet Surg* 2016;69(12):e229–e237
- Nakajima H, Imanishi N, Fukuzumi S, et al. Accompanying arteries of the lesser saphenous vein and sural nerve: anatomic study and its clinical applications. *Plast Reconstr Surg* 1999;103(01):104–120
- Bekara F, Herlin C, Mojallal A, et al. A systematic review and meta-analysis of perforator-pedicled propeller flaps in lower extremity defects: identification of risk factors for complications. *Plast Reconstr Surg* 2016;137(01):314–331
- Khansa I, Momoh AO, Patel PP, Nguyen JT, Miller MJ, Lee BT. Fat necrosis in autologous abdomen-based breast reconstruction: a systematic review. *Plast Reconstr Surg* 2013;131(03):443–452
- Kim HB, Han SJ, Kim EK, Eom JS, Han HH. Comparative study of DIEP and PAP flaps in breast reconstruction: reconstructive outcomes and fat necrosis. *J Reconstr Microsurg* 2023;39(08):627–632
- Bouloumié A, Galitzky J. Angiogenesis in adipose tissue. In: Bastard J-P, Fève B, eds. *Physiology and Pathophysiology of Adipose Tissue*. Paris: Springer; 2013:27–38
- Corvera S, Solivan-Rivera J, Yang Loureiro Z. Angiogenesis in adipose tissue and obesity. *Angiogenesis* 2022;25(04):439–453

- 17 Larsen OA, Lassen NA, Quaade F. Blood flow through human adipose tissue determined with radioactive xenon. *Acta Physiol Scand* 1966;66(03):337–345
- 18 Lempesis IG, Goossens GH, Manolopoulos KN. Measurement of human abdominal and femoral intravascular adipose tissue blood flow using percutaneous Doppler ultrasound. *Adipocyte* 2021;10(01):119–123
- 19 Li K, Zhang Z, Nicoli F, et al. Application of indocyanine green in flap surgery: a systematic review. *J Reconstr Microsurg* 2018;34(02):77–86
- 20 Onode E, Takamatsu K, Kazuki K, Nakamura H. Bipedicled digital artery perforator volar adiposal turned-over flap for reconstruction of transverse fingertip amputation. *JPRAS Open* 2022;32:111–115