L. LI, E. G. KLAUBER, D. SEIDEL* (RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY, PISCATAWAY, USA)

Catalytic Enantioselective Aldol Additions of α -Isothiocyanato Imides to Aldehydes

J. Am. Chem. Soc. **2008**, *130*, 12248-12249.

Gold-Free Aldol-Type Additions of α -Isothiocyanato Imides to Aldehydes

Large-scale synthesis:

Significance: Seidel and co-workers report an organocatalytic diastereo- and enantioselective aldol addition of α -isothiocyanato imides to aldehydes. The reaction is catalyzed by bifunctional thiourea catalyst **1**. A sterically demanding substitution on the imide is necessary for high diastereoselectivities and moreover for suppressing the formation of undesired product **2**. The applicability was shown by preparing the protected β -hydroxy- α -amino acid **3** on a large scale and reducing the catalyst loading to 1 mol%.

Comment: A direct organocatalytic and highly diastereo- and enantioselective approach to protected *syn*- β -hydroxy- α -amino acids under mild conditions with low catalyst loadings is presented. The approach complements one of the first catalytic asymmetric aldol reactions by Ito and Hayashi (*J. Am. Chem. Soc.* **1986**, *108*, 6405). So far, the majority of methods available to synthesize *syn*- β -hydroxy- α -amino acids rely on diastereoselective approaches, for example the use of chiral auxiliaries. A catalytic enantioselective approach to these compounds remains challenging. An expansion of this method to different imides would be desirable.