M. VOLGRAF, J.-P. LUMB, H. C. BRASTIANOS, G. CARR, M. K. W. CHUNG, M. MÜNDEL,
A. G. MAUK, R. J. ANDERSEN,* D. TRAUNER* (UNIVERSITY OF CALIFORNIA, BERKELEY,
USA; UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, CANADA; LUDWIG-
MAXIMILIANS-UNIVERSITÄT MÜNCHEN, GERMANY)

Biomimetic Synthesis of the IDO Inhibitors Exiguamine A and B

Nat. Chem. Biol. 2008, 4, 535-537.

Synthesis of Exiguamine A

Significance: Exiguamine A is isolated from the marine sponge *Neopetrosia exigua* and is a potent inhibitor of indoleamine-2,3-dioxygenase, which in turn metabolizes tryptophan. This biomimetic synthesis relies on a cascade sequence involving intramolecular nucleophilic attack of a tertiary amine onto an *o*-quinone and tautomerization followed by an oxa-6 π -electrocyclization (**I** $\rightarrow$ exiguamine A).

Comment: The phenol resulting from benzyl deprotection of **F** was subjected to salcomine oxidation which gave a 2.5:1 mixture of the regioisomeric *o*- and *p*-quinones, respectively (**F** $\rightarrow$ **G**). Upon treatment with 10 equivalents of AgO, **I** afforded exiguamine A, but with 20 equivalents of AgO exiguamine B was the main product. Treatment of exiguamine A with excess of AgO failed to give exiguamine B.

SYNFACTS Contributors: Philip Kocienski, Indu Dager
Synfacts 2009, 3, 0233-0233 Published online: 19.02.2009
DOI: 10.1055/s-0028-1087689; **Reg-No.:** K00409SF