Endoscopic ultrasound-guided fine-needle aspiration using 22- and 25-gauge needles alternately

During endoscopic ultrasound (EUS)-guided fine-needle aspiration (FNA), the standard size of needle used is a 22-gauge needle. Larger needles have been used to obtain actual core tissue samples [1–3], but their has failed to significantly improve diagnostic accuracy for malignancy [2–4], except perhaps in the case of unusual histology [5]. On the contrary, a new, smaller-caliber (25-gauge) needle has been introduced to the market by Wilson-Cook Medical Inc. (Winston-Salem, North Carolina, USA). The purpose of this study was to compare the 22- and 25-gauge needles for adequacy of tissue acquisition and diagnostic yield.

The study was a retrospective review of all EUS-FNA procedures performed using 22- and 25-gauge needles alternately in the same patient. Of a total of 132 patients undergoing EUS, only 16 met the inclusion criteria. The mean age was 65.1 years. The cytotechnician was present during 75% of the procedures. The needle pass was considered by the endoscopist to be difficult in 37.5% vs. 25.0% of cases using the 22- and 25-gauge needles, respectively ($P=0.7$). The specimen adequacy rates were: cytologic 68.6 vs. 56.3 ($P=0.7$), and histologic 87.5% vs. 75.0% ($P=0.6$) with 22- and 25-gauge needles, respectively. Two patients were lost to follow-up. Out of the remaining 14 patients, a definitive diagnosis was obtained in 85.7% (22-gauge needle) and 50.0% (25-gauge needle) ($P=0.1$). When 22- and 25-gauge needles were combined,

the cytologic and histologic yields, as well as the definitive diagnosis, were higher (81.3%, 93.8%, and 92.9%, respectively). Hence, in conclusion we found no statistically significant difference between needle size despite a relatively easier pass with the 25-gauge needle and higher specimen adequacy and definitive diagnosis with the 22-gauge needle. Although we found the two needles to complement each other when used alternately in the same patient, the differences did not reach statistical significance due to the small number of cases. We recommend large prospective trials.

Endoscopy_UCTN_Code_TTT_1AS_2AF

S. Iqbal^{1,2}, R. N. Mir³, W. Sohn¹

¹ Division of Gastroenterology, New York Methodist Hospital, Brooklyn, New York, USA

² Division of Gastroenterology, Interfaith Medical Center, Brooklyn, New York, USA

³ Department of Pathology, New York Methodist Hospital, Brooklyn, New York, USA

References

- 1 Matsui M, Goto H, Niwa Y et al. Preliminary results of fine needle aspiration biopsy histology in upper gastrointestinal submucosal tumors. *Endoscopy* 1998; 30: 750–755
- 2 Binmoeller KF, Thul R, Rathod V et al. Endoscopic ultrasound-guided 18-gauge, fine needle aspiration biopsy of the pancreas using a 2.8 mm channel convex array echoendoscope. *Gastrointest Endosc* 1998; 47: 121–127
- 3 Levy MJ, Jondal ML, Clain J, Wiersema MJ. Preliminary experience with an EUS-guided trucut biopsy needle compared with EUS-guided FNA. *Gastrointest Endosc* 2003; 57: 101–106
- 4 Solmi L, Muratori R, Bacchini P et al. Comparison between echo-guided fine needle aspiration cytology and microhistology in diagnosing pancreatic masses. *Surg Endosc* 1992; 6: 222–224
- 5 Binmoeller KF, Rathod VD. Difficult pancreatic mass FNA: tips for success. *Gastrointest Endosc* 2002; 56: S86–S93

Bibliography

DOI 10.1055/s-0028-1103444

Endoscopy 2009; 41: E87

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

S. Iqbal, MD

New York Methodist Hospital,
Division of Gastroenterology
753 Classon Ave, 7A
Brooklyn
NY 11238
USA
Fax: +1-718-780-3851
shahzad_iqbal@hotmail.com

Table 1 Comparison of 22- and 25-gauge needles with combined technique.

	22-gauge needle		25-gauge needle		Combined		P-value
	n/N	%	n/N	%	n/N	%	
Tissue yield	16/16	100	15/16	93.6	16/16	100	NS
Difficult needle pass	6/16	37.5	4/16	25.0	N/A		NS
Cytologic adequacy	11/16	68.6	9/16	56.3	13/16	81.3	NS
Histologic adequacy	14/16	87.5	12/16	75.0	15/16	93.8	NS
Definitive diagnosis	12/14	85.7	7/14	50.0	13/14	92.9	NS

N/A, not applicable; NS, not significant.