“Short” double balloon enteroscope for endoscopic retrograde cholangiopancreatography with conventional sphincterotomy and metallic stent placement after Billroth II gastrectomy

Fig. 1 Although the papilla is located at the 11 o'clock position in the visual field (left), the papilla can be moved to the 6 o'clock position by keeping the overtube balloon inflated and rotating the enteroscope, resulting in successful cannulation (right).

Fig. 2 After deep cannulation, a stricture is visualized in the common bile duct.

Fig. 3 After endoscopic sphincterotomy with a conventional sphincterotome, a metallic stent is easily inserted across the stricture.

During endoscopic retrograde cholangiopancreatography (ERCP) in patients with Billroth II gastrectomy, identifying the afferent loop, access to the papilla, and cannulation in the inverted papilla are difficult [1]. With the advent of the double balloon technique, ERCP in such patients can be performed [1], but conventional accessories cannot be used because of short accessory lengths [1–5]. We carried out double balloon enteroscopy for ERCP (DB-ERCP) with a “short” enteroscope in a patient with Billroth II gastrectomy,

and successfully performed ERCP with conventional sphincterotomy and metallic stent placement.

A 79-year-old man, who had undergone Billroth II gastrectomy for gastric cancer, presented with jaundice. Computed tomography (CT) showed a tumor in the common bile duct causing biliary tree dilatation. We performed DB-ERCP with a “short” enteroscope, EC450-BI5 (Fujinon-Toshiba, Osaka, Japan), which has a 2.8-mm working channel and a 152-cm working length and for which conventional ac-

cessories are available. The enteroscope successfully reached the duodenal stump. Despite the inverted papilla being located at the 11 o'clock position in the visual field, the papilla could be moved to the 6 o'clock position by keeping the overtube balloon inflated and rotating the enteroscope, resulting in successful cannulation (● Fig. 1). The diagnosis of malignant biliary structure caused by recurrent gastric cancer was made based on imaging and histologic studies (● Fig. 2). After endoscopic sphincterotomy with a conventional sphincterotome, a metallic stent was easily inserted across the stricture with uneventful recovery (● Fig. 3).

ERCP is technically challenging and often unsuccessful in patients with altered gastrointestinal anatomy [1,3–5]. Several authors have performed DB-ERCP in such patients, with the use of a standard EN-450P5/20 enteroscope or a therapeutic EN-450T5 enteroscope [1–5]. Because both enteroscopes have a working length of 200 cm, specialized custom-made or very limited accessories are available [1–5]. We therefore recommend the “short” enteroscope for diagnostic and therapeutic ERCP in patients with altered gastrointestinal anatomy because all conventional accessories are available.

Endoscopy_UCTN_Code_TTT_1AR_2AK
Endoscopy_UCTN_Code_TTT_1AR_2AZ

**M. Shimatani, M. Matsushita,
M. Takaoka, T. Kusuda, N. Fukata,
M. Koyabu, K. Uchida, K. Okazaki**
Third Department of Internal Medicine,
Kansai Medical University, Osaka, Japan

References

- 1 *Chu YC, Su SJ, Yang CC et al.* ERCP plus papil-
lotomy by use of double-balloon enterosco-
py after Billroth II gastrectomy. *Gastrointest
Endosc* 2007; 66: 1234–1236
- 2 *Maaser C, Lenze F, Bokemeyer M et al.* Double
balloon enteroscopy: a useful tool for diag-
nostic and therapeutic procedures in the
pancreaticobiliary system. *Am J Gastroen-
terol* 2008; 103: 894–900
- 3 *Mönkemüller K, Bellutti M, Neumann H et al.*
Therapeutic ERCP with the double-balloon
enteroscope in patients with Roux-en-Y
anastomosis. *Gastrointest Endosc* 2008;
67: 992–996
- 4 *Aabakken L, Bretthauer M, Line PD.* Double-
balloon enteroscopy for endoscopic retro-
grade cholangiography in patients with a
Roux-en-Y anastomosis. *Endoscopy* 2007;
39: 1068–1071
- 5 *Emmett DS, Mallat DB.* Double-balloon ERCP
in patients who have undergone Roux-en-Y
surgery: a case series. *Gastrointest Endosc*
2007; 66: 1038–1041

Bibliography

DOI 10.1055/s-0028-1103461
Endoscopy 2009; 41: E19–E20
© Georg Thieme Verlag KG Stuttgart · New York ·
ISSN 0013-726X

Corresponding author

M. Matsushita, MD
Third Department of Internal Medicine
Kansai Medical University
2-3-1 Shinmachi
Hirakata
Osaka 573-1191
Japan
Fax: +81-72-8042061
matsumit@hirakata.kmu.ac.jp