Ultrasound Criteria for Diagnosis of Early Pregnancy Failure and Ectopic Pregnancy

Lisa Perriera, M.D.,^{1,3} and Matthew F. Reeves, M.D., M.P.H.^{2,3}

After publication of the above mentioned article in *Seminars in Reproductive Medicine*, Volume 26, Number 5, 2008, pages 372–382, the authors noticed an error in the abstract. The fourth sentence of the abstract reads “An early pregnancy failure can be diagnosed by the absence of a visible yolk sac with a mean sac diameter of 13 mm; the absence of a visible embryo with a mean sac diameter of 13 mm; the absence of cardiac motion with an embryo measuring 5 mm or more in maximal length; or the presence of an empty amnion.”

The sentence should read “An early pregnancy failure can be diagnosed by the absence of a visible yolk sac with a mean sac diameter of 13 mm; the absence of a visible embryo with a mean sac diameter of **20 mm**; the absence of cardiac motion with an embryo measuring 5 mm or more in maximal length; or the presence of an empty amnion.”

¹Clinical Instructor, ²Assistant Professor, ³Department of Obstetrics, Gynecology, and Reproductive Sciences, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania.

Address for correspondence and reprint requests: Matthew F. Reeves, M.D., M.P.H., Department of Obstetrics, Gynecology, and Reproductive Sciences, 300 Halket Street, Pittsburgh, PA 15213-3180 (e-mail: mreeves@upmc.edu).

Growth Factors and the Development and Function of the Reproductive Organs; Guest Editors, Linda C. Giudice, M.D., Ph.D., and Marco Conti, M.D.

Semin Reprod Med 2009;27:103–104. Copyright © 2009 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA. Tel: +1(212) 584-4662.
DOI 10.1055/s-0028-1121901. ISSN 1526-8004.