Capillary Electrophoresis of Natural Products: Current Applications and Recent Advances

Author

Matthias Unger

Affiliation

Institute of Pharmacy and Food Chemistry, Department of Pharmaceutical Chemistry, University of Würzburg, Würzburg, Germany

Key words

- capillary electrophoresis
- capillary electrophoresis-mass spectrometry
- capillary zone electrophoresis
- field-amplified sample injection
- micellar electrokinetic chromatography
- microemulsion electrokinetic chromatography
- sweeping

received Sept. 29, 2008
revised Dec. 23, 2008
accepted February 2, 2009

Bibliography

DOI 10.1055/s-0029-1185453
Published online March 16, 2009
Planta Med 2009; 75: 735–745
© Georg Thieme Verlag KG
Stuttgart · New York ·
ISSN 0032-0943

Correspondence

Matthias Unger
Department of Pharmaceutical Chemistry
Institute of Pharmacy and Food Chemistry
University of Würzburg
Am Hubland
97074 Würzburg
Germany
Phone: + 49 93 18 88 54 63
Fax: + 49 93 18 88 54 94
m.unger@pharmazie.uni-wuerzburg.de

Abstract

Capillary electrophoresis (CE) was introduced as a new analytical technique in the 1970s and rapidly proved to be a powerful tool for the separation and detection of various classes of natural and synthetic compounds. Since the availability of commercially manufactured high-performance instruments, CE represents an interesting alternative to high-pressure liquid chromatography (HPLC), mainly because of its speed and high separation efficiency. In this overview a short description of the basic and widely used CE methods will be given and the applicability of these methods for the analysis of natural products will be discussed. Due to the growing number of publications dealing with CE or CE/MS of secondary plant metabolites, an exhaustive overview of all current applications cannot be given in this contribution. Therefore, the focus of this mini-review will be on the advances and new aspects of recently published CE methods in natural products analysis.

Introduction

Capillary electrophoresis (CE) represents one of the most attractive analytical techniques for the rapid qualitative and quantitative analysis of molecules with a wide range of polarity and molecular weight, including small molecules such as drugs but also macromolecules such as proteins or nucleic acids. Because of its versatility and high separation efficiency, CE is an interesting alternative to the widely used reversed-phase high-pressure liquid chromatography (RP-HPLC) and has gained much interest for the analysis of natural products in herbal extracts, pharmaceutical for-

Abbreviations

ACN:	acetonitrile
AcOH:	acetic acid
CE:	capillary electrophoresis
CD:	cyclodextrin
CEC:	capillary electrochromatography
CE/ESI-MS:	capillary electrophoresis-electrospray ionisation mass spectrometry
CZE:	capillary zone electrophoresis
DMF:	dimethyl formamide
EOF:	electroosmotic flow
ESI:	electrospray ionisation
FASI:	field-amplified sample injection
FESS:	field-enhanced sample stacking
IsoOH:	isopropanol
MEKC:	micellar electrokinetic chromatography
MEEKC:	microemulsion electrokinetic chromatography
MS:	mass spectrometry
NACE:	non-aqueous capillary electrophoresis
MeOH:	methanol
RM-MEKC:	reversed-migration MEKC
THF:	tetrahydrofuran

mulations, and food supplements. Since its introduction in 1979 by Mikkers et al. [1] and in 1981 by Jorgenson and Lukacs [2, 3], the popularity of CE continuously increased so that high-performance CE instruments became rapidly available. Over the years the tremendous progress of this fascinating separation technique has led to a great variety of separation methods and applications that have been outlined in recently published reviews [4–9]. CE has been frequently applied especially for the qualitative and quantitative analysis of secondary plant metabolites in crude plant extracts [10–12]. In the following, a short description of the basic and widely used CE methods

(e.g., capillary zone electrophoresis [CZE], non-aqueous capillary electrophoresis [NACE], micellar electrokinetic chromatography [MEKC], microemulsion electrokinetic chromatography [MEEKC], and capillary electrochromatography [CEC]) will be given, and the applicability of these methods for the analysis of secondary plant metabolites will be discussed.

Capillary Zone Electrophoresis

In the absence of an electroosmotic flow (EOF), the migration velocity v_i of a charged molecule i in an electrical field is proportional to the electrophoretic mobility μ_i and the electrical field strength E (equation 1):

$$v_i = \mu_i \cdot E$$

The electrophoretic mobility μ_i is dependent on the net charge $z_i \cdot e_0$ and the friction coefficient $6 \cdot \pi \cdot r \cdot \eta$ (equation 2):

$$\mu_i = \frac{z_i \cdot e_0}{6 \cdot \pi \cdot r \cdot \eta}$$

Whereas the net charge of a molecule is the result of its pK_a value and of the pH value of the electrolyte, the friction coefficient is determined by its size ($6 \cdot \pi \cdot r$) and the viscosity (η) of the running buffer. As the size and shape of a molecule cannot be influenced, the net charge of the analytes can be manipulated by the pH value of the buffer. Furthermore, if the pK_a values of the analytes are known, the net charge can be calculated via the dissociation rate using the Henderson-Hasselbalch equation (equation 3):

$$pH = pK_a + \log \frac{c(\text{base})}{c(\text{acid})}$$

Ideal candidates for CZE analyses of natural products are permanently charged molecules such as anthocyanins, quaternary alkaloids, and sulphated flavonoids. For such molecules, the pH value of the buffer can be varied over a wide range without loss of the electrophoretic mobility, which is also particularly advantageous for capillary electrophoresis-mass spectrometry (CE/MS). Indeed, CE/MS analyses of anthocyanins and quaternary alkaloids have been published frequently [13–17]. If several permanently charged molecules with similar molecular weights have to be analysed, the permanent charge is a disadvantage because all molecules exhibit a very similar mass-to-charge ratio. In this case the selectivity can be tuned by organic solvents and counterions that have a high affinity to the oppositely charged analytes [18].

Because many secondary plant metabolites (e.g., benzoic or cinnamic acids, coumarins, flavonoids) possess phenolic hydroxy groups, they can be analysed with CZE using running buffers with neutral to basic pH values in the range between 7 and 12. For example, the three major curcuminoids curcumin, demethoxycurcumin, and bis-demethoxycurcumin from *Curcuma domestica*, *Curcuma longa*, and *Curcuma xanthorrhiza* were fully separated and quantified in less than 5 min using a basic running buffer consisting of 20 mM phosphate, 50 mM sodium hydroxide, and 14 mM β -cyclodextrin (β -CD) [19]. Comparable to CEC, MEKC and MEEKC, CE analyses with electrolytes containing cyclodextrins can also be considered electrokinetic chromatography. The addition of β -CD enhanced the solubility of the curcuminoids and influenced the selectivity because the different stability of the curcuminoid-CD inclusion complexes individually influenced the migration behaviour of the analytes. Curcuminoids with a

high affinity to the β -CD molecules have a lower migration velocity to the anode than do curcuminoids that do not easily form inclusion complexes with β -CD. For some natural product classes that are not acidic or basic (e.g., sugars, phenolic glycosides), a charge can be created by complexation. Vicinal hydroxy groups of neutral sugars or glycosides with a *cis*-configuration can form complexes with borate anions, which leads to a migration of the anionic complexes to the anode. In fused silica capillaries and at neutral to basic pH values, the EOF carries the negatively charged complexes to the cathode. In this case the selectivity of the separation is determined by the stability, size, and molecular geometry of the analyte-borate complexes. This separation principle was frequently applied for the analysis of neutral secondary plant metabolites in the past [20,21]. For example, Honda et al. reported the CZE analysis of reducing monosaccharides after derivatisation to their corresponding *N*-2-pyridylglycamines [20]. The derivatised sugar molecules were separated as their borate complexes using 200 mM borate (pH 10.5) as running buffer. Also, Schwaiger et al. separated aldoses and ketoses as their borate complexes using 175 mM borate (pH 10.5) as electrolyte [21]. In order to enhance the UV absorption of the sugars, a reductive amination with 4-aminobenzonitrile was performed.

Among the different classes of natural compounds analysed with CZE are alkaloids [16,22–24], anthraquinones [25,26], anthocyanins [27,28], carbohydrates [29], catechins [30], coumarins [31,32], flavonoids [33–35], glucosinolates [36], phenolic acids [37], proanthocyanidins [38], red wine pigments [39], saponins [40], and xanthenes [41]. Due to the variety of factors influencing the separation selectivity and efficiency in CZE, an extensive discussion on method development strategies cannot be given in this review. Thus, for a comprehensive overview about the optimisation of CZE methods for the analysis of natural products, refer to the recently published review of Li et al. [42].

Non-Aqueous Capillary Electrophoresis

The first paper dealing with CE separations in a pure non-aqueous medium was written by Walbroehl and Jorgenson in 1984 [43]. Since then, non-aqueous buffer systems have been increasingly applied for the separation of small molecules including secondary plant metabolites [44]. The separation media in non-aqueous capillary electrophoresis (NACE) contain water-miscible organic solvents such as acetonitrile (ACN), dimethyl formamide (DMF), or various alcohols and acidic or basic solvents such as formic acid, acetic acid (AcOH), ammonia, or diethylamine [45]. Because the low conductivity of these buffer media normally results in lower currents compared with aqueous media, the electrolyte concentration and the electrical field strength can be increased. Another major advantage of non-aqueous media is the good solubility of less-polar compounds, for example, long-chained fatty acids [46]. According to Porras and Kenndler, the often-cited selectivity and efficiency improvements for CE analyses in non-aqueous media are based on heteroconjugation and/or ion pairing rather than on selective changes of the analyte mobilities or pK_a values [45]. As most buffer salts used for CE separations are not fairly soluble in non-aqueous media, the use of electrolytes in NACE is often restricted to ammonium acetate and ammonium formate, which also allows the application of these buffer systems for CE/MS analyses. In fact, one major reason for the popularity of non-aqueous running buffers is their applicability for CE/MS [47]. The high amount of organic solvents in non-aqueous

Table 1 Selected examples for the analysis of secondary plant metabolites with NACE.

Natural product class	Plant species	Electrolyte	Ref.
Alkaloids	–	50 mM ammonium acetate, 0.6 M AcOH in MeOH–ACN (75:25, v/v)	[48]
Anthraquinones	<i>Xanthophyllum atropvensis</i>	50 mM sodium cholate, 1.0% (v/v) AcOH, 40% (v/v) ACN in MeOH	[51]
Coumarins	Cortex Fraxini, <i>Fraxinus</i> spp.	60 mM sodium cholate, 20 mM ammonium acetate, 3.0% (v/v) AcOH, 20% (v/v) ACN in MeOH	[52]
Flavones	Chinese herbs	10 mM sodium cholate, 80 mM TRIS in ACN–MeOH (60:40, v/v)	[53]
Diterpenes	Danshen (<i>Salvia miltiorrhiza</i>)	250 mM ammonium acetate, 1.0% (v/v) AcOH, 30% (v/v) ACN in MeOH	[55]
Hypericins and hyperforins	St. John's wort (<i>Hypericum perforatum</i>)	50 mM ammonium acetate, 150 mM sodium acetate, 0.002% (w/v) hexadi-methrine bromide in MeOH–DMSO– <i>N</i> -methylformamide (3:2:1, v/v/v)	[57]

buffers increases their volatility and strongly reduces the surface tension of the droplets generated via electrospray ionisation (ESI). Therefore, non-aqueous media often provide a higher sensitivity and compatibility for CE/ESI-MS compared with aqueous electrolytes. Natural products analysed with NACE include alkaloids [48–50], anthraquinones [51], coumarins [52], flavonoids [53], polyphenolic compounds [54], terpenes [55,56], and hypericins and hyperforins from *Hypericum perforatum* [57]. Additional references are listed in the recently published reviews of Scriba [47] and Geiser and Veuthey [58]. Selected examples for NACE separations of natural products are given in **Table 1**.

Electrokinetic Chromatography

Neutral substances without acidic or basic functional groups and which cannot be transferred to charged complexes can be analysed with electrokinetic chromatography (EKC). In contrast to CZE, electrokinetic separation techniques (e.g., micellar electrokinetic chromatography [MEKC], microemulsion electrokinetic chromatography [MEEKC], and capillary electrochromatography [CEC]) are based on the distribution of analytes between a stationary or pseudo-stationary phase and the mobile phase driven by the EOF. In the following paragraphs the characteristics of these methods and their potential for natural products analysis will be explained.

Micellar electrokinetic chromatography

MEKC was introduced by Terabe and coworkers in 1984 [59] and rapidly proved to be the method of choice for the CE analysis of neutral compounds. In MEKC, a pseudo-stationary phase is generated by the addition of a micelle-forming ionic surfactant molecule such as sodium dodecyl sulphate (SDS) or cetyltrimethylammonium bromide. Above the critical micelle concentration, the tenside molecules aggregate to micelles where the polar and water-soluble “head” of the surfactant molecules constitutes the outer surface and the hydrophobic part of the molecule forms a lipophilic core. Whereas the selectivity of CZE separations is based on different electrophoretic mobilities of the analyte ions, the separation mechanism in MEKC is based on hydrophobic interactions of the sample molecules with the applied pseudo-stationary phase and – if the analytes are charged at the pH value of the buffer – different electrophoretic mobilities. In the case of charged compounds, the overlap of the two separation mechanisms (hydrophobic interaction and electrophoretic mobility) can lead to a significant gain in selectivity; therefore, MEKC has been frequently applied for challenging separations of natural products in crude plant extracts and pharmaceutical formulations [10–12]. Because crude plant extracts contain a variety of

acidic, basic, and neutral compounds, MEKC is of particular interest for the simultaneous analysis of charged and uncharged substances in crude plant extracts (e.g., for fingerprint analyses) [10–12,60]. For example, Ganzera et al. recently reported the successful MEKC separation of phenolic acids and flavonoids from *Arnica montana* [60]. Kaempferol 3-*O*-glucoside (**1**), 6-methoxy-kaempferol 3-*O*-glucoside (**2**), hispidulin (**3**), quercetin 3-*O*-glucoside (**4**), patuletin 3-*O*-glucoside (**5**), quercetin 3-*O*-glucuronic acid (**6**), chlorogenic acid (**7**), 3,5-dicaffeoylquinic acid (**8**), and 4,5-dicaffeoylquinic acid (**9**) were separated using an electrolyte consisting of 50 mM borate (pH 6.75), 25 mM SDS, and 30% (v/v) ACN (**Fig. 1**) [60]. The crude plant extracts were hydrodynamically injected (50 mbar for 5 s) into the fused silica capillary (62 cm × 50 μm i.d.), and the separation voltage and capillary temperature were 25 kV and 40 °C, respectively. As can be clearly seen in **Fig. 1**, the analytes migrated in two distinct groups according to their pK_a values. At a pH value of 6.75, the dissociation of hispidulin (**3**) and the flavonoid glycosides **1**, **2**, **4**, and **5** is negligible, so that the migration velocity of these compounds is predominantly determined by their affinity to the pseudo-stationary phase (SDS) and the formation of complexes with borate anions. For quercetin 3-*O*-glucuronic acid (**6**) and the phenolic acids (**7–9**), however, the longer migration times are due to the almost complete dissociation of their carboxylic groups. This deprotonation leads to a strong electrophoretic mobility to the anode but also to a reduced incorporation into the SDS micelles because of repulsion forces and higher polarity.

In addition to the above-mentioned example, MEKC has been frequently applied for the analysis of natural products, for example, alkaloids [61–63], anthraquinones [64,65], cardiac glycosides [66], catechins [67,68], coumarins [69], ecdysones [70], flavonoids [71–73], glucosinolates [74,75], isoflavones [76,77], procyanidins [78,79], saponins [80], and terpenes [81,82]. Examples for MEKC separations of secondary plant metabolites are given in **Table 2**.

Microemulsion electrokinetic chromatography

In recent years microemulsion electrokinetic chromatography (MEEKC) has become an attractive alternative to previously established CE methods such as CZE and particularly MEKC [83]. An important prerequisite for the performance of MEEKC analyses with UV detection is the use of running buffers with optically transparent emulsions containing oil droplets below 10 nm so that light scattering does not occur. Such microemulsions are obtained by dispersion of immiscible liquids in an aqueous buffer in the presence of surfactants that decrease the surface tension between the two liquid layers. The addition of a short-chain alcohol such as butanol (co-surfactant) further lowers the droplet's surface tension. Whereas the oil phase typically consists of heptane

Fig. 1 MEKC separation of two *Arnica montana* extracts under optimised CE conditions. For CE conditions and peak assignment refer to the text. (a) and (b) were tentatively identified as flavonoid and caffeoylquinic acid, respectively. Reprinted with permission from [60].

Table 2 Selected examples for the analysis of natural products with MEKC.

Natural product class	Plant species	Electrolyte	Ref.
Tropane alkaloids	<i>Hyoscyamus muticus</i>	30 mM phosphate–borate, 40 mM SDS, 16.5% (v/v) ACN, pH 8.7	[63]
Anthraquinones	Rhubarb (<i>Rheum raphonticum</i>)	30 mM phosphate, 20 mM SDS, 20 mM sodium cholate, 10 mM β -CD, pH 10.4	[65]
Catechins	Green tea (<i>Camellia sinensis</i>)	5 mM borate, 60 mM phosphate, 50 mM SDS, pH 7.0	[67]
Ecdysteroids and flavonoids	<i>Serratula strangulata</i>	15 mM borate, 60 mM SDS in 20% (v/v) methanol, pH 9.08	[70]
Flavonoids	<i>Ilexidium gracile</i>	15 mM borate, 30 mM SDS in 10% (v/v) ethanol, pH 10.5	[71]
Flavonoids	<i>Astragalus membranaceus</i>	20 mM borate, 100 mM sodium cholate in 25% (v/v) acetonitrile, pH 9.2	[73]
Isoflavones	<i>Medicago</i> spp., <i>Melilotus alba</i>	25 mM borate, 60 mM SDS, 1.6% (v/v) 1,2-hexanediol, pH 9.0	[76]
Flavonoids and terpene lactones	<i>Ginkgo biloba</i>	20 mM phosphoric acid, 40 mM SDS, 12 mM β -CD, pH 2.2	[82]

or octane, the aqueous phase comprises borate, phosphate, or TRIS buffers at basic pH values between 7 and 11; however, MEEKC analyses at low buffer pH values also have been reported [84,85]. Although most MEEKC applications use oil-in-water emulsions, the use of reverse MEEKC with water-in-oil emulsions is also possible and is particularly advantageous for the separation of highly hydrophobic compounds [86]. Comparable to MEKC, enantioselective microemulsions that contain chiral surfactants or chiral alcohols allow the enantioseparation of analytes in racemic mixtures. For example, Aiken and Huie described a novel chiral microemulsion based on (2*R*,3*R*)-di-*n*-butyl tartrate (0.5% w/w) as a water-immiscible chiral selector [87]. The addition of this chiral selector to the running buffer consisting of 0.6% (w/w) SDS and 1.2% (w/w) butanol in 15 mM TRIS (pH 8.1) allowed the enantiomeric separation of a racemic mixture of ephedrine (selectivity factor: 2.6) [87]. Especially because of its applicability for charged, neutral, and lipophilic compounds, MEEKC is an interesting alternative for the analysis of secondary plant metabolites complementary to CEC, MEKC, and RP-HPLC [83].

An impressive example of the MEEKC analysis of lipophilic and medium polar plant constituents has been published by Vanhoenacker et al., who separated the rather unstable hop ingredients humulone, adhumulone, and cohumulone (α -acids) and lupulone, adlupulone, and colupulone (β -acids) as well as the chalcone derivatives isoxanthohumol, 6-prenylnaringenin, and 8-

prenylnaringenin [86]. The separation was obtained using a running buffer containing 10 mM borate adjusted to a pH value of 9.7, 40 mM SDS, 3% (v/v) butanol, and 0.3% (v/v) heptane. Injections were performed hydrodynamically for 3 s at 50 mbar, and between two runs the capillary was rinsed sequentially for 1.5 min with 50 mM borate and then with the separation buffer. The temperature of the capillary was set to 25 °C and the applied separation voltage was 30 kV [86]. The pH value was an extremely important experimental condition, whereas an increase in the SDS concentration or the replacement of butanol by pentanol did not significantly affect the separation. By using the above-mentioned conditions, a baseline separation of the applied hop constituents could be obtained, and it was also possible to detect these compounds in hop extracts [86]. Other natural products that have been analysed with MEEKC include anthraquinones and bianthrone in rhubarb [88], catechins in green and black tea [84,85], ephedrine alkaloids in Chinese herbs [89], sesquiterpene lactones of radix inulae (*Inula helenium*) [90], and xanthones from *Securidaca inappendiculata* [91]. The buffer systems used for these MEEKC separations are given in **Table 3**.

Capillary electrochromatography

Capillary electrochromatography (CEC) can be addressed as a hybrid-technique that combines chromatography with capillary electrophoresis. In CEC the mobile phase is normally driven through the stationary phase by the EOF. In contrast to MEKC

Table 3 Selected examples for the analysis of secondary plant metabolites with MEEKC.

Natural product class	Plant species	Electrolyte	Ref.
Anthraquinones	Rhubarb (<i>Rheum raphonticum</i>)	97.7 % (w/v) 10 mM sodium borate (pH 9.2), 0.6 % (w/v) SDS, 0.5 % (w/v) dibutyl-L-tartrate, 1.2 % (w/v) butanol, 0–40 % (w/v) ACN	[88]
Catechins	Green tea (<i>Camellia sinensis</i>)	86.61–94.13 % (w/v) 50 mM phosphate (pH 2.5), 2.31–3.32 % (w/v) SDS, 1.36 % (w/v) heptane, 7.58–9.72 % (w/v) co-surfactant	[84]
Ephedrine alkaloids	Chinese herbs	20 mM borate, 23.3 mM SDS, 16.4 mM heptane, 180.85 mM butanol in 8 % ACN, pH 9.4	[89]
Phenolic acids and catechins	Black tea (<i>Camellia sinensis</i>)	86.1 % (v/v) 25 mM phosphate (pH 2.0), 2.89 % (w/v) SDS, 1.36 % (w/v) heptane, 7.66 % (w/v) cyclohexanol, 2 % (w/v) ACN	[85]
Sesquiterpene lactones	Radix Inulae (<i>Inula helenium</i>)	85.8 % (w/v) sodium tetraborate, 1.24 % (w/v) SDS, 0.32 % (v/v) hexane, 2.64 % (v/v) butanol, 10 % (v/v) ACN, pH 9.2	[90]
Xanthones	<i>Securidaca inappendiculata</i>	50 mM borate, 120 mM SDS, 80 mM heptane, 10 % (v/v) butanol, 5 mM sulphated β -CD, pH 9.5	[91]

Table 4 Selected examples for the analysis of secondary plant metabolites with CEC.

Substance class	Stationary phase	Mobile phase	Ref.
Purine alkaloids	Silica; 3 μ m	IsoOH, hexane, 1 mM TRIS (pH 8.0) 52:40:8	[93]
Anthraquinones	Hypersil C ₁₈ ; 3 μ m	5 mM AcOH–ACN 20:80	[94]
Boswellic acids	Hypersil C ₁₈ ; 3 μ m	20 mM ammonium formate (pH 6.5), ACN 1:9 (v/v)	[95]
Cannabinoids	Hypersil C _{8/18} ; 3 μ m	ACN, 25 mM phosphate (pH 2.57), 75:25	[96]
Flavanones	Lichrospher C ₁₈ ; 5 μ m	2.5 mM ammonium formate–ACN 80:20	[97]
Isoflavones	Monolith	2.4 mM ammonium formate (pH 2.7), ACN 70:30	[98]
Plant sterols	Hypersil C ₁₈ ; 3 μ m	ACN, THF, 25 mM TRIS (pH 8.0) 60:35:5	[99]
Withanolides	Hypersil C ₁₈ ; 3 μ m	10 mM ammonium acetate (pH 8.0), ACN 40:60	[100]

and MEEKC, the analyses are performed in coated, packed, or monolithic capillaries. There are three common types of stationary phases applied in CEC: (i) capillary columns with packed homogeneous silica or reversed-phase silica particles with a diameter of about 3 μ m; (ii) capillaries coated with the stationary phase; or (iii) monolithic capillary columns made by an *in situ* polymerisation of suitable polymer materials inside the capillary. Whereas for neutral compounds the separation mechanism is primarily based on interactions of the analytes with the applied stationary phase, charged compounds are separated because of their electrophoretic mobility and their affinity to the stationary phase. As for MEKC and MEEKC, this aspect is favourable for fingerprint analyses of crude plant extracts and often reveals excellent selectivities for complex separations. For the most often used reversed-phase stationary phases in CEC, the mobile phase typically consists of a mixture of organic solvents with buffers (e.g., phosphate or borate) [92]. Comparable to NACE and MEEKC, the use of mobile phases with high contents of organic solvents allows the separation of lipophilic natural products (e.g., boswellic acids, cannabinoids, or tocopherols) (► **Table 4**). This aspect contributed to the use of CEC and CEC/MS for the analytical characterisation of plant extracts [92]. Because a detailed overview of the application of CEC for the analysis of natural products was recently given by Scherz et al. [92], single applications will not be presented at this point. Instead, selected examples for the application of CEC in natural products analysis and the corresponding references are given in ► **Table 4**.

CE vs. HPLC in the Analysis of Natural Products

Although the performance of CE instruments and the handling of the capillaries have been improved in recent years, there are still some difficulties such as capillary conditioning, capillary cleaning, and blockage of the capillary due to particles or precipitation of buffer components. Especially the cleaning procedures between runs have to be carefully evaluated and validated, because they have a strong impact on the reproducibility of migration times and peak areas in fused silica capillaries [101]. For example, alkaloids tend to strongly adsorb to the capillary wall, which results in distorted peak shapes or sometimes a complete loss of the signal [18]. Although the use of running buffers with ammonium ions (e.g., ammonium acetate) reduces the adsorption of basic analytes to the capillary wall, exact cleaning procedures with water, sodium hydroxide, or even SDS [102] as well as a sufficient re-equilibration with the electrolyte are essential for reproducible results [101]. Because these cleaning procedures often last several minutes, cleaning and re-equilibration of the inner capillary surface often exceeds the analysis time. Whereas reproducibility, handling, and robustness are major advantages of conventional reversed-phase HPLC (RP-HPLC), the high separation efficiency and selectivity as well as short analysis times are favourable attributes of CE. However, short analysis times and high separation efficiencies were recently achieved for HPLC by the introduction of short HPLC columns with small diameters and particle sizes below 3 μ m. This technical progress led to the development of ultrahigh-pressure liquid chromatography, which was recently reviewed by Wu and Clausen [103]. Perhaps one of the most striking drawbacks of HPLC compared with CE is the limited applicability of normal-phase and RP-HPLC methods for the analysis of highly polar compounds such as phenolic

acids/glycosides, glucuronide and sulphate conjugates, purine and pyrimidine bases, nucleotides, and water-soluble vitamins. These polar compounds are normally not soluble in the non-aqueous solvents used for normal-phase chromatography, and they do not show a sufficient affinity to the most often applied reversed-phase silica gel. Furthermore, the application of amino-alkyl- or diol-based silica columns often does not provide reproducible results or sufficient robustness. Although the introduction of hydrophilic interaction chromatography was an important development for HPLC analysis of polar compounds with acetonitrile-water mixtures containing varying contents of alcohols and electrolytes [104], the analysis of highly water-soluble natural products with HPLC is still a challenge. For CE, however, the analysis of polar substances can be easily done with various methods (e.g., CZE, MEKC, or even NACE). For example, highly polar calystegines could be easily determined both qualitatively and quantitatively with CZE [105], whereas, in this case, the application of RP-HPLC was not possible. Another example is the use of CZE for the quantification of resveratrol in a food supplement in the presence of acesulfame K, riboflavin, ascorbic acid, flavanones, and hydroxycinnamic acids [106]. By using a running buffer consisting of 23 mM borate (pH 10.0) with 7% (v/v) acetonitrile, a baseline separation of all components was obtained in only seven minutes.

Its limited robustness and sometimes difficult manageability has hampered the establishment of CE, and especially CE/MS, in research facilities and quality-control laboratories. But, as outlined in the next section, the number of CE/MS applications has steadily increased as the demand for rapid, efficient, and multidimensional analytical methods increases.

Capillary Electrophoresis-Mass Spectrometry

Since its introduction in 1987 by Olivares et al. [107], CE/MS has evolved to a widely applicable, multidimensional analytical technique complementary to conventional LC/MS methods. One of the major problems that had to be solved for the application of CE/MS was the extremely small liquid flow inside the CE capillaries, which normally ranges from a few nanoliters to several hundred nanoliters per minute. Because even the typical flow rates of microscale LC/MS systems are in the low $\mu\text{L}/\text{min}$ range, the flow-out of the CE capillary has to be increased. For this purpose, a so-called make-up flow (ca. 1–5 $\mu\text{L}/\text{min}$) is added to the capillary flow via a simple tee (liquid junction) or as a coaxial sheath liquid (sheath flow). Although other online interfaces (e.g., sheathless, direct electrode) have been developed, nowadays almost all CE/MS applications use the sheath-flow interface introduced by Smith and coworkers in 1988 [108]. Whereas the sheath-liquid reduces the sensitivity due to additional background noise and the additional solvent, this approach also allows the post-column addition of chemicals in order to improve ESI characteristics and ionisation efficiency. This is particularly relevant if buffer additives with a low volatility or ion-pairing reagents must be used [109].

CE/MS analyses of natural products were first published in 1994 by Hsieh et al. for the qualitative analysis of the protoberberine alkaloids berberine and palmatine [110] and again in 1994 by Henion et al. for the quantitative determination of isoquinoline alkaloids in the bark of *Phellodendron wilsonii* [111]. Both groups used tandem MS and a sheath-flow interface instead of a liquid-junction interface to obtain a stable electrospray. In 1997, a gen-

erally applicable buffer system for the CZE and CE/MS analysis of various alkaloid classes was developed by Unger et al. [16]. A 1 : 1 mixture of 100 mM ammonium acetate, adjusted to pH 3.1, and acetonitrile allowed the CZE separation of monoterpene indole alkaloids, isoquinoline alkaloids, β -carboline, and opium alkaloids. By reducing the ammonium acetate concentration to 80 mM and adjusting the pH value to 4.0 before the solution was mixed with acetonitrile, the electrolyte was also suitable for the CE/MS analysis of the aforementioned alkaloid classes in crude extracts from opium and the cortex of *Aspidosperma quebracho-blanco* and from root and cell suspension cultures of *Rauwolfia serpentina* [112, 113]. Isoquinoline alkaloids were also analysed by Sturm et al., who applied a running buffer consisting of ammonium formate (70 or 100 mM), adjusted to pH 3.0 or 4.0, and methanol or acetonitrile as buffer additives [17]. By using a sheathflow (5 mM formic acid in acetonitrile) at a flow rate of 3 $\mu\text{L}/\text{min}$ and ESI in the positive mode, they obtained $[\text{M}]^+$ ions in the case of quaternary isoquinoline alkaloids such as berberine or sanguinarine and $[\text{M} + \text{H}]^+$ ions for the tertiary amines (e.g., chelidone). This method was successfully applied for the analytical characterisation of the alkaloid pattern in crude methanolic extracts of *Berberis vulgaris*, *Chelidonium majus*, *Eschscholzia californica*, *Jateorhiza palmata*, and *Hydrastis canadensis* [17]. Approximately 10 years later again Stuppner and coworkers applied CE/MS for the analysis of isoquinoline alkaloids [114]. In this case NACE and ion-trap tandem MS with a sheath-flow ESI interface was used for the analysis of isoquinoline alkaloids in central European *Corydalis* species. The non-aqueous electrolyte used for the NACE/ESI-/MS analysis was a mixture of 50 mM ammonium acetate, 1 M acetic acid, and 10% (v/v) methanol in acetonitrile [114]. Among the alkaloid classes also analysed with CE/MS are glycoalkaloids from *Solanum tuberosum* [115], naphthylisoquinoline alkaloids from a Central African *Ancistrocladus* species [116], quinolizidine alkaloids from *Sophora flavescens* [117], and tropane alkaloids from *Atropa belladonna* [118].

Naturally, the application of CE/MS is not restricted to alkaloids. Since 1997 almost all classes of natural products have been analysed with CE coupled to various types of MS instruments [6, 8, 15, 36, 92, 119, 120]. Both ion-trap and time-of-flight (TOF) MS have been applied for the CE/MS analysis of intact glucosinolates in *Arabidopsis thaliana* [36]. Due to the very low and almost identical pK_a values of the permanently charged sulphate groups of the intact glucosinolates, CZE analysis of these compounds at a pH value near the pK_a values of the analytes – which normally provides the optimum selectivity – was not possible. Furthermore, because glucosinolates form relatively stable ion pairs with ammonium ions, thereby reducing the sensitivity for MS detection, the use of ammonium acetate or formate for the running buffer and sheath liquid was avoided. By using 1 M and 0.2% (m/v) formic acid as electrolyte and sheath liquid, respectively, the glucosinolates could be successfully identified with CE/MS in crude extracts without interference from ubiquitous acidic plant constituents such as benzoic or cinnamic acids, which possess much higher pK_a values and thus are not negatively charged under the applied conditions. The intact glucosinolates, however, maintained their electrophoretic mobility even under the acidic conditions of the applied electrolyte (1 M formic acid) and migrated as anions, which are easily detectable with ESI in the negative-ion mode [36]. In **Fig. 2**, the CE/ESI-TOF-MS base peak electropherogram of the glucosinolate pattern of a crude *Arabidopsis thaliana* extract is shown. The application of CE/TOF-MS finally allowed the assignment of glucosinolates that could not be

Fig. 2 Base peak electropherogram as obtained by CE/ESI-TOF-MS of a crude *Arabidopsis thaliana* seed extract. The insert shows a detailed view of extracted ion electropherograms (m/z values) of the major compounds. Reprinted with permission from [36].

identified with ion-trap MS. However, as stated by the authors, even with TOF-MS it was not possible to unequivocally identify all peaks according to their elemental composition because of overlapping isotopic patterns of co-eluting compounds [36].

CE/MS vs. LC/MS in the Analysis of Natural Products

A direct comparison of the applicability of CE/MS and LC/MS for the analysis of natural products was published by Vanhoenacker et al., who analysed phenolic compounds in diethyl ether extracts of red wines [120]. For these analyses both CE/MS and LC/MS were performed with ESI in the negative-ion mode. Despite the higher separation efficiency of CE compared with HPLC, the application of LC/MS in this case was superior because better selectivity and sensitivity were obtained. Moreover, by applying a single-quadrupole mass spectrometer, 23 red wine constituents could be identified using LC/MS, whereas only 13 ingredients could be identified with CE/MS. Interestingly, the application of a more volatile ammonium acetate buffer (25 mM, pH 9.5) instead of an ammonium borate buffer (18.75 mM, pH 9.3) during CE/MS did not result in better sensitivity or improved peak identification. The sensitivity was increased for the catechins and the phenolic acids, but the separation and the signal intensities of the flavonols decreased [120]. The main reason for this result was the significant loss in separation selectivity when borate was replaced by acetate. Obviously, the complexation of aromatic or *cis*-configured vicinal hydroxy groups by borate anions was the key to the successful separation of the flavonols. This complexation also influenced the ionisation behaviour and fragmentation of the phenolic compounds, because the red wine constituents with a vicinal hydroxy group preferentially showed cluster ions

with borate or borate and methanol from the sheath liquid [120]. However, it should be mentioned that for CE/MS analyses the application of single-quadrupole mass spectrometers is particularly unfavourable for fingerprint analyses of natural products because these instruments show a relatively low sensitivity in scan mode. That is why most of the CE/MS analyses of crude plant extracts have been performed with ion-trap MS, which provides a much higher sensitivity in scan mode.

Online Pre-Concentration Methods

Whereas in CE very low absolute sample amounts, typically in the low picogram range, are detected, the low injection volumes and the narrow diameter of the capillaries typically used (25–100 μm) lead to a poor concentration sensitivity for the most often applied UV detection. For molecules with average extinction coefficients, such as flavones, the limit of detection often lies between 1 and 10 $\mu\text{g/mL}$. Even for CE coupled to MS, the concentration sensitivity is only somewhat higher compared with CE with UV detection because the sheath liquid necessary to obtain a stable nanospray dilutes the sample and causes additional background noise [109]. To overcome the low concentration sensitivity of CE analyses, several strategies are currently applied. These strategies involve the use of capillaries with extended detection path lengths (e.g., Z-shaped, multi-reflection, bubble cell) or the application of highly sensitive detection methods such as electrochemical or fluorometric detection. The combination of an extended detection path length with laser-induced fluorescence (LIF) was reported recently for the sensitive detection of riboflavin and other aromatic compounds [121]. Because the laser intensity had to be decreased when bubble cells were used, the sensitivity for riboflavin, for example, was only 8 times higher compared with conventional UV detection [121]. Nevertheless, the use of LIF detectors generally results in a significant increase in sensitivity if the target analytes possess a fluorophoric structure. Although one major drawback is the limited availability of excitation and emission wavelengths for LIF detectors, this technique has been applied several times in the past for the CE analysis of natural products [122–124]. Comparable to catecholamines, many flavonoids can be electrochemically detected with high sensitivity because the catechol structures of, e.g., quercetin or luteolin can be easily oxidised to *ortho*-quinones. Thus, electrochemical detection was preferentially applied for the CE analysis of flavonoids [125, 126].

An alternative strategy to the use of extended detection path lengths or more sensitive detection methods is the application of online pre-concentration methods such as stacking or sweeping. For a comprehensive overview of these methods, refer to the excellent review of Simpson et al. [127]. Originally applied for charged compounds in CZE applications, online pre-concentration techniques are nowadays also widely used for the MEKC or MEEKC analysis of natural products [128, 129]. During online pre-concentration a relatively large sample amount is introduced into the CE capillary with hydrodynamic or electrokinetic injection, and the target analytes are focused in a narrow band before the separation begins. The focusing of sample molecules most often results from (i) a different field strength in the sample zone and running buffer (e.g., stacking, field-enhanced sample stacking, FESS); (ii) a change in the effective charge of the analyte (e.g., dynamic pH junction); or (iii) the partitioning of the sample molecules in a pseudo-stationary phase such as SDS (sweeping).

The application of these online pre-concentration techniques provides a 10- to 1000-fold enhancement in concentration sensitivity, or even higher if large sample amounts can be introduced into the capillary [127]. The first application of a simple and effective online pre-concentration method for the CE analysis of natural products in crude plant extracts was published by Unger and Stöckigt in 1997 [130]. They used field-amplified sample injection (FASI) for the determination of alkaloids in crude methanolic extracts from the roots of *Berberis vulgaris* and *Hydrastis canadensis*. By introducing a short plug of methanol 70% (v/v) before electrokinetic injection of the alkaloids at 16 kV for 8 s, the concentration sensitivity was 1000 times higher compared with hydrodynamic injection at 345 mbar for 1 s [130].

The online enrichment of charged substances with FASI is not limited to CZE separations and can be combined with other CE separation methods such as MEEKC. The combination of FASI with MEEKC was used by Yu et al. to analyse trace amounts of the quinolizidine alkaloids sophoridine, matrine, oxymatrine, oxysophocarpine, and cytosine in the roots of *Sophora flavescens* [129]. In this case the obtained LODs (defined as signal-to-noise ratio = 3) for the analytes were as low as 0.1 ng/mL, which is approximately 10000 times higher compared with conventional hydrodynamic injection.

Among the various online pre-concentration methods, FESS, FASI, dynamic pH junction, and sweeping can be considered the most useful techniques for the analysis of trace amounts of natural products in crude plant extracts, pharmaceutical formulations, and even body fluids such as urine or plasma [127]. For example, a combination of electrokinetic injection, dynamic pH junction, and sweeping was used for the sensitive determination of sinapic, ferulic, coumarinic, caffeic, syringic, vanillic, and 4-hydroxybenzoic acid in crude acetone extracts from *Majorana hortensis* [131]. The sample was dissolved in a basic borate buffer (50 mM, pH 9.5), and both the junction electrolyte and the running buffer (mobilisation electrolyte) consisted of 50 mM phosphate, pH 2.5. The latter also contained 60 mM SDS because the mobilisation and separation of the organic acids were done by reversed-migration MEKC (RM-MEKC). In RM-MEKC the running buffer is adjusted to a low pH value (<2.5) where virtually no EOF is present or the EOF is suppressed by coating the capillaries [132]. Therefore, the micelles with the partially included analytes migrate to the oppositely charged electrode without being swept by the EOF. For online pre-concentration, the phenolic acids were electrokinetically injected (–10 kV, 30 min) from the alkaline sample buffer (pH 9.5) into the junction electrolyte (pH 2.5); at this point the phenolic acids were neutralised and stacked (accumulation part). After accumulation of the phenolic acids, the sample solution (inlet vial) was replaced by the running buffer (50 mM phosphate, 60 mM SDS, pH 2.5) and the separation voltage (–10 kV) was applied. The negatively charged SDS micelles of the inlet vial migrated to the anode and penetrated the junction electrolyte, where the neutralised phenolic acids were stacked due to the loss in electrophoretic mobility at the acidic pH value of 2.5. Because of their higher lipophilicity, the neutralised phenolic acids were efficiently incorporated into the SDS micelles and swept to the anode. The obtained LODs between 0.4 and 4.2 ng/mL mainly resulted from the long duration used for electrokinetic injection (30 min) and the stacking process during the accumulation of the negatively charged phenolic acids in the acidic junction electrolyte [131]. However, it can be speculated that the application of a permanently or dynamically coated capillary and a simple electrokinetic injection or FASI at relatively

high electrical field strengths may have provided similar results regarding sensitivity enhancement and separation efficiency.

Sweeping was originally developed for the online concentration of neutral analytes in MEKC [133]. In this online concentration technique, neutral or charged analytes are picked up and accumulated in micelles of a pseudo-stationary phase that penetrates the sample zone. Because the analytes are incorporated into the micelles, lipophilic molecules are more efficiently concentrated than hydrophilic substances. Also, in order to obtain a strong incorporation of the analytes into the pseudo-stationary phase, buffer additives such as cyclodextrins or organic solvents should be avoided in the sample matrix. In recent years, sweeping has been applied several times for the online pre-concentration of natural products, e.g., alkaloids [134], catechins [135], flavonoids [128], and phenolic acids [136]. By introducing a large sample amount into the capillary via hydrodynamic or electrokinetic injection, sweeping can lead to a dramatic increase in sensitivity (> 1000-fold) [127, 133]. For example, a 1500-fold improvement in detection sensitivity was obtained by a sweeping technique applied for the MEKC analysis of *trans*-resveratrol in red wine [137].

Future Perspectives and Concluding Remarks

▼ The ongoing technical advances and future developments in all fields of instrumental analytics will lead to further miniaturisation and improvement of CE instruments. Such improvements may soon lead to the routine CE analysis of very small absolute sample amounts in the low femtogram range. This sensitivity may be achieved by a combination of various online pre-concentration techniques with highly sensitive detection methods, such as fluorometry or electrochemiluminescence [138].

Because the interest in nanoscale separation techniques is high and still growing, CE will be the method of choice for future applications in all areas of bioanalytical research. Due to the widespread use of a great variety of medicinal plants for the production of herbal supplements and phytomedicinal products, one of the future challenges in natural products analysis will be the detection and exact quantification of trace amounts of species-specific constituents (marker compounds) for the unambiguous identification of authentic plant material. Furthermore, CE will be an important tool for the detection and quantification of trace amounts of harmful plant constituents or impurities from toxic plants, for example, aristolochic acids or colchicine. In this regard, it can be speculated that the future application of CE in natural products analysis is directly linked to the availability of robust and highly sensitive detection methods.

Acknowledgements

▼ The assistance of A. Frank in the preparation of the manuscript is gratefully acknowledged.

References

- 1 Mikkers FEP, Everaerts FM, Verheggen ThPEM. High-performance zone electrophoresis. *J Chromatogr* 1979; 169: 11–20
- 2 Jorgenson JW, Lukacs KD. Zone electrophoresis in open-tubular glass capillaries. *Anal Chem* 1981; 53: 1298–1302
- 3 Jorgenson JW, Lukacs KD. Free-zone electrophoresis in glass capillaries. *Clin Chem* 1981; 27: 1551–1553

- 4 Altria KD, Elder D. Overview of the status and applications of capillary electrophoresis to the analysis of small molecules. *J Chromatogr A* 2004; 1023: 1–14
- 5 Cheung RH, Marriot PJ, Small DM. CE methods applied to the analysis of micronutrients in foods. *Electrophoresis* 2007; 28: 3390–3413
- 6 Huck CW, Stecher G, Scherz H, Bonn G. Analysis of drugs, natural and bioactive compounds containing phenolic groups by capillary electrophoresis coupled to mass spectrometry. *Electrophoresis* 2005; 26: 1319–1333
- 7 Kostal V, Katzenmeyer J, Arriaga EA. Capillary electrophoresis in bioanalysis. *Anal Chem* 2008; 80: 4533–4550
- 8 Huo Y, Kok WT. Recent applications in CEC. *Electrophoresis* 2008; 29: 80–93
- 9 Kraly J, Fazal MA, Schoenherr RM, Bonn R, Harwood MM, Turner E, Jones M, Dovichi NJ. Bioanalytical applications of capillary electrophoresis. *Anal Chem* 2006; 78: 4097–4110
- 10 Issaq HJ. Capillary electrophoresis of natural products. *Electrophoresis* 1997; 18: 2438–2452
- 11 Issaq HJ. Capillary electrophoresis of natural products-II. *Electrophoresis* 1999; 20: 3190–3202
- 12 Suntornsuk L. Capillary electrophoresis of phytochemical substances. *J Pharm Biomed Anal* 2002; 27: 679–698
- 13 Bednár P, Papoušková B, Müller L, Barták P, Stávek J, Pavloušek P, Lemr K. Utilization of capillary electrophoresis/mass spectrometry (CE/MSn) for the study of anthocyanin dyes. *J Sep Sci* 2005; 28: 1291–1299
- 14 Petersson EV, Puerta A, Bergquist J, Turner C. Analysis of anthocyanins in red onion using capillary electrophoresis-time of flight-mass spectrometry. *Electrophoresis* 2008; 29: 2723–2730
- 15 Seguera-Carretero A, Puertas-Mejía MA, Cortacero-Ramírez S, Beltrán R, Alonso-Villaverde C, Joven J, Dinelli G, Fernández-Gutiérrez A. Selective extraction, separation, and identification of anthocyanins from *Hibiscus sabdariffa* L. using solid phase extraction-capillary electrophoresis-mass spectrometry (time-of-flight/ion trap). *Electrophoresis* 2008; 29: 2852–2861
- 16 Unger M, Stöckigt D, Belder D, Stöckigt J. General approach for the analysis of various alkaloid classes using capillary electrophoresis and capillary electrophoresis-mass spectrometry. *J Chromatogr A* 1997; 767: 263–276
- 17 Sturm S, Stuppner H. Analysis of isoquinoline alkaloids in medical plants by capillary electrophoresis-mass spectrometry. *Electrophoresis* 1998; 19: 3026–3032
- 18 Unger M. Capillary zone electrophoresis of alkaloids – influence of structure on electrophoretic mobility. *J Chromatogr A* 1998; 807: 81–87
- 19 Lechtenberg M, Quandt B, Nahrstedt A. Quantitative determination of curcuminoids in *Curcuma* rhizomes and rapid differentiation of *Curcuma domestica* Val. and *Curcuma xanthorrhiza* Roxb. by capillary electrophoresis. *Phytochem Anal* 2004; 15: 152–158
- 20 Honda S, Iwase S, Makino A, Fujiwara S. Simultaneous determination of reducing monosaccharides by capillary zone electrophoresis as the borate complexes of N-2-pyridylglycines. *Anal Biochem* 1989; 176: 72–77
- 21 Schwaiger H, Oefner PJ, Huber C, Grill E, Bonn GK. Capillary zone electrophoresis and micellar electrokinetic chromatography of 4-aminobenzonitrile carbohydrate derivatives. *Electrophoresis* 1994; 15: 941–952
- 22 Zong YY, Che CT. Determination of strychnine and brucine by capillary zone electrophoresis. *Planta Med* 1995; 61: 456–458
- 23 Eeva M, Salo JP, Oksman-Caldentey KM. Determination of the main tropane alkaloids from transformed *Hyoscyamus muticus* plants by capillary zone electrophoresis. *J Pharm Biomed Anal* 1998; 16: 717–722
- 24 Unger M, Laug S, Holzgrabe U. Capillary zone electrophoresis as a tool for the quality control of goldenseal extracts. *Electrophoresis* 2005; 26: 2430–2436
- 25 Koyama J, Morita I, Fujiyoshi H, Kobayashi N. Simultaneous determination of anthraquinones, their 8- β -D-glucosides, and sennosides of Rhei Rhizoma by capillary electrophoresis. *Chem Pharm Bull* 2005; 53: 573–575
- 26 Tian K, Zhang H, Chen X, Hu Z. Determination of five anthraquinones in medicinal plants by capillary zone electrophoresis with beta-cyclodextrin addition. *J Chromatogr A* 2006; 1123: 134–137
- 27 Sáenz-López R, Fernández-Zurbano P, Tena MT. Development and validation of a capillary zone electrophoresis method for the quantitative determination of anthocyanins in wine. *J Chromatogr A* 2003; 990: 247–258
- 28 Ichiyanagi T, Kashiwada Y, Ikeshiro Y, Hatano Y, Shida Y, Horie Y, Matsugo S, Konishi T. Complete assignment of bilberry (*Vaccinium myrtillus* L.) anthocyanins separated by capillary zone electrophoresis. *Chem Pharm Bull* 2004; 51: 226–229
- 29 Vopli N, Maccari F, Linhardt RJ. Capillary electrophoresis of complex natural polysaccharides. *Electrophoresis* 2008; 29: 3095–3106
- 30 Arce L, Ríos A, Valcárcel M. Determination of anti-carcinogenic polyphenols present in green tea using capillary electrophoresis coupled to a flow injection system. *J Chromatogr A* 1998; 827: 113–120
- 31 Günaydin K, Erim FB. Determination of khellin and visnagin in *Ammi visnaga* fruits by capillary electrophoresis. *J Chromatogr A* 2002; 954: 291–294
- 32 Yue ME, Jiang TF, Liu X, Shi YP. Separation and determination of coumarins from *Cacalia tangutica* by capillary zone electrophoresis. *Biomed Chromatogr* 2005; 19: 250–254
- 33 Seitz U, Oefner PJ, Nathakarnkitkool S, Popp M, Bonn GK. Capillary electrophoretic analysis of flavonoids. *Electrophoresis* 1992; 13: 35–38
- 34 Quaglia MG, Bossù E, Donati E, Mazzanti G, Brandt A. Determination of silymarin in the extract from the dried silybum marianum fruits by high performance liquid chromatography and capillary electrophoresis. *J Pharm Biomed Anal* 1999; 19: 435–442
- 35 Marchart E, Krenn L, Kopp B. Quantification of the flavonoid glycosides in *Passiflora incarnata* by capillary electrophoresis. *Planta Med* 2003; 69: 452–456
- 36 Bringmann G, Kajahn I, Neuss C, Pelzing M, Laug S, Unger M, Holzgrabe U. Analysis of the glucosinolate pattern of *Arabidopsis thaliana* seeds by capillary zone electrophoresis coupled to electrospray ionization-mass spectrometry. *Electrophoresis* 2005; 26: 1513–1522
- 37 Trute A, Nahrstedt A. Identification and quantitative analysis of phenolic compounds from the dry extract of *Hedera helix*. *Planta Med* 1997; 63: 177–179
- 38 Kreimeyer J, Peterleit F, Nahrstedt A. Separations of flavan-3-ols and dimeric proanthocyanidins by capillary electrophoresis. *Planta Med* 1998; 64: 63–67
- 39 Sáenz-López R, Fernández-Zurbano P, Tena MT. Analysis of aged red wine pigments by capillary zone electrophoresis. *J Chromatogr A* 2004; 1052: 191–197
- 40 Emara S, Mohamed KM, Masujima T, Yamasaki K. Separation of naturally occurring triterpenoidal saponins by capillary zone electrophoresis. *Biomed Chromatogr* 2001; 15: 252–256
- 41 Wu X, Gong S, Bo T, Liao Y, Liu H. Determination of dissociation constants of pharmacologically active xanthenes by capillary zone electrophoresis with diode array detection. *J Chromatogr A* 2004; 1061: 217–223
- 42 Li P, Li SP, Wang YT. Optimization of CZE for analysis of phytochemical bioactive compounds. *Electrophoresis* 2006; 27: 4808–4819
- 43 Walbroehl Y, Jorgenson JW. On-column UV absorption detector for open tubular capillary zone electrophoresis. *J Chromatogr* 1984; 315: 135–143
- 44 Bjornsdottir I, Tjornelund J, Jansen SH. Nonaqueous capillary electrophoresis – its applicability in the analysis of food, pharmaceuticals and biological fluids. *Electrophoresis* 1998; 19: 2179–2186
- 45 Porras SP, Kenndler E. Are the asserted advantages of organic solvents in capillary electrophoresis real? A critical discussion. *Electrophoresis* 2005; 26: 3203–3220
- 46 Wang T, Wei H, Li SF. Nonaqueous capillary zone electrophoresis for separation of free fatty acids with indirect fluorescence detection. *Electrophoresis* 1998; 19: 2187–2192
- 47 Scriba GK. Non-aqueous capillary electrophoresis-mass spectrometry. *J Chromatogr A* 2007; 1159: 28–41
- 48 Barthe L, Ribet JP, Pélissou M, Degude MJ, Fahy J, Duflos A. Optimization of the separation of Vinca alkaloids by nonaqueous capillary electrophoresis. *J Chromatogr A* 2002; 968: 241–250
- 49 Li Y, He X, Qi S, Gao W, Chen X, Hu Z. Separation and determination of strychnine and brucine in *Strychnos nux-vomica* L. and its preparation by nonaqueous capillary electrophoresis. *J Pharm Biomed Anal* 2006; 41: 400–407
- 50 Liu Q, Liu Y, Li Y, Yao S. Nonaqueous capillary electrophoresis coupled with laser-induced native fluorescence detection for the analysis of berberine, palmatine, and jatrorrhizine in Chinese herbal medicines. *J Sep Sci* 2006; 29: 1268–1274
- 51 Li Y, Qi S, Chen X, Hu Z. Separation and determination of the anthraquinones in *Xanthophyllum atropvensis* Pierre by nonaqueous capillary electrophoresis. *Talanta* 2005; 65: 15–20

- 52 Li C, Chen A, Chen X, Ma X, Chen X, Hu Z. Non-aqueous capillary electrophoresis for separation and simultaneous determination of fraxin, esculetin and esculetin in *Cortex fraxini* and its medicinal preparations. *Biomed Chromatogr* 2005; 19: 696–702
- 53 Qi S, Li Y, Wu S, Chen X, Hu Z. Novel nonaqueous capillary electrophoresis separation and determination of bioactive flavone derivatives in Chinese herbs. *J Sep Sci* 2005; 28: 2180–2186
- 54 Demianová Z, Sirén H, Kuldvee R, Riekkola ML. Nonaqueous capillary electrophoresis separation of polyphenolic compounds in wine using coated capillaries at high pH in methanol. *Electrophoresis* 2003; 24: 4264–4271
- 55 Che AJ, Zhang JY, Li CH, Chen XF, Hu ZD, Chen XG. Separation and determination of active components in *Radix Salviae miltiorrhizae* and its medicinal preparations by nonaqueous capillary electrophoresis. *J Sep Sci* 2004; 27: 569–575
- 56 Qi S, Ding L, Tian K, Chen X, Hu Z. Novel and simple nonaqueous capillary electrophoresis separation and determination of bioactive triterpenes in Chinese herbs. *J Pharm Biomed Anal* 2006; 40: 35–41
- 57 Jensen AG, Hansen SH. Separation of hypericins and hyperforins in extracts of *Hypericum perforatum* L. using non-aqueous capillary electrophoresis with reversed electroosmotic flow. *J Pharm Biomed Anal* 2002; 27: 167–176
- 58 Geiser L, Veuthey JL. Nonaqueous capillary electrophoresis in pharmaceutical analysis. *Electrophoresis* 2007; 28: 45–57
- 59 Terabe S, Otsuka K, Ichikawa K, Tsuchiya A, Ando T. Electrokinetic separations with micellar solutions and open-tubular capillaries. *Anal Chem* 1984; 56: 111–113
- 60 Ganzera M, Egger C, Zidorn C, Stuppner H. Quantitative analysis of flavonoids and phenolic acids in *Arnica montana* L. by micellar electrokinetic capillary chromatography. *Anal Chim Acta* 2008; 614: 196–200
- 61 Kuo CH, Sun SW. Separation of bisbenzylisoquinoline alkaloids by micellar electrokinetic chromatography. *Phytochem Anal* 2002; 13: 63–68
- 62 Yu L, Xu Y, Feng H, Li SF. Separation and determination of toxic pyrrolizidine alkaloids in traditional Chinese herbal medicines by micellar electrokinetic chromatography with organic modifier. *Electrophoresis* 2005; 26: 3397–3404
- 63 Mateus L, Cherkaoui S, Christen P, Oksman-Caldentey KM. Simultaneous determination of scopalamine, hyoscyamine and littorine in plants and different hairy root clones of *Hyoscyamus muticus* by micellar electrokinetic chromatography. *Phytochemistry* 2000; 54: 517–523
- 64 Kuzuya H, Tamai I, Beppu H, Shimpo K, Chibara T. Determination of aloenin, barbaloin and isobarbaloin in aloe species by micellar electrokinetic chromatography. *J Chromatogr B* 2001; 752: 91–97
- 65 Shang X, Yuan Z. Determination of hydroxyanthraquinoids in Rhubarb by cyclodextrin-modified micellar electrokinetic chromatography using a mixed micellar system of sodium dodecyl sulfate and sodium cholate. *J Pharm Biomed Anal* 2003; 31: 75–81
- 66 Debusschère L, Demesmay C, Rocca JL, Lachatre G, Lofti H. Separation of cardiac glycosides by micellar electrokinetic chromatography and microemulsion electrokinetic chromatography. *J Chromatogr A* 1997; 779: 227–233
- 67 Weiss DJ, Austria EJ, Anderton CR, Hompesch R, Jander A. Analysis of green tea extract supplements by micellar electrokinetic chromatography. *J Chromatogr A* 2006; 1117: 103–108
- 68 Gotti R, Furlanetto S, Pinzauti S, Carvrini V. Analysis of catechins in *Theobroma cacao* beans by cyclodextrin-modified micellar electrokinetic chromatography. *J Chromatogr A* 2006; 1112: 345–352
- 69 Wang SF, Ju Y, Chen XG, Hu ZD. Separation and determination of coumarins in the root bark of three citrus plants by micellar electrokinetic capillary chromatography. *Planta Med* 2003; 69: 483–486
- 70 Wang S, Dai J, Chen X, Hu Z. Identification and determination of eclysones and flavonoids in *Serratula stragulata* by micellar electrokinetic capillary chromatography. *Planta Med* 2002; 68: 1029–1033
- 71 Zhang Y, Chen J, Ma XM, Shi YP. Simultaneous determination of flavonoids in *Ixeridium gracile* by micellar electrokinetic chromatography. *J Pharm Biomed Anal* 2007; 45: 742–746
- 72 Pietta P, Bruno A, Mauri P, Rava A. Separation of flavonol-2-O-glycosides from *Calendula officinalis* and *Sambucus nigra* by high-performance liquid and micellar electrokinetic capillary chromatography. *J Chromatogr* 1992; 593: 165–170
- 73 Song Y, Li P, Wang D, Cheng YY. Micellar electrokinetic chromatography for the quantitative analysis of flavonoids in the radix of *Astragalus membranaceus* var. *mongholicus*. *Planta Med* 2008; 74: 84–89
- 74 Karcher A, El Rassi Z. Capillary electrophoresis of glucosinolates and their degradation products. *Electrophoresis* 1999; 20: 3181–3189
- 75 Michaelsen S, Moller P, Sorensen H. Factors influencing the separation and quantitation of intact glucosinolates and desulphoglucosinolates by micellar electrokinetic capillary chromatography. *J Chromatogr* 1992; 608: 363–374
- 76 Baggett BR, Cooper JD, Hogan ET, Carper J, Paiva NL, Smith JT. Profiling isoflavonoids found in legume root extracts using capillary electrophoresis. *Electrophoresis* 2002; 23: 1642–1651
- 77 Zhang Y, Chen J, Zhao L, Shi YP. Separation and determination of isoflavones in red clover by micellar electrokinetic capillary chromatography. *Biomed Chromatogr* 2007; 21: 987–992
- 78 Cifuentes A, Bartolomé B, Gómez-Cardovés C. Fast determination of procyanidins and other phenolic compounds in food samples by micellar electrokinetic chromatography using acidic buffers. *Electrophoresis* 2001; 22: 1561–1567
- 79 Herrero-Martínez JM, Ràfols C, Rosés M, Bosch E, Lozano C, Torres JL. Micellar electrokinetic chromatography estimation of size and composition of procyanidins after thiolysis with cysteine. *Electrophoresis* 2003; 24: 1404–1410
- 80 Glöckl I, Veit M, Blaschke G. Determination of ginsenosides from *Panax ginseng* using micellar electrokinetic chromatography. *Planta Med* 2002; 68: 158–161
- 81 Petriska I, Polásek M, Sladkovský R. Assay of terpene alcohols in pharmacopoeial essential oils by micellar electrokinetic capillary chromatography (MEKC). *Pharmazie* 2006; 61: 989–993
- 82 Dubber MJ, Kanfer I. Application of reverse-flow micellar electrokinetic chromatography for the simultaneous determination of flavonols and terpene trilactones in *Ginkgo biloba* dosage forms. *J Chromatogr A* 2006; 1122: 266–274
- 83 Huie CW. Recent applications of microemulsion electrokinetic chromatography. *Electrophoresis* 2006; 27: 60–75
- 84 Pomponio R, Gotti R, Luppi B, Cavrini V. Microemulsion electrokinetic chromatography for the analysis of green tea catechins: effect of the cosurfactant on the separation selectivity. *Electrophoresis* 2003; 24: 1658–1667
- 85 Huang HY, Lien WC, Chiu CW. Comparison of microemulsion electrokinetic chromatography and micellar electrokinetic chromatography methods for the analysis of phenolic compounds. *J Sep Sci* 2005; 28: 973–981
- 86 Vanhoenacker G, Rong H, De Keukeleire D, Baeyens W, Van Der Weken G, Sandra P. Simultaneous analysis of hop acids and prenylated flavanones by microemulsion electrokinetic chromatography with diode array detection. *Biomed Chromatogr* 2000; 14: 34–36
- 87 Aiken JH, Huie CW. Use of a microemulsion system to incorporate a lipophilic chiral selector in electrokinetic capillary chromatography. *Chromatographia* 1993; 35: 448–503
- 88 Sun SW, Yeh PC. Analysis of rhubarb anthraquinones and bianthrone by microemulsion electrokinetic chromatography. *J Pharm Biomed Anal* 2005; 36: 995–1001
- 89 Zhang J, Xie J, Liu J, Tian J, Chen X, Hu Z. Microemulsion electrokinetic chromatography with laser-induced fluorescence detection for sensitive determination of ephedrine and pseudoephedrine. *Electrophoresis* 2004; 25: 74–79
- 90 Wenhua G, Yaoen C, Yegao Y, Xingguo C, Zhide H. Separation and determination of two sesquiterpene lactones in *Radix inulae* and *Liuwei Anxian* San by microemulsion electrokinetic chromatography. *Biomed Chromatogr* 2004; 18: 826–832
- 91 Bo T, Yang X, Li KA, Xu L, Liu H. Comparison of micellar electrokinetic chromatography and microemulsion electrokinetic chromatography for separation of pharmacologically active xanthenes from *Securidaca inappendiculata*. *J Sep Sci* 2003; 26: 133–136
- 92 Scherz H, Huck CW, Bonn GK. CEC and EKC of natural compounds. *Electrophoresis* 2007; 28: 1645–1657
- 93 Lai EP, Dabek-Zlotorzynska E. Separation of theophylline, caffeine and related drugs by normal-phase capillary electrochromatography. *Electrophoresis* 1999; 20: 2366–2372
- 94 Li Y, Liu H, Ji X, Li J. Optimized separation of pharmacologically active anthraquinones in Rhubarb by capillary electrochromatography. *Electrophoresis* 2000; 21: 3109–3115
- 95 Ganzera M, Stöggel WM, Bonn GK, Khan IA, Stuppner H. Capillary electrochromatography of boswellic acids in *Boswellia serrata* Roxb. *J Sep Sci* 2003; 26: 1383–1388
- 96 Lurie IS, Meyers RP, Conner TS. Capillary electrochromatography of cannabinoids. *Anal Chem* 1998; 70: 3255–3260

- 97 Desiderio C, De Rossi A, Sinibaldi M. Analysis of flavanone-7-O-glycosides in citrus juices by short-end capillary electrochromatography. *J Chromatogr A* 2005; 1081: 99–104
- 98 Starkey JA, Mechref Y, Byun CK, Steinmetz R, Fuqua JS, Pescovitz OH, Novotny MV. Determination of trace isoflavone phytoestrogens in biological materials by capillary electrochromatography. *Anal Chem* 2002; 71: 5998–6005
- 99 Abidi SL. Capillary electrochromatography of sterols and related sterol esters derived from vegetable oils. *J Chromatogr A* 2004; 1059: 199–208
- 100 Cherkaoui S, Cahours X, Veuthey JL. Analysis of selected withanolides in plant extracts by capillary electrochromatography and microemulsion electrokinetic chromatography. *Electrophoresis* 2003; 24: 336–342
- 101 Wätzig H, Degenhardt M, Kunkel A. Strategies for capillary electrophoresis: method development and validation for pharmaceutical and biological applications. *Electrophoresis* 1998; 19: 2695–2752
- 102 Lloyd DK, Wätzig H. Sodium dodecyl sulfate solution is an effective between-run rinse for capillary electrophoresis of samples in biological matrices. *J Chromatogr B* 1995; 663: 400–405
- 103 Wu N, Clausen AM. Fundamental and practical aspects of ultrahigh pressure liquid chromatography for fast separations. *J Sep Sci* 2007; 30: 1167–1182
- 104 Hemström P, Irgum K. Hydrophilic interaction chromatography. *J Sep Sci* 2006; 29: 1784–1821
- 105 Dräger B. Analysis of tropane related alkaloids. *J Chromatogr A* 2002; 978: 1–35
- 106 Orlandini S, Giannini I, Pinzauti S, Furlanetto S. Multivariate optimisation and validation of a capillary electrophoresis method for the analysis of resveratrol in a nutraceutical. *Talanta* 2008; 74: 570–577
- 107 Olivares JA, Nguyen NT, Yonker CR, Smith RD. On-line mass spectrometric detection for capillary zone electrophoresis. *Anal Chem* 1987; 59: 1230–1232
- 108 Smith RD, Barinaga CJ, Udseth HR. Improved electrospray ionization interface for capillary zone electrophoresis-mass spectrometry. *Anal Chem* 1988; 60: 1948–1952
- 109 Banks JF. Recent advances in capillary electrophoresis/electrospray/mass spectrometry. *Electrophoresis* 1997; 18: 2255–2266
- 110 Hsieh FY, Cai J, Henion J. Determination of trace impurities of peptides and alkaloids by capillary electrophoresis-ion spray mass spectrometry. *J Chromatogr A* 1994; 679: 206–211
- 111 Henion JD, Mordehai AV, Cai J. Quantitative capillary electrophoresis ion spray mass spectrometry on a benchtop ion trap for the determination of isoquinoline alkaloids. *Anal Chem* 1994; 66: 2103–2109
- 112 Unger M, Stöckigt D, Belder D, Stöckigt J. Alkaloid determination in crude extracts from cortex quebracho and opium applying capillary electrophoresis and capillary electrophoresis-mass spectrometry coupling. *Pharmazie* 1997; 52: 691–695
- 113 Stöckigt D, Unger M, Belder D, Stöckigt J. Analysis of *Rauwolfia* alkaloids employing capillary electrophoresis-mass spectrometry. *Nat Prod Lett* 1997; 9: 265–272
- 114 Sturm S, Seger C, Stuppner H. Analysis of Central European *Corydalis* species by nonaqueous capillary electrophoresis-electrospray ion trap mass spectrometry. *J Chromatogr A* 2007; 1159: 42–50
- 115 Bianco G, Schmitt-Kopplin P, De Benedetto G, Kettrup A, Cataldi TR. Determination of glycoalkaloids and relative aglycones by nonaqueous capillary electrophoresis coupled with electrospray ionization-ion trap mass spectrometry. *Electrophoresis* 2002; 23: 2904–2912
- 116 Unger M, Dreyer M, Specker S, Laug S, Pelzing M, Neuss C, Holzgrabe U, Bringmann G. Analytical characterisation of crude extracts from an African *Ancistrocladus* species using high-performance liquid chromatography and capillary electrophoresis coupled to ion trap mass spectrometry. *Phytochem Anal* 2004; 15: 21–26
- 117 Wang S, Qu H, Cheng Y. NACE-ESI-MS combined with on-line concentration for high-sensitivity analysis of quinolizidine alkaloids. *Electrophoresis* 2007; 28: 1399–1406
- 118 Arráez-Román D, Zurek G, Bässmann C, Segura-Carretero A, Fernández-Gutiérrez A. Characterisation of *Atropa belladonna* L. compounds by capillary electrophoresis-electrospray ionization-time of flight-mass spectrometry and capillary electrophoresis-electrospray ionization-ion trap-mass spectrometry. *Electrophoresis* 2008; 29: 2112–2116
- 119 Simó C, Barbas C, Cifuentes A. Capillary electrophoresis-mass spectrometry in food analysis. *Electrophoresis* 2005; 26: 1306–1318
- 120 Vanhoenacker G, DeVilliers A, Lazou K, DeKeukeleire D, Sandra P. Comparison of high-performance liquid chromatography – mass spectroscopy and capillary electrophoresis – mass spectroscopy for the analysis of phenolic compounds in diethyl ether extracts of red wines. *Chromatographia* 2001; 54: 309–315
- 121 Rodat A, Gavard P, Couderc F. Improving detection in capillary electrophoresis with laser induced fluorescence via a bubble cell capillary and laser power adjustment. *Biomed Chromatogr* 2009; 23: 42–47
- 122 Driedger DR, LeBlanc RJ, LeBlanc EL, Sporns P. A capillary electrophoresis laser-induced fluorescence method for analysis of potato glycoalkaloids based on a solution-phase immunoassay. 2. Performance evaluation. *J Agric Food Chem* 2000; 48: 4079–4082
- 123 Zhang ZL, Liu X, Li DF, Lu YT. Determination of jasmonic acid in bark extracts from *Hevea brasiliensis* by capillary electrophoresis with laser-induced fluorescence detection. *Anal Bioanal Chem* 2005; 382: 1616–1619
- 124 Wang W, Tang J, Wang S, Zhou L, Hu Z. Method development for the determination of coumarin compounds by capillary electrophoresis with indirect laser-induced fluorescence detection. *J Chromatogr A* 2007; 1148: 108–114
- 125 Geng CH, Wang WY, Lin M, Ye JN. Determination of flavonoids in *Plumula nelumbini* by micellar electrokinetic capillary electrophoresis with electrochemical detection. *J Capill Electrophor Microchip Technol* 2007; 10: 57–61
- 126 Yu LS, Xu XQ, Huang L, Ling JM, Chen GN. Separation and determination of flavonoids using microemulsion EKC with electrochemical detection. *Electrophoresis* 2008; 29: 726–733
- 127 Simpson Jr SL, Quirino JP, Terabe S. On-line sample pre-concentration in capillary electrophoresis. Fundamentals and applications. *J Chromatogr A* 2008; 1184: 504–541
- 128 Zhu J, Yu K, Chen X, Hu Z. Comparison of two sample pre-concentration strategies for the sensitivity enhancement of flavonoids found in Chinese herbal medicine in micellar electrokinetic chromatography with UV detection. *J Chromatogr A* 2007; 1166: 191–200
- 129 Yu L, Xu X, Huang L, Lin J, Chen G. Microemulsion electrokinetic chromatography coupling with field amplified sample injection and electroosmotic flow suppressant for analysis of some quinolizidine alkaloids. *J Chromatogr A* 2008; 1198: 220–225
- 130 Unger M, Stöckigt J. Improved detection of alkaloids in crude extracts applying capillary electrophoresis with field amplified sample injection. *J Chromatogr A* 1997; 791: 323–331
- 131 Petr J, Vitkocvá K, Ranc V, Znaleznia J, Maier V, Knob R, Sevcik J. Determination of some phenolic acids in *Majorana hortensis* by capillary electrophoresis with online electrokinetic pre-concentration. *J Agric Food Chem* 2008; 56: 3940–3944
- 132 Quirino JP, Otsuka K, Terabe S. On-line concentration of neutral analytes for micellar electrokinetic chromatography VI. Stacking using reverse migrating micelles and a water plug. *J Chromatogr B* 1998; 714: 29–38
- 133 Quirino JP, Terabe S. Exceeding 5000-fold concentration of dilute analytes in micellar electrokinetic chromatography. *Science* 1998; 282: 465–468
- 134 Huang HY, Hsieh SH. Analyses of tobacco alkaloids by cation-selective exhaustive injection sweeping microemulsion electrokinetic chromatography. *J Chromatogr A* 2007; 1164: 313–319
- 135 Huang HY, Huang IY, Liang HH, Lee S. Sample stacking for the analysis of catechins by microemulsion EKC. *Electrophoresis* 2007; 28: 1735–1743
- 136 Huang HY, Lien WC, Huang IY. Anion-selective exhaustive injection-sweeping microemulsion electrokinetic chromatography. *Electrophoresis* 2006; 27: 3202–3209
- 137 Hsieh MC, Lin CH. On-line identification of trans-resveratrol in red wine using a sweeping technique combined with capillary electrophoresis/77 K fluorescence spectroscopy. *Electrophoresis* 2004; 25: 677–682
- 138 Zhou M, Ma YJ, Ren XN, Zhou XY, Li L, Chen H. Determination of sinomenine in *Sinomenium acutum* by capillary electrophoresis with electrochemiluminescence detection. *Anal Chim Acta* 2007; 587: 104–109