Wire-guided endoscopic snare resection of a choledochocoele-associated villous adenoma

Fig. 1 Endoscopic view of the choledochocoele.

Adenomas arising in a choledochocoele are usually resected surgically because they are difficult to completely remove endoscopically and they are associated with a high malignancy potential [1–2]. We successfully performed an endoscopic snare resection for choledochocoele-associated villous adenoma with high-grade dysplasia.

A 76-year-old woman was admitted complaining of epigastric pain lasting for several days. Duodenoscopy revealed a soft cystic-mass-like lesion covered with normal mucosa above the ampulla of Vater (● **Fig. 1**).

Intraductal ultrasound showed an anechogenic cystic mass with diffuse wall thickening and preserved wall layer (● **Fig. 2**).

Magnetic resonance cholangiopancreatography showed an oval-shaped cystic mass above the ampulla of Vater. A major endoscopic sphincterotomy was conducted. Following sphincterotomy, duodenoscopy revealed mucosal nodules on the inner surface of the choledochocoele (● **Fig. 3**).

Histopathologic findings of the biopsy specimen showed an adenoma with high-grade dysplasia (● **Fig. 4**).

Because the patient wanted to undergo an endoscopic treatment, we performed a wire-guided endoscopic snare resection for the choledochocoele with the adenoma (● **Fig. 5**, ● **Video 1**).

Fig. 3 Endoscopic view after a major endoscopic sphincterotomy.

During the 1-year follow-up duodenoscopy, no remnant or recurrence of the adenoma was found.

The surgical approach may be too invasive for a choledochocoele with adenoma. Itoi et al. [3] performed a balloon-catheter-assisted endoscopic snare resection using a double-channel duodenoscope for choledochocoele. We attempted to perform a wire-guided endoscopic snare resection using the previously reported method for adenomas of the major papilla [4]. En bloc resection was successfully achieved without any procedure-related complications or residual tumor. In conclusion, endoscopic snare resection may be a feasible and effective solution for selected patients with choledochocoele associated with a villous adenoma.

Endoscopy_UCTN_Code_CCL_1AZ_2AK
Endoscopy_UCTN_Code_TTT_1AR_2AK

Fig. 2 Intraductal ultrasound showing an anechogenic cystic mass with diffuse wall thickening.

Video 1

Wire-guided endoscopic snare resection. The guide wire was inserted into the main pancreatic duct. An electrosurgical snare was then inserted along the guide wire. The snare was manipulated to grasp the base of the choledochocoele before en bloc resection was conducted. After the excision had been completed, a pancreatic stent was immediately placed along the indwelling guide wire. Histopathologic examination of the resected specimen showed a villous adenoma with a clear resection margin (● **Fig. 6**).

Fig. 4 Histopathologic findings of the biopsy specimen (hematoxylin and eosin stain, × 200).

Fig. 5 Endoscopic view for the wire-guided endoscopic snare resection.

Fig. 6 Histopathologic picture of the endoscopic resected specimen showing villous adenoma (hematoxylin and eosin stain, × 10).

Bibliography

DOI 10.1055/s-0029-1214429

Endoscopy 2009; 41: E78–E79

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

J. H. Moon, MD, PhD

Soon Chun Hyang University School of Medicine
Digestive Disease Center

Soon Chun Hyang University Bucheon Hospital
1174 Jung-Dong

Wonmi-Ku

Bucheon 420-767

Korea

Fax: +82-32-6215080

jhmoon@schbc.ac.kr

J. K. Park¹, J. H. Moon¹, H. K. Kim¹,
H. J. Choi¹, J. H. Kang¹, H. K. Kim²,
B. M. Ko¹, Y. D. Cho¹, M. S. Lee¹,
C. S. Shim¹

¹ Digestive Disease Center, Department
of Internal Medicine, Soon Chun Hyang
University School of Medicine, Bucheon
and Seoul, Korea

² Department of Pathology, Soon Chun
Hyang University School of Medicine,
Bucheon, Korea

References

- 1 Kawakami H, Kuwatani M, Onodera M et al. Villous adenoma arising in choledochocoele. *Gastrointest Endosc* 2007; 66: 1231–1232
- 2 Ohtsuka T, Inoue K, Ohuchida J et al. Carcinoma arising in choledochocoele. *Endoscopy* 2001; 33: 614–619
- 3 Itoi T, Gotoda T, Yasuda I et al. Balloon-catheter-assisted complete endoscopic snare resection for choledochocoele by using double-channel duodenoscope (with videos). *Gastrointest Endosc* 2007; 66: 622–625
- 4 Moon JH, Cha SW, Cho YD et al. Wire-guided endoscopic snare papillectomy for tumors of the major duodenal papilla. *Gastrointest Endosc* 2005; 61: 461–466