New endoscopic images of mucosal prolapse syndrome

Fig. 1 Colonoscopy showing a polypoid lesion with central ulceration in the lower rectum of a patient with bleeding per rectum and fecal incontinence.

Fig. 2 Dilated, brownish pits and widened pericryptal space in the region of the lesion visualized on magnifying colonoscopy.

Fig. 3 Autofluorescence imaging (AFI) findings: a magenta-colored elevation surrounds the yellowish-green ulcerated area.

Fig. 4 Microscopic examination shows elongation and distortion of the crypts and fibromuscular obliteration in the mucosa.

A 65-year-old man presented with hematochezia and fecal incontinence. For a long time he had been experiencing severe constipation and had needed to train to evacuate the bowels. Conventional colonoscopy revealed a hyperemic broad-based polypoid lesion with central ulceration in the lower rectum (● Fig. 1). Magnifying colonoscopy with narrow-band image system (NBI) revealed dilated brownish, oval-to-long pits and widening of the pericryptal space around the polypoid lesion (● Fig. 2). There was no destruction of or irregularity in the pit pattern and no abnormalities in the microvessels. Autofluorescence imaging (AFI) revealed a magenta-colored elevation surrounding the yellowish-green ulcerated area (● Fig. 3). Histological examination of biopsy specimens taken from

the polypoid lesion revealed elongation and distortion of the crypts and fibromuscular obliteration in the mucosa (● Fig. 4). On the basis of the clinicopathological features, a diagnosis of mucosal prolapse syndrome was made. The patient was successfully treated with bowel retraining to avoid straining at defecation and dependence on laxatives.

Endoscopy_UCTN_Code_CCL_1AD_2AJ

C. C. Chen¹, H. Isomoto¹, H. Ishii¹, T. Hayashi², Y. Mizuta¹, S. Kohn¹

¹ Second Department of Internal Medicine, Nagasaki University School of Medicine, Nagasaki, Japan

² Department of Pathology, Nagasaki University School of Medicine, Nagasaki, Japan

Bibliography

DOI 10.1055/s-0029-1214624

Endoscopy 2009; 41: E136

© Georg Thieme Verlag KG Stuttgart · New York · ISSN 0013-726X

Corresponding author

C. C. Chen, MD, H. Isomoto, MD

Second Department of Internal Medicine

Nagasaki University School of Medicine

1-7-1 Sakamoto

Nagasaki 852-8501

Japan

Fax: +81-95-8497285

cszen@hotmail.com

hajimei2002@yahoo.co.jp